

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
27 March 2003 (27.03.2003)

PCT

(10) International Publication Number  
**WO 03/025131 A2**

(51) International Patent Classification<sup>7</sup>: **C12N**

(21) International Application Number: **PCT/US02/29221**

(22) International Filing Date:  
13 September 2002 (13.09.2002)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:

60/322,196	14 September 2001 (14.09.2001)	US
60/324,134	21 September 2001 (21.09.2001)	US
60/327,233	5 October 2001 (05.10.2001)	US
60/346,198	26 October 2001 (26.10.2001)	US
60/343,980	2 November 2001 (02.11.2001)	US
60/348,887	9 November 2001 (09.11.2001)	US
60/332,423	16 November 2001 (16.11.2001)	US
60/334,145	28 November 2001 (28.11.2001)	US
60/334,229	28 November 2001 (28.11.2001)	US
60/337,451	6 December 2001 (06.12.2001)	US
60/351,928	25 January 2002 (25.01.2002)	US
60/366,837	21 March 2002 (21.03.2002)	US

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SPRAGUE, William, W.** [US/US]; 611 13th Street # C, Sacramento, CA 95814 (US). **CHAWLA, Narinder, K.** [US/US]; 33 Union Square, #712, Union City, CA 94587 (US). **WARREN, Bridget, A.** [US/US]; 1810 S. El Camino Real, #B103, Encinitas, CA 92024 (US). **TANG, Y., Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **ELLIOTT, Vicki, S.** [US/US]; 3770 Polton Place Way, San Jose, CA 95121 (US). **MARQUIS, Joseph, P.** [US/US]; 4428 Lazy Lane, San Jose, CA 95135 (US). **LI, Joana, X.** [US/US]; 1264 Geneva Avenue, San Francisco, CA 94112 (US). **GRIFFIN, Jennifer, A.** [US/US]; 33691 Mello Way, Fremont, CA 94555 (US). **GIETZEN, Kimberly, J.** [US/US]; 691 Los Huecos Drive, San Jose, CA 95123 (US). **YANG, Junming** [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). **LU, Dyung Aina, M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **EMERLING,**

**Brooke, M.** [US/US]; 1735 Woodland Avenue # 71, Palo Alto, CA 94303 (US). **DUGGAN, Brendan, M.** [AU/US]; 243 Buena Vista Avenue # 306, Sunnyvale, CA 94086 (US). **RICHARDSON, Thomas, W.** [US/US]; 616 Canyon Road #107, Redwood City, CA 94062 (US). **LEE, Soo, Yeun** [KR/US]; 40 Westdale Avenue, Daly City, CA 94015 (US). **RAMKUMAR, Jayalaxmi** [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). **BECHA, Shanya, D.** [US/US]; 21062 Gary Drive # 117, Castro Valley, CA 94546 (US). **LEHR-MASON, Patricia, M.** [US/US]; 360 Clarke Lane, Morgan Hill, CA 95037 (US). **SWARNAKAR, Anita** [CA/US]; 8 Locksley Avenue #5D, San Francisco, CA 94122 (US). **TRAN, Uyen, K.** [US/US]; 2638 Mabury Square, San Jose, CA 95133 (US). **KABLE, Amy, E.** [US/US]; 2345 Polk Street #4, San Francisco, CA 94109 (US). **HAFALIA, April, J.A.** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). **KHARE, Reena** [US/US]; 12650 Orella Court, Saratoga, CA 95070 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PROTEIN MODIFICATION AND MAINTENANCE MOLECULES**

(57) Abstract: Various embodiments of the invention provide human protein modification and maintenance molecules (PMMM) and polynucleotides which identify and encode PMMM. Embodiments of the invention also provide expression vectors, host cells, antibodies, agonists, and antagonists. Other embodiments provide methods for diagnosing, treating, or preventing disorders associated with aberrant expression of PMMM.a.

WO 03/025131 A2

## PROTEIN MODIFICATION AND MAINTENANCE MOLECULES

### TECHNICAL FIELD

The invention relates to novel nucleic acids, protein modification and maintenance molecules  
5 encoded by these nucleic acids, and to the use of these nucleic acids and proteins in the diagnosis,  
treatment, and prevention of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell  
proliferative, developmental, epithelial, neurological, reproductive, endocrine, metabolic, pancreatic  
disorders, disorders associated with the adrenals, disorders associated with gonadal steroid hormones,  
cancers, and infections. The invention also relates to the assessment of the effects of exogenous  
10 compounds on the expression of nucleic acids and protein modification and maintenance molecules.

### BACKGROUND OF THE INVENTION

The cellular processes regulating modification and maintenance of protein molecules  
coordinate their function, conformation, stabilization, and degradation. Each of these processes  
15 is mediated by key enzymes or proteins such as kinases, phosphatases, proteases, protease  
inhibitors, isomerases, transferases, and molecular chaperones.

#### Kinases

Kinases catalyze the transfer of high-energy phosphate groups from adenosine  
triphosphate (ATP) to target proteins on the hydroxyamino acid residues serine, threonine, or  
20 tyrosine. Addition of a phosphate group alters the local charge on the acceptor molecule, causing  
internal conformational changes and potentially influencing intermolecular contacts. Reversible  
protein phosphorylation is the ubiquitous strategy used to control many of the intracellular events  
in eukaryotic cells. It is estimated that more than ten percent of proteins active in a typical  
mammalian cell are phosphorylated. Extracellular signals including hormones,  
25 neurotransmitters, and growth and differentiation factors can activate kinases, which can occur as  
cell surface receptors or as the activators of the final effector protein, as well as elsewhere along  
the signal transduction pathway. Kinases are involved in all aspects of a cell's function, from  
basic metabolic processes, such as glycolysis, to cell-cycle regulation, differentiation, and  
communication with the extracellular environment through signal transduction cascades.  
30 Inappropriate phosphorylation of proteins in cells has been linked to changes in cell cycle  
progression and cell differentiation. Changes in the cell cycle have been linked to induction of  
apoptosis or cancer. Changes in cell differentiation have been linked to diseases and disorders of  
the reproductive system, immune system, and skeletal muscle.

There are two classes of protein kinases. One class, protein tyrosine kinases (PTKs),

phosphorylates tyrosine residues, and the other class, protein serine/threonine kinases (STKs), phosphorylates serine and threonine residues. Some PTKs and STKs possess structural characteristics of both families and have dual specificity for both tyrosine and serine/threonine residues. Almost all kinases contain a conserved 250-300 amino acid catalytic domain

- 5 containing specific residues and sequence motifs characteristic of the kinase family. (Reviewed in Hardie, G. and S. Hanks (1995) The Protein Kinase Facts Book, Vol I, Academic Press, San Diego, CA, pp. 17-20).

### Phosphatases

- Phosphatases hydrolytically remove phosphate groups from proteins. Phosphatases are  
10 essential in determining the extent of phosphorylation in the cell and, together with kinases, regulate key cellular processes such as metabolic enzyme activity, proliferation, cell growth and differentiation, cell adhesion, and cell cycle progression. Protein phosphatases are characterized as either serine/threonine- or tyrosine-specific based on their preferred phospho-amino acid substrate. Some phosphatases (DSPs, for dual specificity phosphatases) can act on  
15 phosphorylated tyrosine, serine, or threonine residues. The protein serine/threonine phosphatases (PSPs) are important regulators of many cAMP-mediated hormone responses in cells. Protein tyrosine phosphatases (PTPs) play a significant role in cell cycle and cell signaling processes.

### Proteases

- Proteases cleave proteins and peptides at the peptide bond that forms the backbone of the  
20 protein or peptide chain. Proteolysis is one of the most important and frequent enzymatic reactions that occurs both within and outside of cells. Proteolysis is responsible for the activation and maturation of nascent polypeptides, the degradation of misfolded and damaged proteins, and the controlled turnover of peptides within the cell. Proteases participate in digestion, endocrine function, tissue remodeling during embryonic development, wound healing, and normal growth.  
25 Proteases can play a role in regulatory processes by affecting the half life of regulatory proteins. Proteases are involved in the etiology or progression of disease states such as inflammation, angiogenesis, tumor dispersion and metastasis, cardiovascular disease, neurological disease, and bacterial, parasitic, and viral infections.

Proteases can be categorized on the basis of where they cleave their substrates.

- 30 Exopeptidases, which include aminopeptidases, dipeptidyl peptidases, tripeptidases, carboxypeptidases, peptidyl-di-peptidases, dipeptidases, and omega peptidases, cleave residues at the termini of their substrates. Endopeptidases, including serine proteases, cysteine proteases, and metalloproteases, cleave at residues within the peptide. Four principal categories of mammalian proteases have been identified based on active site structure, mechanism of action,

and overall three-dimensional structure. (See Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York NY, pp. 1-5.)

### Serine Proteases

The serine proteases (SPs) are a large, widespread family of proteolytic enzymes that include the digestive enzymes trypsin and chymotrypsin, components of the complement and blood-clotting cascades, and enzymes that control the degradation and turnover of macromolecules within the cell and in the extracellular matrix. Most of the more than 20 subfamilies can be grouped into six clans, each with a common ancestor. These six clans are hypothesized to have descended from at least four evolutionarily distinct ancestors. SPs are named for the presence of a serine residue found in the active catalytic site of most families. The active site is defined by the catalytic triad, a set of conserved asparagine, histidine, and serine residues critical for catalysis. These residues form a charge relay network that facilitates substrate binding. Other residues outside the active site form an oxyanion hole that stabilizes the tetrahedral transition intermediate formed during catalysis. SPs have a wide range of substrates and can be subdivided into subfamilies on the basis of their substrate specificity. The main subfamilies are named for the residue(s) after which they cleave: trypases (after arginine or lysine), aspases (after aspartate), chymases (after phenylalanine or leucine), metases (methionine), and serases (after serine) (Rawlings, N.D. and A.J. Barrett (1994) *Methods Enzymol.* 244:19-61).

Most mammalian serine proteases are synthesized as zymogens, inactive precursors that are activated by proteolysis. For example, trypsinogen is converted to its active form, trypsin, by enteropeptidase. Enteropeptidase is an intestinal protease that removes an N-terminal fragment from trypsinogen. The remaining active fragment is trypsin, which in turn activates the precursors of the other pancreatic enzymes. Likewise, proteolysis of prothrombin, the precursor of thrombin, generates three separate polypeptide fragments. The N-terminal fragment is released while the other two fragments, which comprise active thrombin, remain associated through disulfide bonds.

The two largest SP subfamilies are the chymotrypsin (S1) and subtilisin (S8) families. Some members of the chymotrypsin family contain two structural domains unique to this family. Kringle domains are triple-looped, disulfide cross-linked domains found in varying copy number. Kringle domains are thought to play a role in binding mediators such as membranes, other proteins or phospholipids, and in the regulation of proteolytic activity (PROSITE PDOC00020). Apple domains are 90 amino-acid repeated domains, each containing six conserved cysteines. Three disulfide bonds link the first and sixth, second and fifth, and third and fourth cysteines



(PROSITE PDOC00376). Apple domains are involved in protein-protein interactions. S1 family members include trypsin, chymotrypsin, coagulation factors IX-XII, complement factors B, C, and D, granzymes, kallikrein, and tissue- and urokinase-plasminogen activators. The subtilisin family has members found in the eubacteria, archaeobacteria, eukaryotes, and viruses. Subtilisins include the proprotein-processing endopeptidases kexin and furin and the pituitary prohormone convertases PC1, PC2, PC3, PC6, and PACE4 (Rawlings and Barrett, *supra*).

SPs have functions in many normal processes and some have been implicated in the etiology or treatment of disease. Enterokinase, the initiator of intestinal digestion, is found in the intestinal brush border, where it cleaves the acidic propeptide from trypsinogen to yield active trypsin (Kitamoto, Y. et al. (1994) Proc. Natl. Acad. Sci. USA 91:7588-7592). Prolylcarboxypeptidase, a lysosomal serine peptidase that cleaves peptides such as angiotensin II and III and [des-Arg<sup>9</sup>] bradykinin, shares sequence homology with members of both the serine carboxypeptidase and prolylendopeptidase families (Tan, F. et al. (1993) J. Biol. Chem. 268:16631-16638). The protease neuropsin may influence synapse formation and neuronal connectivity in the hippocampus in response to neural signaling (Chen, Z.-L. et al. (1995) J. Neurosci. 15:5088-5097). Tissue plasminogen activator is useful for acute management of stroke (Zivin, J.A. (1999) Neurology 53:14-19) and myocardial infarction (Ross, A.M. (1999) Clin. Cardiol. 22:165-171). Some receptors (PAR, for proteinase-activated receptor), highly expressed throughout the digestive tract, are activated by proteolytic cleavage of an extracellular domain. The major agonists for PARs, thrombin, trypsin, and mast cell tryptase, are released in allergy and inflammatory conditions. Control of PAR activation by proteases has been suggested as a promising therapeutic target (Vergnolle, N. (2000) Aliment. Pharmacol. Ther. 14:257-266; Rice, K.D. et al. (1998) Curr. Pharm. Des. 4:381-396). Prostate-specific antigen (PSA) is a kallikrein-like serine protease synthesized and secreted exclusively by epithelial cells in the prostate gland. Serum PSA is elevated in prostate cancer and is the most sensitive physiological marker for monitoring cancer progression and response to therapy. PSA can also identify the prostate as the origin of a metastatic tumor (Brawer, M.K. and P.H. Lange (1989) Urology 33:11-16).

The signal peptidase is a specialized class of SP found in all prokaryotic and eukaryotic cell types that serves in the processing of signal peptides from certain proteins. Signal peptides are amino-terminal domains of a protein which direct the protein from its ribosomal assembly site to a particular cellular or extracellular location. Once the protein has been exported, removal of the signal sequence by a signal peptidase and posttranslational processing, e.g., glycosylation or phosphorylation, activate the protein. Signal peptidases exist as multi-subunit complexes in both yeast and mammals. The canine signal peptidase complex is composed of five subunits, all

associated with the microsomal membrane and containing hydrophobic regions that span the membrane one or more times (Shelness, G.S. and G. Blobel (1990) J. Biol. Chem. 265:9512-9519). Some of these subunits serve to fix the complex in its proper position on the membrane while others contain the actual catalytic activity.

5       The mechanism for the translocation process into the ER involves the recognition of an N-terminal signal peptide on the elongating protein. The signal peptide directs the protein and attached ribosome to a receptor on the ER membrane. The polypeptide chain passes through a pore in the ER membrane into the lumen while the N-terminal signal peptide remains attached at the membrane surface. The process is completed when signal peptidase located inside the ER cleaves the  
10   signal peptide from the protein and releases the protein into the lumen.

Thrombin is a serine protease with an essential role in the process of blood coagulation. Prothrombin, synthesized in the liver, is converted to active thrombin by Factor Xa. Activated thrombin then cleaves soluble fibrinogen to polymer-forming fibrin, a primary component of blood clots. In addition, thrombin activates Factor XIIIa, which plays a role in cross-linking  
15   fibrin.

Thrombin also stimulates platelet aggregation through proteolytic processing of a 41-residue amino-terminal peptide from protease-activated receptor 1 (PAR-1), formerly known as the thrombin receptor. The cleavage of the amino-terminal peptide exposes a new amino terminus and may also be associated with PAR-1 internalization (Stubbs, M.T. and W. Bode  
20   (1994) Curr. Opin. Struct. Biol. 4:823-832; and Ofoso, F.A. et al. (1998) Biochem. J. 336:283-285). In addition to stimulating platelet activation through cleavage of the PAR-1 receptor, thrombin also induces platelet aggregation following cleavage of glycoprotein V, also on the surface of platelets. Glycoprotein V appears to be the major thrombin substrate on intact platelets. Platelets deficient for glycoprotein V are hypersensitive to thrombin, which is still  
25   required to cleave PAR-1. While platelet aggregation is required for normal hemostasis in mammals, excessive platelet aggregation can result in arterial thrombosis, atherosclerotic arteries, acute myocardial infarction, and stroke (Ramakrishnan, V. et al. (1999) Proc. Natl. Acad. Sci. U.S.A. 96:13336-13341 and references within).

Proteases in another family have a serine in their active site and are dependent on the  
30   hydrolysis of ATP for their activity. These proteases contain proteolytic core domains and regulatory ATPase domains which can be identified by the presence of the P-loop, an ATP/GTP-binding motif (PROSITE PDOC00803). Members of this family include the eukaryotic mitochondrial matrix proteases, Clp protease and the proteasome. Clp protease was originally found in plant chloroplasts but is believed to be widespread in both prokaryotic and eukaryotic

cells. The gene for early-onset torsion dystonia encodes a protein related to Clp protease (Ozelius, L.J. et al. (1998) Adv. Neurol. 78:93-105).

The proteasome is an intracellular protease complex found in some bacteria and in all eukaryotic cells, and plays an important role in cellular physiology. The proteasome is a large (~2000 kDa) multisubunit complex composed of a central catalytic core containing a variety of proteases arranged in four seven-membered rings with the active sites facing inwards into the central cavity, and terminal ATPase subunits covering the outer port of the cavity and regulating substrate entry (for review, see Schmidt, M. et al. (1999) Curr. Opin. Chem. Biol. 3:584-591). Proteasomes are associated with the ubiquitin conjugation system (UCS), a major pathway for the degradation of cellular proteins of all types, including proteins that function to activate or repress cellular processes such as transcription and cell cycle progression (Ciechanover, A. (1994) Cell 79:13-21). In the UCS pathway, proteins targeted for degradation are conjugated to ubiquitin, a small heat stable protein. The ubiquitinated protein is then recognized and degraded by the proteasome. The resultant ubiquitin-peptide complex is hydrolyzed by a ubiquitin carboxyl terminal hydrolase, and free ubiquitin is released for reutilization by the UCS. Ubiquitin-proteasome systems are implicated in the degradation of mitotic cyclic kinases, oncoproteins, tumor suppressor genes (p53), cell surface receptors associated with signal transduction, transcriptional regulators, and mutated or damaged proteins (Ciechanover, *supra*). This pathway has been implicated in a number of diseases, including cystic fibrosis, Angelman's syndrome, and Liddle syndrome (reviewed in Schwartz, A.L. and A. Ciechanover (1999) Annu. Rev. Med. 50:57-74). A murine proto-oncogene, Unp, encodes a nuclear ubiquitin protease whose overexpression leads to oncogenic transformation of NIH3T3 cells. The human homolog of this gene is consistently elevated in small cell tumors and adenocarcinomas of the lung (Gray, D.A. (1995) Oncogene 10:2179-2183). Ubiquitin carboxyl terminal hydrolase is involved in the differentiation of a lymphoblastic leukemia cell line to a non-dividing mature state (Maki, A. et al. (1996) Differentiation 60:59-66). In neurons, ubiquitin carboxyl terminal hydrolase (PGP 9.5) expression is strong in the abnormal structures that occur in human neurodegenerative diseases (Lowe, J. et al. (1990) J. Pathol. 161:153-160). The proteasome is a large (~2000 kDa) multisubunit complex composed of a central catalytic core containing a variety of proteases arranged in four seven-membered rings with the active sites facing inwards into the central cavity, and terminal ATPase subunits covering the outer port of the cavity and regulating substrate entry (for review, see Schmidt, M. et al. (1999) Curr. Op. Chem. Biol. 3:584-591).

#### Cysteine Proteases

Cysteine proteases (CPs) are involved in diverse cellular processes ranging from the

processing of precursor proteins to intracellular degradation. Nearly half of the CPs known are present only in viruses. CPs have a cysteine as the major catalytic residue at the active site where catalysis proceeds via a thioester intermediate and is facilitated by nearby histidine and asparagine residues. A glutamine residue is also important, as it helps to form an oxyanion hole.

- 5 Two important CP families include the papain-like enzymes (C1) and the calpains (C2). Papain-like family members are generally lysosomal or secreted and therefore are synthesized with signal peptides as well as propeptides. Most members bear a conserved motif in the propeptide that may have structural significance (Karrer, K.M. et al. (1993) Proc. Natl. Acad. Sci. USA 90:3063-3067). Three-dimensional structures of papain family members show a bilobed molecule with  
10 the catalytic site located between the two lobes. Papains include cathepsins B, C, H, L, and S, certain plant allergens and dipeptidyl peptidase (for a review, see Rawlings, N.D. and A.J. Barrett (1994) Methods Enzymol. 244:461-486).

- Some CPs are expressed ubiquitously, while others are produced only by cells of the immune system. Of particular note, CPs are produced by monocytes, macrophages and other  
15 cells which migrate to sites of inflammation and secrete molecules involved in tissue repair. Overabundance of these repair molecules plays a role in certain disorders. In autoimmune diseases such as rheumatoid arthritis, secretion of the cysteine peptidase cathepsin C degrades collagen, laminin, elastin and other structural proteins found in the extracellular matrix of bones. Bone weakened by such degradation is also more susceptible to tumor invasion and metastasis.  
20 Cathepsin L expression may also contribute to the influx of mononuclear cells which exacerbates the destruction of the rheumatoid synovium (Keyszer, G.M. (1995) Arthritis Rheum. 38:976-984).

- Calpains are calcium-dependent cytosolic endopeptidases which contain both an N-terminal catalytic domain and a C-terminal calcium-binding domain. Calpain is expressed as a  
25 proenzyme heterodimer consisting of a catalytic subunit unique to each isoform and a regulatory subunit common to different isoforms. Each subunit bears a calcium-binding EF-hand domain. The regulatory subunit also contains a hydrophobic glycine-rich domain that allows the enzyme to associate with cell membranes. Calpains are activated by increased intracellular calcium concentration, which induces a change in conformation and limited autolysis. The resultant  
30 active molecule requires a lower calcium concentration for its activity (Chan, S.L. and M.P. Mattson (1999) J. Neurosci. Res. 58:167-190). Calpain expression is predominantly neuronal, although it is present in other tissues. Several chronic neurodegenerative disorders, including ALS, Parkinson's disease and Alzheimer's disease are associated with increased calpain expression (Chan and Mattson, *supra*). Calpain-mediated breakdown of the cytoskeleton has

been proposed to contribute to brain damage resulting from head injury (McCracken, E. et al. (1999) J. Neurotrauma 16:749-761). Calpain-3 is predominantly expressed in skeletal muscle, and is responsible for limb-girdle muscular dystrophy type 2A (Minami, N. et al. (1999) J. Neurol. Sci. 171:31-37).

5 Another family of thiol proteases is the caspases, which are involved in the initiation and execution phases of apoptosis. A pro-apoptotic signal can activate initiator caspases that trigger a proteolytic caspase cascade, leading to the hydrolysis of target proteins and the classic apoptotic death of the cell. Two active site residues, a cysteine and a histidine, have been implicated in the catalytic mechanism. Caspases are among the most specific endopeptidases, cleaving after  
10 aspartate residues. Caspases are synthesized as inactive zymogens consisting of one large (p20) and one small (p10) subunit separated by a small spacer region, and a variable N-terminal prodomain. This prodomain interacts with cofactors that can positively or negatively affect apoptosis. An activating signal causes autoproteolytic cleavage of a specific aspartate residue (D297 in the caspase-1 numbering convention) and removal of the spacer and prodomain, leaving  
15 a p10/p20 heterodimer. Two of these heterodimers interact via their small subunits to form the catalytically active tetramer. The long prodomains of some caspase family members have been shown to promote dimerization and auto-processing of procaspases. Some caspases contain a "death effector domain" in their prodomain by which they can be recruited into self-activating complexes with other caspases and FADD protein associated death receptors or the TNF receptor  
20 complex. In addition, two dimers from different caspase family members can associate, changing the substrate specificity of the resultant tetramer. Endogenous caspase inhibitors (inhibitor of apoptosis proteins, or IAPs) also exist. All these interactions have clear effects on the control of apoptosis (reviewed in Chan and Mattson, *supra*; Salveson, G.S. and V.M. Dixit (1999) Proc. Natl. Acad. Sci. USA 96:10964-10967).

25 Caspases have been implicated in a number of diseases. Mice lacking some caspases have severe nervous system defects due to failed apoptosis in the neuroepithelium and suffer early lethality. Others show severe defects in the inflammatory response, as caspases are responsible for processing IL-1b and possibly other inflammatory cytokines (Chan and Mattson, *supra*). Cowpox virus and baculoviruses target caspases to avoid the death of their host cell and  
30 promote successful infection. In addition, increases in inappropriate apoptosis have been reported in AIDS, neurodegenerative diseases and ischemic injury, while a decrease in cell death is associated with cancer (Salveson and Dixit, *supra*; Thompson, C.B. (1995) Science 267:1456-1462).

#### Aspartyl proteases

Aspartyl proteases (APs) include the lysosomal proteases cathepsins D and E, as well as chymosin, renin, and the gastric pepsins. Most retroviruses encode an AP, usually as part of the *pol* polyprotein. APs, also called acid proteases, are monomeric enzymes consisting of two domains, each domain containing one half of the active site with its own catalytic aspartic acid residue. APs are most active in the range of pH 2–3, at which one of the aspartate residues is ionized and the other neutral. The pepsin family of APs contains many secreted enzymes, and all are likely to be synthesized with signal peptides and propeptides. Most family members have three disulfide loops, the first ~5 residue loop following the first aspartate, the second 5–6 residue loop preceding the second aspartate, and the third and largest loop occurring toward the C terminus. Retropepsins, on the other hand, are analogous to a single domain of pepsin, and become active as homodimers with each retropepsin monomer contributing one half of the active site. Retropepsins are required for processing the viral polyproteins.

APs have roles in various tissues, and some have been associated with disease. Renin mediates the first step in processing the hormone angiotensin, which is responsible for regulating electrolyte balance and blood pressure (reviewed in Crews, D.E. and S.R. Williams (1999) *Hum. Biol.* 71:475–503). Abnormal regulation and expression of cathepsins are evident in various inflammatory disease states. Expression of cathepsin D is elevated in synovial tissues from patients with rheumatoid arthritis and osteoarthritis. The increased expression and differential regulation of the cathepsins are linked to the metastatic potential of a variety of cancers (Chambers, A.F. et al. (1993) *Crit. Rev. Oncol.* 4:95–114).

#### Metalloproteases

Metalloproteases require a metal ion for activity, usually manganese or zinc. Examples of manganese metalloenzymes include aminopeptidase P and human proline dipeptidase (PEPD). Aminopeptidase P can degrade bradykinin, a nonapeptide activated in a variety of inflammatory responses. Aminopeptidase P has been implicated in coronary ischemia/reperfusion injury. Administration of aminopeptidase P inhibitors has been shown to have a cardioprotective effect in rats (Ersahin, C. et al (1999) *J. Cardiovasc. Pharmacol.* 34:604–611).

Most zinc-dependent metalloproteases share a common sequence in the zinc-binding domain. The active site is made up of two histidines which act as zinc ligands and a catalytic glutamic acid C-terminal to the first histidine. Proteins containing this signature sequence are known as the metzincins and include aminopeptidase N, angiotensin-converting enzyme, neurolysin, the matrix metalloproteases and the adamalysins (ADAMS). An alternate sequence is found in the zinc carboxypeptidases, in which all three conserved residues – two histidines and a glutamic acid – are involved in zinc binding.

A number of the neutral metalloendopeptidases, including angiotensin converting enzyme and the aminopeptidases, are involved in the metabolism of peptide hormones. High aminopeptidase B activity, for example, is found in the adrenal glands and neurohypophyses of hypertensive rats (Prieto, I. et al. (1998) *Horm. Metab. Res.* 30:246-248). Oligopeptidase M/neurolysin can hydrolyze bradykinin as well as neurotensin (Serizawa, A. et al. (1995) *J. Biol. Chem.* 270:2092-2098). Neurotensin is a vasoactive peptide that can act as a neurotransmitter in the brain, where it has been implicated in limiting food intake (Tritos, N.A. et al. (1999) *Neuropeptides* 33:339-349).

The matrix metalloproteases (MMPs) are a family of at least 23 enzymes that can degrade components of the extracellular matrix (ECM). They are  $Zn^{2+}$  endopeptidases with an N-terminal catalytic domain. Nearly all members of the family have a hinge peptide and a C-terminal domain which can bind to substrate molecules in the ECM or to inhibitors produced by the tissue (TIMPs, for tissue inhibitor of metalloprotease; Campbell, I.L. and A. Pagenstecher (1999) *Trends Neurosci.* 22:285-287). The presence of fibronectin-like repeats, transmembrane domains, or C-terminal hemopexinase-like domains can be used to separate MMPs into collagenase, gelatinase, stromelysin and membrane-type MMP subfamilies. In the inactive form, the  $Zn^{2+}$  ion in the active site interacts with a cysteine in the pro-sequence. Activating factors disrupt the  $Zn^{2+}$ -cysteine interaction, or "cysteine switch," exposing the active site. This partially activates the enzyme, which then cleaves off its propeptide and becomes fully active. MMPs are often activated by the serine proteases plasmin and furin. MMPs are often regulated by stoichiometric, noncovalent interactions with inhibitors; the balance of protease to inhibitor, then, is very important in tissue homeostasis (reviewed in Yong, V.W. et al. (1998) *Trends Neurosci.* 21:75-80).

MMPs are implicated in a number of diseases including osteoarthritis (Mitchell, P. et al. (1996) *J. Clin. Invest.* 97:761-768), atherosclerotic plaque rupture (Sukhova, G.K. et al. (1999) *Circulation* 99:2503-2509), aortic aneurysm (Schneiderman, J. et al. (1998) *Am. J. Path.* 152:703-710), non-healing wounds (Saarialho-Kere, U.K. et al. (1994) *J. Clin. Invest.* 94:79-88), bone resorption (Blavier, L. and J.M. Delaisse (1995) *J. Cell Sci.* 108:3649-3659), age-related macular degeneration (Steen, B. et al. (1998) *Invest. Ophthalmol. Vis. Sci.* 39:2194-2200), emphysema (Finlay, G.A. et al. (1997) *Thorax* 52:502-506), myocardial infarction (Rohde, L.E. et al. (1999) *Circulation* 99:3063-3070) and dilated cardiomyopathy (Thomas, C.V. et al. (1998) *Circulation* 97:1708-1715). MMP inhibitors prevent metastasis of mammary carcinoma and experimental tumors in rat, and Lewis lung carcinoma, hemangioma, and human ovarian carcinoma xenografts in mice (Eccles, S.A. et al. (1996) *Cancer Res.* 56:2815-2822; Anderson et al. (1996) *Cancer Res.*

56:715-718; Volpert, O.V. et al. (1996) J. Clin. Invest. 98:671-679; Taraboletti, G. et al. (1995) J. Natl. Cancer Inst. 87:293-298; Davies, B. et al. (1993) Cancer Res. 53:2087-2091). MMPs may be active in Alzheimer's disease. A number of MMPs are implicated in multiple sclerosis, and administration of MMP inhibitors can relieve some of its symptoms (reviewed in Yong et al.,  
 5 *supra*).

The astacin family of metalloendopeptidases have been detected in species ranging from hydra to humans, in mature and in developmental systems, performing functions involved in activation of growth factors, degradation of polypeptides, and processing of extracellular proteins. Astacin family proteases are synthesized with NH<sub>2</sub>-terminal signal and proenzyme  
 10 sequences, and many (such as meprins, BMP-1, tolloid) contain multiple domains COOH-terminal to the protease domain. They may be secreted from cells or are plasma membrane-associated enzymes. They have a signature sequence in the protease domain and a unique type of zinc binding, with pentacoordination, as well as a protease domain tertiary structure that contains common attributes with serralsins, matrix metalloendopeptidases, and  
 15 snake venom proteases. Astacins cleave peptide bonds in polypeptides such as insulin B chain and bradykinin and in proteins such as casein and gelatin; and they have arylamidase activity. Meprins are unique proteases in the astacin family, due to their oligomeric structure; they are dimers of disulfide-linked dimers and are highly glycosylated, type I integral membrane proteins that have many attributes of receptors or integrins with adhesion, epidermal growth factor-like,  
 20 and transmembrane domains. The alpha and beta subunits are differentially expressed and processed to yield latent and active proteases as well as membrane-associated and secreted forms. Meprins are regulated at the transcriptional and posttranslational levels (Bond, J.S. and Beynon, R.J. (1995) Protein Sci. 4:1247-1261).

Another family of metalloproteases is the ADAMs, for A Disintegrin and Metalloprotease  
 25 Domain, which they share with their close relatives the adamalysins, snake venom metalloproteases (SVMPs). ADAMs combine features of both cell surface adhesion molecules and proteases, containing a prodomain, a protease domain, a disintegrin domain, a cysteine rich domain, an epidermal growth factor repeat, a transmembrane domain, and a cytoplasmic tail. The first three domains listed above are also found in the SVMPs. The ADAMs possess four  
 30 potential functions: proteolysis, adhesion, signaling and fusion. The ADAMs share the metzincin zinc binding sequence and are inhibited by some MMP antagonists such as TIMP-1.

ADAMs are implicated in such processes as sperm-egg binding and fusion, myoblast fusion, and protein-ectodomain processing or shedding of cytokines, cytokine receptors, adhesion proteins and other extracellular protein domains (Schlöndorff, J. and C.P. Blobel (1999) J. Cell.



Sci. 112:3603-3617). The Kuzbanian protein cleaves a substrate in the NOTCH pathway (possibly NOTCH itself), activating the program for lateral inhibition in *Drosophila* neural development. Two ADAMs, TACE (ADAM 17) and ADAM 10, are proposed to have analogous roles in the processing of amyloid precursor protein in the brain (Schlöndorff and Blobel, *supra*).

- 5 TACE has also been identified as the TNF activating enzyme (Black, R.A. et al. (1997) Nature 385:729-733). TNF is a pleiotropic cytokine that is important in mobilizing host defenses in response to infection or trauma, but can cause severe damage in excess and is often overproduced in autoimmune disease. TACE cleaves membrane-bound pro-TNF to release a soluble form. Other ADAMs may be involved in a similar type of processing of other membrane-bound
- 10 molecules.

- Proteins of the ADAMTS sub-family have all of the features of ADAM family metalloproteases and contain an additional thrombospondin domain (TS). The prototypic ADAMTS was identified in mouse, and found to be expressed in heart and kidney and upregulated by proinflammatory stimuli (Kuno, K. et al. (1997) J. Biol. Chem. 272:556-562). To
- 15 date eleven members are recognized by the Human Genome Organization (HUGO; <http://www.gene.ucl.ac.uk/users/hester/adamts.html#Approved>). Members of this family have the ability to degrade aggrecan, a high molecular weight proteoglycan which provides cartilage with important mechanical properties including compressibility, and which is lost during the development of arthritis. Enzymes which degrade aggrecan are thus considered attractive targets
- 20 to prevent and slow the degradation of articular cartilage (See, e.g., Tortorella, M.D. (1999) Science 284:1664-1666; Abbaszade, I. (1999) J. Biol. Chem. 274:23443-23450). Other members are reported to have antiangiogenic potential (Kuno et al., *supra*) and/or procollagen processing (Colige, A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2374-2379).

#### Protease inhibitors

- 25 Protease inhibitors and other regulators of protease activity control the activity and effects of proteases. Protease inhibitors have been shown to control pathogenesis in animal models of proteolytic disorders (Murphy, G. (1991) Agents Actions Suppl. 35:69-76). In patients with HIV disease protease inhibitors have been shown to be effective in preventing disease progression and reducing mortality (Barry, M. et al. (1997) Clin. Pharmacokinet. 32:194-209). Low levels of the
- 30 cystatins, low molecular weight inhibitors of the cysteine proteases, correlate with malignant progression of tumors (Calkins, C. et al. (1995) Biol. Biochem. Hoppe Seyler 376:71-80). The cystatin superfamily of protease inhibitors is characterized by a particular pattern of linearly arranged and tandemly repeated disulfide loops (Kellermann, J. et al. (1989) J. Biol. Chem. 264:14121-14128). An example of a representative of a structural prototype of a novel family

among the cystatin superfamily is human alpha 2-HS glycoprotein (AHSG), a plasma protein synthesized in liver and selectively concentrated in bone matrix, dentine, and other mineralized tissues (Triffitt, J.T. (1976) *Calcif. Tissue Res.* 22:27-33), which is also classified as belonging to the fetuin family. Fetuins are characterized by the presence of 2 N-terminally located cystatin-  
 5 like repeats and a unique C-terminal domain which is not present in other proteins of the cystatin superfamily (PROSITE PDOC00966). AHSG has been reported to be involved in bone formation and resorption as well as immune responses (Yang, F. et al. (1992) 1130:149-156; Lee, C.C. et al. (1987) *PNAS USA* 84:4403-4407; Nakamura, O. et al. (1999) *Biosci. Biotechnol. Biochem.* 63:1383-1391). Additionally, AHSG has been implicated in infertility associated with  
 10 endometriosis (Mathur, S.P. (2000) *Am. J. Reprod. Immunol.* 44:89-95; Mathur, S.P. et al. (1999) *Autoimmunity* 29:121-127) and inhibition of osteogenesis (Binkert, C. et al, (1999) *J. Biol Chem.* 274:28514-28520). Decreased serum levels of AHSG have been detected in patients with acute leukemias, chronic granulocyte and myelomonocyte leukemias, lymphomas, myelofibrosis, multiple myeloma, metastatizing solid tumors, systemic lupus erythematosus, rheumatoid  
 15 arthritis, acute alcoholic hepatitis, fatty liver, chronic active hepatitis, liver cirrhosis, acute and chronic pancreatitis, and Crohn's disease (Kalabay, L. et al. (1992) *Orv. Hetil.* 133:1553-1554; 1559-1560).

Serpins are inhibitors of mammalian plasma serine proteases. Many serpins serve to regulate the blood clotting cascade and/or the complement cascade in mammals. Sp32 is a  
 20 positive regulator of the mammalian acrosomal protease, acrosin, that binds the proenzyme, proacrosin, and thereby aides in packaging the enzyme into the acrosomal matrix (Baba, T. et al. (1994) *J. Biol. Chem.* 269:10133-10140). The Kunitz family of serine protease inhibitors are characterized by one or more "Kunitz domains" containing a series of cysteine residues that are regularly spaced over approximately 50 amino acid residues and form three intrachain disulfide  
 25 bonds. Members of this family include aprotinin, tissue factor pathway inhibitor (TFPI-1 and TFPI-2), inter-a-trypsin inhibitor, and bikunin (Marlor, C.W. et al. (1997) *J. Biol. Chem.* 272:12202-12208). Members of this family are potent inhibitors (in the nanomolar range) against serine proteases such as kallikrein and plasmin. Aprotinin has clinical utility in reduction of perioperative blood loss. ITI has been found to inactivate human trypsin, chymotrypsin,  
 30 neutrophil elastase and cathepsin G (Morii, M. et al. (1985) *Biol. Chem. Hoppe Seyler* 366:19-21); and is suspected of playing a key role in the biology of the extracellular matrix and in the pathophysiology of chronic bronchopulmonary diseases or lung cancer progression (Cuvelier, A. et al. (2000) *Rev. Mal. Respir.* 17:437-446).

A major portion of all proteins synthesized in eukaryotic cells are synthesized on the cytosolic surface of the endoplasmic reticulum (ER). Before these immature proteins are distributed to other organelles in the cell or are secreted, they must be transported into the interior lumen of the ER where post-translational modifications are performed. These modifications include protein folding and the formation of disulfide bonds, and N-linked glycosylations.

#### **Protein Isomerases**

Protein folding in the ER is aided by two principal types of protein isomerases, protein disulfide isomerase (PDI), and peptidyl-prolyl isomerase (PPI). PDI catalyzes the oxidation of free sulfhydryl groups in cysteine residues to form intramolecular disulfide bonds in proteins.

PPI, an enzyme that catalyzes the isomerization of certain proline imidic bonds in oligopeptides and proteins, is considered to govern one of the rate limiting steps in the folding of many proteins to their final functional conformation. The cyclophilins represent a major class of PPI that was originally identified as the major receptor for the immunosuppressive drug cyclosporin A (Handschumacher, R.E. et al. (1984) *Science* 226: 544-547).

#### **Protein Glycosylation**

The glycosylation of most soluble secreted and membrane-bound proteins by oligosaccharides linked to asparagine residues in proteins is also performed in the ER. This reaction is catalyzed by a membrane-bound enzyme, oligosaccharyl transferase. Although the exact purpose of this "N-linked" glycosylation is unknown, the presence of oligosaccharides tends to make a glycoprotein resistant to protease digestion. In addition, oligosaccharides attached to cell-surface proteins called selectins are known to function in cell-cell adhesion processes (Alberts, B. et al. (1994) Molecular Biology of the Cell Garland Publishing Co., New York, NY, p. 608). "O-linked" glycosylation of proteins also occurs in the ER by the addition of N-acetylgalactosamine to the hydroxyl group of a serine or threonine residue followed by the sequential addition of other sugar residues to the first. This process is catalyzed by a series of glycosyltransferases, each specific for a particular donor sugar nucleotide and acceptor molecule (Lodish, H. et al. (1995) Molecular Cell Biology, W. H. Freeman and Co., New York, NY, pp. 700-708). For example, one of the glycosyltransferases in the dolichol pathway, dolichol phosphate mannosyl synthase, is required in N-glycosylation, O-mannosylation, and glycosylphosphatidylinositol membrane anchoring of protein (Tomita, S. et al. (1998) *J. Biol. Chem.* 9249-9254). Thus, in many cases, both N- and O-linked oligosaccharides appear to be required for the secretion of proteins or the movement of plasma membrane glycoproteins to the cell surface.

An additional glycosylation mechanism operates in the ER specifically to target lysosomal enzymes to lysosomes and prevent their secretion. Lysosomal enzymes in the ER receive an N-linked oligosaccharide, like plasma membrane and secreted proteins, but are then phosphorylated on one or two mannose residues. The phosphorylation of mannose residues occurs in two steps, the first step being the addition of an N-acetylglucosamine phosphate residue by N-acetylglucosamine phosphotransferase, and the second the removal of the N-acetylglucosamine group by phosphodiesterase. The phosphorylated mannose residue then targets the lysosomal enzyme to a mannose 6-phosphate receptor which transports it to a lysosome vesicle (Lodish et al. *supra*, pp. 708-711).

## 10 **Chaperones**

Molecular chaperones are proteins that aid in the proper folding of immature proteins and refolding of improperly folded ones, the assembly of protein subunits, and in the transport of unfolded proteins across membranes. Chaperones are also called heat-shock proteins (hsp) because of their tendency to be expressed in dramatically increased amounts following brief exposure of cells to elevated temperatures. This latter property most likely reflects their need in the refolding of proteins that have become denatured by the high temperatures. Chaperones may be divided into several classes according to their location, function, and molecular weight, and include hsp60, TCP1, hsp70, hsp40 (also called DnaJ), and hsp90. For example, hsp90 binds to steroid hormone receptors, represses transcription in the absence of the ligand, and provides proper folding of the ligand-binding domain of the receptor in the presence of the hormone (Burston, S.G. and A.R. Clarke (1995) *Essays Biochem.* 29:125-136). Hsp60 and hsp70 chaperones aid in the transport and folding of newly synthesized proteins. Hsp70 acts early in protein folding, binding a newly synthesized protein before it leaves the ribosome and transporting the protein to the mitochondria or ER before releasing the folded protein. Hsp60, along with hsp10, binds misfolded proteins and gives them the opportunity to refold correctly. All chaperones share an affinity for hydrophobic patches on incompletely folded proteins and the ability to hydrolyze ATP. The energy of ATP hydrolysis is used to release the hsp-bound protein in its properly folded state (Alberts et al., *supra*, pp. 214, 571-572). Hsp40/DnaJ homologs include mDj3, mDj4, mDj5, mDj6, mDj7, mDj8, mDj9, mDj10, and mDj11 (Ohtsuka, K. and Hata, M. (2000) *Cell Stress Chaperones* 5:98-112).

## Lysyl Hydroxylases

Lysyl hydroxylase is an enzyme involved in collagen biosynthesis. Collagens are a family of fibrous structural proteins that are found in essentially all tissues. Collagens are the most abundant proteins in mammals, and are essential for the formation of connective tissue such as

skin, bone, tendon, cartilage, blood vessels and teeth. Members of the collagen family can be distinguished from one another by the degree of cross-linking between collagen fibers and by the number of carbohydrate units (e.g., galactose or glucosylgalactose) attached to the collagen fibers. Hydroxylated lysine residues (hydroxylysine) are essential for stability of cross-linking and as attachment points for carbohydrate units.

The enzyme lysyl hydroxylase catalyzes the hydroxylation of lysine residues to form hydroxylysine. Lysyl hydroxylase targets the lysine residue of the sequence, X-lys-gly (lys = lysine, gly = glycine, and X = any amino acid residue). Three isoforms of lysyl hydroxylase have been characterized, termed LH1 (or PLOD; procollagen-lysine, 2-oxoglutarate 5-dioxygenase), LH2 (or PLOD2), and LH3. The three enzymes share 60% sequence identity overall, with even higher similarity in the C-terminal region. In addition, there are regions in the middle of the molecule that have an identity of more than 80% (Valtavaara, M. et al. (1998) J. Biol. Chem. 273:12881-12886).

Diminished lysyl hydroxylase activity is involved in certain connective tissue disorders. In particular mutations, including a truncation and duplications within the coding region of the gene for PLOD, have been described in patients with type VI Ehlers-Danos syndrome (Hyland, J. et al. (1992) Nature Genet. 2:228-31; Hautala, T. et al. (1993) Genomics 15:399-404).

#### Expression profiling

Microarrays are analytical tools used in bioanalysis. A microarray has a plurality of molecules spatially distributed over, and stably associated with, the surface of a solid support. Microarrays of polypeptides, polynucleotides, and/or antibodies have been developed and find use in a variety of applications, such as gene sequencing, monitoring gene expression, gene mapping, bacterial identification, drug discovery, and combinatorial chemistry.

One area in particular in which microarrays find use is in gene expression analysis. Array technology can provide a simple way to explore the expression of a single polymorphic gene or the expression profile of a large number of related or unrelated genes. When the expression of a single gene is examined, arrays are employed to detect the expression of a specific gene or its variants. When an expression profile is examined, arrays provide a platform for identifying genes that are tissue specific, are affected by a substance being tested in a toxicology assay, are part of a signaling cascade, carry out housekeeping functions, or are specifically related to a particular

#### Alzheimer's Disease

The potential application of gene expression profiling is also relevant to improving diagnosis, prognosis, and treatment of diseases such as Alzheimer's disease. For example, both the levels and sequences expressed in tissues from subjects with Alzheimer's disease may be compared with the

levels and sequences expressed in normal brain tissue. Alzheimer's disease is a progressive neurodegenerative disorder that is characterized by the formation of senile plaques and neurofibrillary tangles containing amyloid beta peptide. These plaques are found in limbic and association cortices of the brain. The hippocampus is part of the limbic system and plays an important role in learning and memory. In subjects with Alzheimer's disease, accumulating plaques damage the neuronal architecture in limbic areas and eventually cripple the memory process.

#### Steroid Hormones

The potential application of gene expression profiling is relevant to measuring the toxic response to potential therapeutic compounds and of the metabolic response to therapeutic agents. For instance, diseases treated with steroids and disorders caused by the metabolic response to treatment with steroids include adenomatosis, cholestasis, cirrhosis, hemangioma, Henoch-Schonlein purpura, hepatitis, hepatocellular and metastatic carcinomas, idiopathic thrombocytopenic purpura, porphyria, sarcoidosis, and Wilson disease. It is desirable to measure the toxic response to potential therapeutic compounds and of the metabolic response to therapeutic agents.

Steroids are a class of lipid-soluble molecules, including cholesterol, bile acids, vitamin D, and hormones, that share a common four-ring structure based on cyclopentanoperhydrophenanthrene and that carry out a wide variety of functions. Steroid hormones, produced by the adrenal cortex, ovaries, and testes, include glucocorticoids, mineralocorticoids, androgens, and estrogens. Steroid hormones are widely used for fertility control and in anti-inflammatory treatments for physical injuries and diseases such as arthritis, asthma, and auto-immune disorders. Progesterone, a naturally occurring progestin, is primarily used to treat amenorrhea, abnormal uterine bleeding, or as a contraceptive. Medroxyprogesterone (MAH), also known as 6 $\alpha$ -methyl-17-hydroxyprogesterone, is a synthetic progestin with a pharmacological activity about 15 times greater than progesterone. MAH is used for the treatment of renal and endometrial carcinomas, amenorrhea, abnormal uterine bleeding, and endometriosis associated with hormonal imbalance. MAH has a stimulatory effect on respiratory centers and has been used in cases of low blood oxygenation caused by sleep apnea, chronic obstructive pulmonary disease, or hypercapnia. Beclomethasone is a synthetic glucocorticoid that is used to treat steroid-dependent asthma, to relieve symptoms associated with allergic or nonallergic (vasomotor) rhinitis, or to prevent recurrent nasal polyps following surgical removal. Budesonide is a corticosteroid used to control symptoms associated with allergic rhinitis or asthma. Dexamethasone is a synthetic glucocorticoid used in anti-inflammatory or immunosuppressive compositions. Prednisone is metabolized in the liver to its active form, prednisolone, a glucocorticoid with anti-inflammatory properties. Betamethasone is a synthetic glucocorticoid with anti-inflammatory and immunosuppressive activity and is used to treat psoriasis and fungal infections, such as athlete's foot and ringworm. By comparing both the levels and sequences expressed in tissues

from subjects exposed to or treated with steroid compounds with the levels and sequences expressed in normal untreated tissue it is possible to determine tissue responses to steroids.

### Breast Cancer

Array technology can provide a simple way to explore the expression of a single polymorphic gene or the expression profile of a large number of related or unrelated genes. When the expression of a single gene is examined, arrays are employed to detect the expression of a specific gene or its variants. When an expression profile is examined, arrays provide a platform for identifying genes that are tissue specific, are affected by a substance being tested in a toxicology assay, are part of a signaling cascade, carry out housekeeping functions, or are specifically related to a particular genetic predisposition, condition, disease, or disorder.

The potential application of gene expression profiling is particularly relevant to improving diagnosis, prognosis, and treatment of cancers, such as breast cancer, colon cancer, lung cancer, ovarian cancer and prostate cancer. Breast cancer is the most frequently diagnosed type of cancer in American women and the second most frequent cause of cancer death. The lifetime risk of an American woman developing breast cancer is 1 in 8, and one-third of women diagnosed with breast cancer die of the disease. A number of risk factors have been identified, including hormonal and genetic factors. One genetic defect associated with breast cancer results in a loss of heterozygosity (LOH) at multiple loci such as p53, Rb, BRCA1, and BRCA2. Another genetic defect is gene amplification involving genes such as c-myc and c-erbB2 (Her2-neu gene). Steroid and growth factor pathways are also altered in breast cancer, notably the estrogen, progesterone, and epidermal growth factor (EGF) pathways. Breast cancer evolves through a multi-step process whereby premalignant mammary epithelial cells undergo a relatively defined sequence of events leading to tumor formation. An early event in tumor development is ductal hyperplasia. Cells undergoing rapid neoplastic growth gradually progress to invasive carcinoma and become metastatic to the lung, bone, and potentially other organs. Variables that may influence the process of tumor progression and malignant transformation include genetic factors, environmental factors, growth factors, and hormones.

### Colon Cancer

Colon cancer evolves through a multi-step process whereby pre-malignant colonocytes undergo a relatively defined sequence of events leading to tumor formation. While soft tissue sarcomas are relatively rare, more than 50% of new patients diagnosed with the disease will die from it. The molecular pathways leading to the development of sarcomas are relatively unknown, due to the rarity of the disease and variation in pathology. Several factors participate in the process of tumor progression and malignant transformation including genetic factors, mutations, and selection.

To understand the nature of gene alterations in colorectal cancer, a number of studies have focused on the inherited syndromes. Familial adenomatous polyposis (FAP), is caused by mutations

in the adenomatous polyposis coli gene (APC), resulting in truncated or inactive forms of the protein. This tumor suppressor gene has been mapped to chromosome 5q. Hereditary nonpolyposis colorectal cancer (HNPCC) is caused by mutations in mismatch repair genes. Although hereditary colon cancer syndromes occur in a small percentage of the population and most colorectal cancers are considered sporadic, knowledge from studies of the hereditary syndromes can be generally applied. For instance, somatic mutations in APC occur in at least 80% of sporadic colon tumors. APC mutations are thought to be the initiating event in the disease. Other mutations occur subsequently. Approximately 50% of colorectal cancers contain activating mutations in ras, while 85% contain inactivating mutations in p53. Changes in all of these genes lead to gene expression changes in colon cancer.

#### 10 Lung Cancer

Lung cancer is the leading cause of cancer death in the United States, affecting more than 100,000 men and 50,000 women each year. Nearly 90% of the patients diagnosed with lung cancer are cigarette smokers. Tobacco smoke contains thousands of noxious substances that induce carcinogen metabolizing enzymes and covalent DNA adduct formation in the exposed bronchial epithelium. In nearly 80% of patients diagnosed with lung cancer, metastasis has already occurred. Most commonly lung cancers metastasize to pleura, brain, bone, pericardium, and liver. The decision to treat with surgery, radiation therapy, or chemotherapy is made on the basis of tumor histology, response to growth factors or hormones, and sensitivity to inhibitors or drugs. With current treatments, most patients die within one year of diagnosis. Earlier diagnosis and a systematic approach to identification, staging, and treatment of lung cancer could positively affect patient outcome.

Lung cancers progress through a series of morphologically distinct stages from hyperplasia to invasive carcinoma. Malignant lung cancers are divided into two groups comprising four histopathological classes. The Non Small Cell Lung Carcinoma (NSCLC) group includes squamous cell carcinomas, adenocarcinomas, and large cell carcinomas and accounts for about 70% of all lung cancer cases. Adenocarcinomas typically arise in the peripheral airways and often form mucin secreting glands. Squamous cell carcinomas typically arise in proximal airways. The histogenesis of squamous cell carcinomas may be related to chronic inflammation and injury to the bronchial epithelium, leading to squamous metaplasia. The Small Cell Lung Carcinoma (SCLC) group accounts for about 20% of lung cancer cases. SCLCs typically arise in proximal airways and exhibit a number of paraneoplastic syndromes including inappropriate production of adrenocorticotropin and anti-diuretic hormone.

Lung cancer cells accumulate numerous genetic lesions, many of which are associated with cytologically visible chromosomal aberrations. The high frequency of chromosomal deletions associated with lung cancer may reflect the role of multiple tumor suppressor loci in the etiology of



this disease. Deletion of the short arm of chromosome 3 is found in over 90% of cases and represents one of the earliest genetic lesions leading to lung cancer. Deletions at chromosome arms 9p and 17p are also common. Other frequently observed genetic lesions include overexpression of telomerase, activation of oncogenes such as K-ras and c-myc, and inactivation of tumor suppressor genes such as RB, p53 and CDKN2.

Genes differentially regulated in lung cancer have been identified by a variety of methods. Using mRNA differential display technology, Manda *et al.* (1999; Genomics 51:5-14) identified five genes differentially expressed in lung cancer cell lines compared to normal bronchial epithelial cells. Among the known genes, pulmonary surfactant apoprotein A and alpha 2 macroglobulin were down regulated whereas nm23H1 was upregulated. Petersen *et al.* (2000; Int J. Cancer, 86:512-517) used suppression subtractive hybridization to identify 552 clones differentially expressed in lung tumor derived cell lines, 205 of which represented known genes. Among the known genes, thrombospondin-1, fibronectin, intercellular adhesion molecule 1, and cytokeratins 6 and 18 were previously observed to be differentially expressed in lung cancers. Wang *et al.* (2000; Oncogene 19:1519-1528) used a combination of microarray analysis and subtractive hybridization to identify 17 genes differentially overexpressed in squamous cell carcinoma compared with normal lung epithelium. Among the known genes they identified were keratin isoform 6, KOC, SPRC, IGFb2, connexin 26, plakophilin 1 and cytokeratin 13.

#### Ovarian Cancer

Ovarian cancer is the leading cause of death from a gynecologic cancer. The majority of ovarian cancers are derived from epithelial cells, and 70% of patients with epithelial ovarian cancers present with late-stage disease. As a result, the long-term survival rates for this disease is very low. Identification of early-stage markers for ovarian cancer would significantly increase the survival rate. Genetic variations involved in ovarian cancer development include mutation of p53 and microsatellite instability. Gene expression patterns likely vary when normal ovary is compared to ovarian tumors.

#### Prostate Cancer

Prostate cancer is a common malignancy in men over the age of 50, and the incidence increases with age. In the US, there are approximately 132,000 newly diagnosed cases of prostate cancer and more than 33,000 deaths from the disorder each year. Once cancer cells arise in the prostate, they are stimulated by testosterone to a more rapid growth. Thus, removal of the testes can indirectly reduce both rapid growth and metastasis of the cancer. Over 95 percent of prostatic cancers are adenocarcinomas which originate in the prostatic acini. The remaining 5 percent are divided between squamous cell and transitional cell carcinomas, both of which arise in the prostatic ducts or other parts of the prostate gland.

As with most tumors, prostate cancer develops through a multistage progression ultimately resulting in an aggressive tumor phenotype. The initial step in tumor progression involves the hyperproliferation of normal luminal and/or basal epithelial cells. Androgen responsive cells become hyperplastic and evolve into early-stage tumors. Although early-stage tumors are often androgen sensitive and respond to androgen ablation, a population of androgen independent cells evolve from the hyperplastic population. These cells represent a more advanced form of prostate tumor that may become invasive and potentially become metastatic to the bone, brain, or lung. A variety of genes may be differentially expressed during tumor progression. For example, loss of heterozygosity (LOH) is frequently observed on chromosome 8p in prostate cancer. Fluorescence in situ hybridization (FISH) revealed a deletion for at least 1 locus on 8p in 29 (69%) tumors, with a significantly higher frequency of the deletion on 8p21.2-p21.1 in advanced prostate cancer than in localized prostate cancer, implying that deletions on 8p22-p21.3 play an important role in tumor differentiation, while 8p21.2-p21.1 deletion plays a role in progression of prostate cancer (Oba, K. et al. (2001) Cancer Genet. Cytogenet. 124: 20-26).

A primary diagnostic marker for prostate cancer is prostate specific antigen (PSA). PSA is a tissue-specific serine protease almost exclusively produced by prostatic epithelial cells. The quantity of PSA correlates with the number and volume of the prostatic epithelial cells, and consequently, the levels of PSA are an excellent indicator of abnormal prostate growth. Men with prostate cancer exhibit an early linear increase in PSA levels followed by an exponential increase prior to diagnosis. However, since PSA levels are also influenced by factors such as inflammation, androgen and other growth factors, some scientists maintain that changes in PSA levels are not useful in detecting individual cases of prostate cancer.

#### Leukocytes

Leukocytes comprise lymphocytes, granulocytes, and monocytes. Lymphocytes include T- and B-cells, which specifically recognize and respond to foreign pathogens. T-cells fight viral infections and activate other leukocytes, while B-cells secrete antibodies that neutralize bacteria and other microbes. Granulocytes and monocytes are primarily migratory, phagocytic cells that exit the bloodstream to fight infection in tissues. Monocytes, which are derived from immature promonocytes, further differentiate into macrophages that engulf and digest microorganisms and damaged or dead cells. Monocytes and macrophages modulate the immune response by secreting signaling molecules such as growth factors and cytokines. Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), for example, is a macrophage-secreted protein with anti-tumor and anti-viral activity. In addition, monocytes and macrophages are recruited to sites of infection and inflammation by signaling proteins secreted by other leukocytes. The differentiation of the monocyte blood cell lineage can be studied in vitro using cultured cell lines. For example, THP-1 is a human promonocyte cell line that can be

activated by treatment with both phorbol ester such as phorbol myristate acetate (PMA), and lipopolysaccharide (LPS). PMA is a broad activator of the protein kinase C-dependent pathways.

Monocytes are involved in the initiation and maintenance of inflammatory immune responses. The outer membrane of gram-negative bacteria expresses lipopolysaccharide (LPS) complexes called endotoxins. Toxicity is associated with the lipid component (Lipid A) of LPS, and immunogenicity is associated with the polysaccharide components of LPS. LPS elicits a variety of inflammatory responses, and because it activates complement by the alternative (properdin) pathway, it is often part of the pathology of gram-negative bacterial infections. For the most part, endotoxins remain associated with the cell wall until the bacteria disintegrate. LPS released into the bloodstream by lysing gram-negative bacteria is first bound by certain plasma proteins identified as LPS-binding proteins. The LPS-binding protein complex interacts with CD14 receptors on monocytes, macrophages, B cells, and other types of receptors on endothelial cells. Activation of human B cells with LPS results in mitogenesis as well as immunoglobulin synthesis. In monocytes and macrophages three types of events are triggered during their interaction with LPS: 1) Production of cytokines, including IL-1, IL-6, IL-8, TNF- $\alpha$ , and platelet-activating factor, which stimulate production of prostaglandins and leukotrienes that mediate inflammation and septic shock; 2) Activation of the complement cascade; and 3) Activation of the coagulation cascade.

There is a need in the art for new compositions, including nucleic acids and proteins, for the diagnosis, prevention, and treatment of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, reproductive, endocrine, metabolic, pancreatic disorders, disorders associated with the adrenals, disorders associated with gonadal steroid hormones, cancers, and infections.

## SUMMARY OF THE INVENTION

Various embodiments of the invention provide purified polypeptides, protein modification and maintenance molecules, referred to collectively as 'PMMM' and individually as 'PMMM-1,' 'PMMM-2,' 'PMMM-3,' 'PMMM-4,' 'PMMM-5,' 'PMMM-6,' 'PMMM-7,' 'PMMM-8,' 'PMMM-9,' 'PMMM-10,' 'PMMM-11,' 'PMMM-12,' 'PMMM-13,' 'PMMM-14,' 'PMMM-15,' 'PMMM-16,' 'PMMM-17,' 'PMMM-18,' 'PMMM-19,' 'PMMM-20,' 'PMMM-21,' 'PMMM-22,' 'PMMM-23,' 'PMMM-24,' 'PMMM-25,' 'PMMM-26,' 'PMMM-27,' 'PMMM-28,' 'PMMM-29,' 'PMMM-30,' and 'PMMM-31' and methods for using these proteins and their encoding polynucleotides for the detection, diagnosis, and treatment of diseases and medical conditions. Embodiments also provide methods for utilizing the purified protein modification and maintenance molecules and/or their encoding polynucleotides for facilitating the drug discovery process, including determination of efficacy, dosage, toxicity, and pharmacology. Related embodiments provide methods for utilizing the

purified protein modification and maintenance molecules and/or their encoding polynucleotides for investigating the pathogenesis of diseases and medical conditions.

An embodiment provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. Another embodiment provides an isolated polypeptide comprising an amino acid sequence of SEQ ID NO:1-31.

Still another embodiment provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. In another embodiment, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-31. In an alternative embodiment, the polynucleotide is selected from the group consisting of SEQ ID NO:32-62.

Still another embodiment provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. Another embodiment provides a cell transformed with the recombinant polynucleotide. Yet another embodiment provides a transgenic organism comprising the recombinant polynucleotide.

Another embodiment provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group

consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Yet another embodiment provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31.

Still yet another embodiment provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In other embodiments, the polynucleotide can comprise at least about 20, 30, 40, 60, 80, or 100 contiguous nucleotides.

Yet another embodiment provides a method for detecting a target polynucleotide in a sample, said target polynucleotide being selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex. In a related embodiment, the

method can include detecting the amount of the hybridization complex. In still other embodiments, the probe can comprise at least about 20, 30, 40, 60, 80, or 100 contiguous nucleotides.

Still yet another embodiment provides a method for detecting a target polynucleotide in a sample, said target polynucleotide being selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof. In a related embodiment, the method can include detecting the amount of the amplified target polynucleotide or fragment thereof.

Another embodiment provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and a pharmaceutically acceptable excipient. In one embodiment, the composition can comprise an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. Other embodiments provide a method of treating a disease or condition associated with decreased or abnormal expression of functional PMMM, comprising administering to a patient in need of such treatment the composition.

Yet another embodiment provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. Another embodiment provides a composition comprising an agonist compound identified by the method and a

pharmaceutically acceptable excipient. Yet another embodiment provides a method of treating a disease or condition associated with decreased expression of functional PMMM, comprising administering to a patient in need of such treatment the composition.

Still yet another embodiment provides a method for screening a compound for effectiveness  
5 as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group  
10 consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. Another embodiment provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. Yet another embodiment  
15 provides a method of treating a disease or condition associated with overexpression of functional PMMM, comprising administering to a patient in need of such treatment the composition.

Another embodiment provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a  
20 naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31. The method comprises a) combining the  
25 polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

Yet another embodiment provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino  
30 acid sequence selected from the group consisting of SEQ ID NO:1-31, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical or at least about 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-31, and d) an immunogenic fragment of a polypeptide having an amino acid sequence  
35 selected from the group consisting of SEQ ID NO:1-31. The method comprises a) combining the

polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

Still yet another embodiment provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, b) detecting altered expression of the target polynucleotide, and c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

Another embodiment provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical or at least about 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide can comprise a fragment of a polynucleotide selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.



## BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for full length polynucleotide and polypeptide embodiments of the invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog, and the PROTEOME database identification numbers and annotations of PROTEOME database homologs, for polypeptide embodiments of the invention. The probability scores for the matches between each polypeptide and its homolog(s) are also shown.

Table 3 shows structural features of polypeptide embodiments, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide embodiments, along with selected fragments of the polynucleotides.

Table 5 shows representative cDNA libraries for polynucleotide embodiments.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze polynucleotides and polypeptides, along with applicable descriptions, references, and threshold parameters.

Table 8 shows single nucleotide polymorphisms found in polynucleotide sequences of the invention, along with allele frequencies in different human populations.

## DESCRIPTION OF THE INVENTION

Before the present proteins, nucleic acids, and methods are described, it is understood that embodiments of the invention are not limited to the particular machines, instruments, materials, and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the invention.

As used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing

the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with various embodiments of the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

## 5 DEFINITIONS

"PMMM" refers to the amino acid sequences of substantially purified PMMM obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of PMMM. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PMMM either by directly interacting with PMMM or by acting on components of the biological pathway in which PMMM participates.

An "allelic variant" is an alternative form of the gene encoding PMMM. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding PMMM include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as PMMM or a polypeptide with at least one functional characteristic of PMMM. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding PMMM, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide encoding PMMM. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent PMMM. Deliberate amino acid substitutions may be made on the basis of one or more similarities in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of PMMM is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" can refer to an oligopeptide, a peptide, a polypeptide, or a protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid. Amplification may be carried out using polymerase chain reaction (PCR) technologies or other nucleic acid amplification technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of PMMM. Antagonists may include proteins such as antibodies, anticalins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PMMM either by directly interacting with PMMM or by acting on components of the biological pathway in which PMMM participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind PMMM polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "aptamer" refers to a nucleic acid or oligonucleotide molecule that binds to a specific molecular target. Aptamers are derived from an *in vitro* evolutionary process (e.g., SELEX (Systematic Evolution of Ligands by EXponential Enrichment), described in U.S. Patent No. 5,270,163), which selects for target-specific aptamer sequences from large combinatorial libraries. Aptamer compositions may be double-stranded or single-stranded, and may include deoxyribonucleotides, ribonucleotides, nucleotide derivatives, or other nucleotide-like molecules.

The nucleotide components of an aptamer may have modified sugar groups (e.g., the 2'-OH group of a ribonucleotide may be replaced by 2'-F or 2'-NH<sub>2</sub>), which may improve a desired property, e.g., resistance to nucleases or longer lifetime in blood. Aptamers may be conjugated to other molecules, e.g., a high molecular weight carrier to slow clearance of the aptamer from the circulatory system.

5 Aptamers may be specifically cross-linked to their cognate ligands, e.g., by photo-activation of a cross-linker (Brody, E.N. and L. Gold (2000) J. Biotechnol. 74:5-13).

The term "intramer" refers to an aptamer which is expressed *in vivo*. For example, a vaccinia virus-based RNA expression system has been used to express specific RNA aptamers at high levels in the cytoplasm of leukocytes (Blind, M. et al. (1999) Proc. Natl. Acad. Sci. USA 96:3606-3610).

10 The term "spiegelmer" refers to an aptamer which includes L-DNA, L-RNA, or other left-handed nucleotide derivatives or nucleotide-like molecules. Aptamers containing left-handed nucleotides are resistant to degradation by naturally occurring enzymes, which normally act on substrates containing right-handed nucleotides.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a polynucleotide having a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

25 The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic PMMM, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

30 "Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide" and a "composition comprising a given polypeptide" can refer to any composition containing the given polynucleotide or polypeptide. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising

35

polynucleotides encoding PMMM or fragments of PMMM may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (Accelrys, Burlington MA) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
20	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
25	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
30	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
35	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide.

Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

"Exon shuffling" refers to the recombination of different coding regions (exons). Since an exon may represent a structural or functional domain of the encoded protein, new proteins may be assembled through the novel reassortment of stable substructures, thus allowing acceleration of the evolution of new protein functions.

A "fragment" is a unique portion of PMMM or a polynucleotide encoding PMMM which can be identical in sequence to, but shorter in length than, the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from about 5 to about 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:32-62 can comprise a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:32-62, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:32-62 can be employed in one or more embodiments of methods of the invention, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:32-62 from related

polynucleotides. The precise length of a fragment of SEQ ID NO:32-62 and the region of SEQ ID NO:32-62 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

5 A fragment of SEQ ID NO:1-31 is encoded by a fragment of SEQ ID NO:32-62. A fragment of SEQ ID NO:1-31 can comprise a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-31. For example, a fragment of SEQ ID NO:1-31 can be used as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-31. The precise length of a fragment of SEQ ID NO:1-31 and the region of SEQ ID NO:1-31 to which the fragment corresponds can be determined based on the intended purpose for the fragment using one or more  
10 analytical methods described herein or otherwise known in the art.

A "full length" polynucleotide is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, alternatively, sequence identity, between two or  
15 more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of identical residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences,  
20 and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using one or more computer algorithms or programs known in the art or described herein. For example, percent identity can be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the  
25 LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989; CABIOS 5:151-153) and in Higgins, D.G. et al. (1992; CABIOS 8:189-191). For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default.

30 Alternatively, a suite of commonly used and freely available sequence comparison algorithms which can be used is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence  
35 analysis programs including "blastn," that is used to align a known polynucleotide sequence with

other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

*Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of identical residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide. The phrases "percent similarity" and "% similarity," as applied to polypeptide sequences, refer to the percentage of residue matches, including identical residue matches and conservative substitutions, between at least two polypeptide sequences aligned using a standardized algorithm. In contrast,



conservative substitutions are not included in the calculation of percent identity between polypeptide sequences.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity.

Specific hybridization complexes form under permissive annealing conditions and remain hybridized

after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. and D.W. Russell (2001; Molecular Cloning: A Laboratory Manual, 3rd ed., vol. 1-3, Cold Spring Harbor Press, Cold Spring Harbor NY, ch. 9).

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acids by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid present in solution and another nucleic acid immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or polynucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

5 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of PMMM which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of PMMM which is useful in any of the antibody production methods disclosed herein or known in the art.

10 The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, antibodies, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, antibody, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of PMMM. For example, modulation  
15 may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of PMMM.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the  
20 antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where  
25 necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript  
30 elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an PMMM may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of PMMM.

“Probe” refers to nucleic acids encoding PMMM, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acids. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. “Primers” are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in, for example, Sambrook, J. and D.W. Russell (2001; Molecular Cloning: A Laboratory Manual, 3rd ed., vol. 1-3, Cold Spring Harbor Press, Cold Spring Harbor NY), Ausubel, F.M. et al. (1999; Short Protocols in Molecular Biology, 4<sup>th</sup> ed., John Wiley & Sons, New York NY), and Innis, M. et al. (1990; PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA). PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a “mispriming library,” in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user’s specific

needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both  
5 unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

10 A "recombinant nucleic acid" is a nucleic acid that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook and Russell (*supra*). The term  
15 recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a  
20 vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription,  
25 translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

30 An "RNA equivalent," in reference to a DNA molecule, is composed of the same linear sequence of nucleotides as the reference DNA molecule with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing PMMM,  
35 nucleic acids encoding PMMM, or fragments thereof may comprise a bodily fluid; an extract from a

cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably at least about 75% free, and most preferably at least about 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" or "expression profile" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with

a recombinant virus. In another embodiment, the nucleic acid can be introduced by infection with a recombinant viral vector, such as a lentiviral vector (Lois, C. et al. (2002) Science 295:868-872). The term genetic manipulation does not include classical cross-breeding, or *in vitro* fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook and Russell (*supra*).

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotides that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity or sequence similarity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity or sequence similarity over a certain defined length of one of the polypeptides.

## THE INVENTION

Various embodiments of the invention include new human protein modification and maintenance molecules (PMMM), the polynucleotides encoding PMMM, and the use of these compositions for the diagnosis, treatment, or prevention of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, reproductive, endocrine, metabolic, pancreatic disorders, disorders associated with the adrenals, disorders associated with gonadal steroid hormones, cancers, and infections.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide embodiments of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown. Column 6 shows the Incyte ID numbers of physical, full length clones corresponding to the polypeptide and polynucleotide sequences of the invention. The full length clones encode polypeptides which have at least 95% sequence identity to the polypeptide sequences shown in column 3.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database and the PROTEOME database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (GenBank ID NO:) of the nearest GenBank homolog and the PROTEOME database identification numbers (PROTEOME ID NO:) of the nearest PROTEOME database homologs. Column 4 shows the probability scores for the matches between each polypeptide and its homolog(s). Column 5 shows the annotation of the GenBank and PROTEOME database homolog(s) along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Accelrys, Burlington MA). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column



7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are protein modification and maintenance molecules. For example, SEQ ID NO:2 is 86% identical, from residue M1 to residue E738 and 96% identical, from residue K607 to residue L900, to human inter-alpha-trypsin inhibitor family heavy chain-related protein (GenBank ID g4096840) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability scores are 0.0 and 7.3e-152, which indicate the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:2 also contains a von Willebrand factor type A domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and additional BLAST analyses provide further corroborative evidence that SEQ ID NO:2 is a protease inhibitor.

In another example, SEQ ID NO:9 is 50% identical, from residue M1 to residue G378, to Mus musculus mDj10 (GenBank ID g6567172) as determined by BLAST. The BLAST probability score is 9.7e-102. SEQ ID NO:9 also contains a DnaJ domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database. Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:9 is a molecular chaperone.

In another example, SEQ ID NO:12 is 100% identical, from residue M1 to residue N344, to human phosphatidyl inositol glycan class T (GenBank ID g14456615) as determined by BLAST. The BLAST probability score is 5.4e-280. Data from BLAST-PRODOM analysis provides further corroborative evidence that SEQ ID NO:12 is a phosphatidyl inositol glycan. In an alternative example, SEQ ID NO:13 is 100% identical, from residue D63 to residue L476, to human phosphatidyl inositol glycan class T (GenBank ID g14456615) as determined by BLAST. The BLAST probability score is 4.7e-261. Data from BLAST-PRODOM analysis provides further corroborative evidence that SEQ ID NO:13 is a phosphatidyl inositol glycan.

In yet another example, SEQ ID NO:15 is 97% identical, from residue D50 to residue D121, to human ubiquitin-conjugating enzyme HR6B (GenBank ID g11037550) as determined by BLAST. The BLAST probability score is 2.1e-58. SEQ ID NO:15 is localized to the subcellular region, has ubiquitination function, and is a protein conjugation factor as determined by BLAST analysis using the PROTEOME database. SEQ ID NO:15 also contains an ubiquitin-conjugating enzyme domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database. Data from BLAST-PRODOM, BLAST-DOMO, and PROFILESCAN

analyses provide further corroborative evidence that SEQ ID NO:15 is a ubiquitin-conjugating enzyme.

In a further example, SEQ ID NO:19 is 100% identical, from residue M1 to residue G82, and 100% identical, from residue G82 to residue A652, to the large subunit of human CANP (GenBank ID g29664, residues M1-G82 and G144-A714 respectively) as determined by BLAST. The BLAST probability score is 0.0. SEQ ID NO:19 is homologous to other proteins, such as calpain, the large subunit of a cysteine protease, having cysteine protease activity and localized to the plasma membrane, as determined by BLAST analysis using the PROTEOME database. SEQ ID NO:19 also contains calpain and EF hand domains as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. Data from BLIMPS, MOTIFS, and BLAST analyses provide further corroborative evidence that SEQ ID NO:19 is a calpain cysteine protease. SEQ ID NO:1, SEQ ID NO:3-8, SEQ ID NO:10-11, SEQ ID NO:14, SEQ ID NO:16-18, and SEQ ID NO:20-31 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-31 are described in Table 7.

As shown in Table 4, the full length polynucleotide embodiments were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Column 1 lists the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:), the corresponding Incyte polynucleotide consensus sequence number (Incyte ID) for each polynucleotide of the invention, and the length of each polynucleotide sequence in basepairs. Column 2 shows the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences used to assemble the full length polynucleotide embodiments, and of fragments of the polynucleotides which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:32-62 or that distinguish between SEQ ID NO:32-62 and related polynucleotides.

The polynucleotide fragments described in Column 2 of Table 4 may refer specifically, for example, to Incyte cDNAs derived from tissue-specific cDNA libraries or from pooled cDNA libraries. Alternatively, the polynucleotide fragments described in column 2 may refer to GenBank cDNAs or ESTs which contributed to the assembly of the full length polynucleotides. In addition, the polynucleotide fragments described in column 2 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (*i.e.*, those sequences including the designation "ENST"). Alternatively, the polynucleotide fragments described in column 2 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (*i.e.*, those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (*i.e.*, those sequences including the designation "NP"). Alternatively, the polynucleotide fragments described in column 2 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon

stitching" algorithm. For example, a polynucleotide sequence identified as

FL\_XXXXXX\_N<sub>1</sub>\_N<sub>2</sub>\_YYYYY\_N<sub>3</sub>\_N<sub>4</sub> represents a "stitched" sequence in which XXXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYY is the number of the prediction generated by the algorithm, and N<sub>1,2,3...</sub>, if present, represent specific

exons that may have been manually edited during analysis (See Example V). Alternatively, the polynucleotide fragments in column 2 may refer to assemblages of exons brought together by an "exon-stretching" algorithm. For example, a polynucleotide sequence identified as

FLXXXXXX\_gAAAAA\_gBBBBB\_1\_N is a "stretched" sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human

genomic sequence to which the "exon-stretching" algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq sequence was used as a protein homolog for the "exon-stretching" algorithm, a RefSeq identifier (denoted by "NM," "NP," or "NT") may be used in place of the GenBank identifier (i.e., gBBBBB).

Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis methods associated with the prefixes (see Example IV and Example V).

Prefix	Type of analysis and/or examples of programs
GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGENES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
GBI	Hand-edited analysis of genomic sequences.
FL	Stitched or stretched genomic sequences (see Example V).
INCY	Full length transcript and exon prediction from mapping of EST sequences to the genome. Genomic location and EST composition data are combined to predict the exons and resulting transcript.

In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in Table 4 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotides which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to

assemble and confirm the above polynucleotides. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

Table 8 shows single nucleotide polymorphisms (SNPs) found in polynucleotide sequences of the invention, along with allele frequencies in different human populations. Columns 1 and 2 show the polynucleotide sequence identification number (SEQ ID NO:) and the corresponding Incyte project identification number (PID) for polynucleotides of the invention. Column 3 shows the Incyte identification number for the EST in which the SNP was detected (EST ID), and column 4 shows the identification number for the SNP (SNP ID). Column 5 shows the position within the EST sequence at which the SNP is located (EST SNP), and column 6 shows the position of the SNP within the full-length polynucleotide sequence (CB1 SNP). Column 7 shows the allele found in the EST sequence. Columns 8 and 9 show the two alleles found at the SNP site. Column 10 shows the amino acid encoded by the codon including the SNP site, based upon the allele found in the EST. Columns 11-14 show the frequency of allele 1 in four different human populations. An entry of n/d (not detected) indicates that the frequency of allele 1 in the population was too low to be detected, while n/a (not available) indicates that the allele frequency was not determined for the population.

The invention also encompasses PMMM variants. Various embodiments of PMMM variants can have at least about 80%, at least about 90%, or at least about 95% amino acid sequence identity to the PMMM amino acid sequence, and can contain at least one functional or structural characteristic of PMMM.

Various embodiments also encompass polynucleotides which encode PMMM. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:32-62, which encodes PMMM. The polynucleotide sequences of SEQ ID NO:32-62, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses variants of a polynucleotide encoding PMMM. In particular, such a variant polynucleotide will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a polynucleotide encoding PMMM. A particular aspect of the invention encompasses a variant of a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NO:32-62 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:32-62. Any one of the polynucleotide variants described above can encode a polypeptide which contains at least one functional or structural characteristic of PMMM.

In addition, or in the alternative, a polynucleotide variant of the invention is a splice variant of a polynucleotide encoding PMMM. A splice variant may have portions which have significant sequence identity to a polynucleotide encoding PMMM, but will generally have a greater or lesser number of polynucleotides due to additions or deletions of blocks of sequence arising from alternate splicing of exons during mRNA processing. A splice variant may have less than about 70%, or alternatively less than about 60%, or alternatively less than about 50% polynucleotide sequence identity to a polynucleotide encoding PMMM over its entire length; however, portions of the splice variant will have at least about 70%, or alternatively at least about 85%, or alternatively at least about 95%, or alternatively 100% polynucleotide sequence identity to portions of the polynucleotide encoding PMMM. For example, a polynucleotide comprising a sequence of SEQ ID NO:43 and a polynucleotide comprising a sequence of SEQ ID NO:44 are splice variants of each other. Any one of the splice variants described above can encode a polypeptide which contains at least one functional or structural characteristic of PMMM.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding PMMM, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring PMMM, and all such variations are to be considered as being specifically disclosed.

Although polynucleotides which encode PMMM and its variants are generally capable of hybridizing to polynucleotides encoding naturally occurring PMMM under appropriately selected conditions of stringency, it may be advantageous to produce polynucleotides encoding PMMM or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PMMM and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of polynucleotides which encode PMMM and PMMM derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic polynucleotide may be inserted into any of the many available expression vectors and cell

systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a polynucleotide encoding PMMM or any fragment thereof.

Embodiments of the invention can also include polynucleotides that are capable of hybridizing to the claimed polynucleotides, and, in particular, to those having the sequences shown in SEQ ID NO:32-62 and fragments thereof, under various conditions of stringency (Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511). Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Biosciences, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Invitrogen, Carlsbad CA). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Amersham Biosciences), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art (Ausubel et al., *supra*, ch. 7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853).

The nucleic acids encoding PMMM may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector (Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322). Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences (Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186). A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA (Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119). In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art (Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may

use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or  
5 another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T)  
10 library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the  
15 emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments  
20 which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotides or fragments thereof which encode PMMM may be cloned in recombinant DNA molecules that direct expression of PMMM, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other polynucleotides which encode substantially the same or a functionally  
25 equivalent polypeptides may be produced and used to express PMMM.

The polynucleotides of the invention can be engineered using methods generally known in the art in order to alter PMMM-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be  
30 used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent No.  
35 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat.

Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of PMMM, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then  
5 subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are  
10 optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, polynucleotides encoding PMMM may be synthesized, in whole or in  
15 part, using one or more chemical methods well known in the art (Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232). Alternatively, PMMM itself or a fragment thereof may be synthesized using chemical methods known in the art. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques (Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH  
20 Freeman, New York NY, pp. 55-60; Roberge, J.Y. et al. (1995) Science 269:202-204). Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of PMMM, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

25 The peptide may be substantially purified by preparative high performance liquid chromatography (Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421). The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing (Creighton, *supra*, pp. 28-53).

In order to express a biologically active PMMM, the polynucleotides encoding PMMM or  
30 derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotides encoding PMMM. Such elements may vary in their strength and specificity. Specific initiation signals may  
35 also be used to achieve more efficient translation of polynucleotides encoding PMMM. Such signals



include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where a polynucleotide sequence encoding PMMM and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

Methods which are well known to those skilled in the art may be used to construct expression vectors containing polynucleotides encoding PMMM and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination (Sambrook and Russell, *supra*, ch. 1-4, and 8; Ausubel et al., *supra*, ch. 1, 3, and 15).

A variety of expression vector/host systems may be utilized to contain and express polynucleotides encoding PMMM. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems (Sambrook and Russell, *supra*; Ausubel et al., *supra*; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355). Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of polynucleotides to the targeted organ, tissue, or cell population (Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5:350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6340-6344; Buller, R.M. et al. (1985) *Nature* 317:813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31:219-226; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242). The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotides encoding PMMM. For example, routine cloning, subcloning, and propagation of polynucleotides encoding PMMM can be achieved using a

multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Invitrogen). Ligation of polynucleotides encoding PMMM into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for  
5 *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence (Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509). When large quantities of PMMM are needed, e.g. for the production of antibodies, vectors which direct high level expression of PMMM may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

10 Yeast expression systems may be used for production of PMMM. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign polynucleotide sequences into the host genome for stable propagation (Ausubel  
15 et al., *supra*; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; Scorer, C.A. et al. (1994) Bio/Technology 12:181-184).

Plant systems may also be used for expression of PMMM. Transcription of polynucleotides encoding PMMM may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J.  
20 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection (The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New  
25 York NY, pp. 191-196).

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, polynucleotides encoding PMMM may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be  
30 used to obtain infective virus which expresses PMMM in host cells (Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of  
35 DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are

constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes (Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355).

For long term production of recombinant proteins in mammalian systems, stable expression of PMMM in cell lines is preferred. For example, polynucleotides encoding PMMM can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively (Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823). Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14). Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites (Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051). Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$ -glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding PMMM is inserted within a marker gene sequence, transformed cells containing polynucleotides encoding PMMM can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PMMM under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the polynucleotide encoding PMMM and that express PMMM may be identified by a variety of procedures known to those of skill in the art. These

procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of PMMM using either  
5 specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PMMM is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art  
10 (Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ).

A wide variety of labels and conjugation techniques are known by those skilled in the art and  
15 may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding PMMM include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, polynucleotides encoding PMMM, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially  
20 available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Biosciences, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or  
25 chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with polynucleotides encoding PMMM may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors  
30 containing polynucleotides which encode PMMM may be designed to contain signal sequences which direct secretion of PMMM through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted polynucleotides or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation,  
35 glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves

a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct  
5 modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant polynucleotides encoding PMMM may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric PMMM protein containing a heterologous moiety that can be recognized by a commercially available antibody may  
10 facilitate the screening of peptide libraries for inhibitors of PMMM activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their  
15 cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the PMMM encoding sequence and the heterologous protein  
20 sequence, so that PMMM may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel et al. (*supra*, ch. 10 and 16). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In another embodiment, synthesis of radiolabeled PMMM may be achieved *in vitro* using the  
25 TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

PMMM, fragments of PMMM, or variants of PMMM may be used to screen for compounds  
30 that specifically bind to PMMM. One or more test compounds may be screened for specific binding to PMMM. In various embodiments, 1, 2, 3, 4, 5, 10, 20, 50, 100, or 200 test compounds can be screened for specific binding to PMMM. Examples of test compounds can include antibodies, anticalins, oligonucleotides, proteins (e.g., ligands or receptors), or small molecules.

In related embodiments, variants of PMMM can be used to screen for binding of test  
35 compounds, such as antibodies, to PMMM, a variant of PMMM, or a combination of PMMM and/or

one or more variants PMMM. In an embodiment, a variant of PMMM can be used to screen for compounds that bind to a variant of PMMM, but not to PMMM having the exact sequence of a sequence of SEQ ID NO:1-31. PMMM variants used to perform such screening can have a range of about 50% to about 99% sequence identity to PMMM, with various embodiments having 60%, 70%,  
5 75%, 80%, 85%, 90%, and 95% sequence identity.

In an embodiment, a compound identified in a screen for specific binding to PMMM can be closely related to the natural ligand of PMMM, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner (Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2):Chapter 5). In another embodiment, the compound thus identified can  
10 be a natural ligand of a receptor PMMM (Howard, A.D. et al. (2001) Trends Pharmacol. Sci.22:132-140; Wise, A. et al. (2002) Drug Discovery Today 7:235-246).

In other embodiments, a compound identified in a screen for specific binding to PMMM can be closely related to the natural receptor to which PMMM binds, at least a fragment of the receptor, or a fragment of the receptor including all or a portion of the ligand binding site or binding pocket.  
15 For example, the compound may be a receptor for PMMM which is capable of propagating a signal, or a decoy receptor for PMMM which is not capable of propagating a signal (Ashkenazi, A. and V.M. Divit (1999) Curr. Opin. Cell Biol. 11:255-260; Mantovani, A. et al. (2001) Trends Immunol. 22:328-336). The compound can be rationally designed using known techniques. Examples of such techniques include those used to construct the compound etanercept (ENBREL; Amgen Inc.,  
20 Thousand Oaks CA), which is efficacious for treating rheumatoid arthritis in humans. Etanercept is an engineered p75 tumor necrosis factor (TNF) receptor dimer linked to the Fc portion of human IgG<sub>1</sub> (Taylor, P.C. et al. (2001) Curr. Opin. Immunol. 13:611-616).

In one embodiment, two or more antibodies having similar or, alternatively, different specificities can be screened for specific binding to PMMM, fragments of PMMM, or variants of  
25 PMMM. The binding specificity of the antibodies thus screened can thereby be selected to identify particular fragments or variants of PMMM. In one embodiment, an antibody can be selected such that its binding specificity allows for preferential identification of specific fragments or variants of PMMM. In another embodiment, an antibody can be selected such that its binding specificity allows for preferential diagnosis of a specific disease or condition having increased, decreased, or otherwise  
30 abnormal production of PMMM.

In an embodiment, anticalins can be screened for specific binding to PMMM, fragments of PMMM, or variants of PMMM. Anticalins are ligand-binding proteins that have been constructed based on a lipocalin scaffold (Weiss, G.A. and H.B. Lowman (2000) Chem. Biol. 7:R177-R184; Skerra, A. (2001) J. Biotechnol. 74:257-275). The protein architecture of lipocalins can include a  
35 beta-barrel having eight antiparallel beta-strands, which supports four loops at its open end. These

loops form the natural ligand-binding site of the lipocalins, a site which can be re-engineered *in vitro* by amino acid substitutions to impart novel binding specificities. The amino acid substitutions can be made using methods known in the art or described herein, and can include conservative substitutions (e.g., substitutions that do not alter binding specificity) or substitutions that modestly, moderately, or significantly alter binding specificity.

In one embodiment, screening for compounds which specifically bind to, stimulate, or inhibit PMMM involves producing appropriate cells which express PMMM, either as a secreted protein or on the cell membrane. Preferred cells can include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing PMMM or cell membrane fractions which contain PMMM are then contacted with a test compound and binding, stimulation, or inhibition of activity of either PMMM or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with PMMM, either in solution or affixed to a solid support, and detecting the binding of PMMM to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

An assay can be used to assess the ability of a compound to bind to its natural ligand and/or to inhibit the binding of its natural ligand to its natural receptors. Examples of such assays include radio-labeling assays such as those described in U.S. Patent No. 5,914,236 and U.S. Patent No. 6,372,724. In a related embodiment, one or more amino acid substitutions can be introduced into a polypeptide compound (such as a receptor) to improve or alter its ability to bind to its natural ligands (Matthews, D.J. and J.A. Wells. (1994) Chem. Biol. 1:25-30). In another related embodiment, one or more amino acid substitutions can be introduced into a polypeptide compound (such as a ligand) to improve or alter its ability to bind to its natural receptors (Cunningham, B.C. and J.A. Wells (1991) Proc. Natl. Acad. Sci. USA 88:3407-3411; Lowman, H.B. et al. (1991) J. Biol. Chem. 266:10982-10988).

PMMM, fragments of PMMM, or variants of PMMM may be used to screen for compounds that modulate the activity of PMMM. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for PMMM activity, wherein PMMM is combined with at least one test compound, and the activity of PMMM in the presence of a test compound is compared with the activity of PMMM in the absence of the test compound. A change in the activity of PMMM in the presence of the test compound is

indicative of a compound that modulates the activity of PMMM. Alternatively, a test compound is combined with an *in vitro* or cell-free system comprising PMMM under conditions suitable for PMMM activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of PMMM may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding PMMM or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease (see, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337). For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (*neo*; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding PMMM may also be manipulated *in vitro* in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding PMMM can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding PMMM is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress PMMM, e.g., by secreting PMMM in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).



## THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of PMMM and protein modification and maintenance molecules. In addition, examples of tissues expressing PMMM can be found in Table 6 and can also be found in Example XI.

- 5 Therefore, PMMM appears to play a role in gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, reproductive, endocrine, metabolic, pancreatic disorders, disorders associated with the adrenals, disorders associated with gonadal steroid hormones, cancers, and infections. In the treatment of disorders associated with increased PMMM expression or activity, it is desirable to decrease the expression or  
10 activity of PMMM. In the treatment of disorders associated with decreased PMMM expression or activity, it is desirable to increase the expression or activity of PMMM.

- Therefore, in one embodiment, PMMM or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMMM. Examples of such disorders include, but are not limited to, a gastrointestinal  
15 disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis,  
20 passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha<sub>1</sub>-  
25 antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis,  
30 Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular  
35 calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective

endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; an autoimmune/inflammatory disease, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, atherosclerotic plaque rupture, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, degradation of articular cartilage, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, bone resorption, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, age-related macular degeneration, and sensorineural hearing loss; an epithelial disorder, such as dyshidrotic eczema, allergic contact dermatitis, keratosis pilaris, melasma, vitiligo, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, seborrheic keratosis, folliculitis, herpes simplex, herpes zoster, varicella, candidiasis, dermatophytosis, scabies, insect bites, cherry angioma, keloid, dermatofibroma, acrochordons, urticaria, transient acantholytic dermatosis, xerosis, eczema, atopic dermatitis, contact dermatitis, hand eczema, nummular eczema, lichen simplex chronicus, asteatotic eczema, stasis dermatitis and stasis ulceration, seborrheic dermatitis, psoriasis, lichen planus, pityriasis rosea, impetigo, ecthyma, dermatophytosis, tinea versicolor, warts, acne vulgaris, acne rosacea, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, bullous pemphigoid, herpes gestationis, dermatitis

- herpetiformis, linear IgA disease, epidermolysis bullosa acquisita, dermatomyositis, lupus erythematosus, scleroderma and morphea, erythroderma, alopecia, figurate skin lesions, telangiectasias, hypopigmentation, hyperpigmentation, vesicles/bullae, exanthems, cutaneous drug reactions, papulonodular skin lesions, chronic non-healing wounds, photosensitivity diseases,
- 5 epidermolysis bullosa simplex, epidermolytic hyperkeratosis, epidermolytic and nonepidermolytic palmoplantar keratoderma, ichthyosis bullosa of Siemens, ichthyosis exfoliativa, keratosis palmaris et plantaris, keratosis palmoplantaris, palmoplantar keratoderma, keratosis punctata, Meesmann's corneal dystrophy, pachyonychia congenita, white sponge nevus, steatocystoma multiplex, epidermal nevi/epidermolytic hyperkeratosis type, monilethrix, trichothiodystrophy, chronic
- 10 hepatitis/cryptogenic cirrhosis, and colorectal hyperplasia; a neurological disorder, such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and
- 15 viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental
- 20 retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety,
- 25 and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a reproductive disorder, such as infertility, including tubal disease, ovulatory defects, and endometriosis, a disorder of prolactin production, a disruption of the estrous
- 30 cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the
- 35 male breast, and gynecomastia; an endocrine disorder such as a disorder of the hypothalamus and/or

pituitary resulting from lesions such as a primary brain tumor, adenoma, infarction associated with pregnancy, hypophysectomy, aneurysm, vascular malformation, thrombosis, infection, immunological disorder, and complication due to head trauma; a disorder associated with hypopituitarism including hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease,

5 Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism; a disorder associated with hyperpituitarism including acromegaly, gigantism, and syndrome of inappropriate antidiuretic hormone (ADH) secretion (SIADH) often caused by benign adenoma; a disorder associated with hypothyroidism including goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis

10 (Hashimoto's disease), and cretinism; a disorder associated with hyperthyroidism including thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter, thyroid carcinoma, and Plummer's disease; a disorder associated with hyperparathyroidism including Conn disease (chronic hypercalcemia); a metabolic disorder such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis,

15 diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, mannosidosis, neuraminidase deficiency, obesity, pentosuria

20 phenylketonuria, pseudovitamin D-deficiency rickets; a disorder of carbohydrate metabolism such as congenital type II dyserythropoietic anemia, diabetes, insulin-dependent diabetes mellitus, non-insulin-dependent diabetes mellitus, fructose-1,6-diphosphatase deficiency, galactosemia, glucagonoma, hereditary fructose intolerance, hypoglycemia, mannosidosis, neuraminidase deficiency, obesity, galactose epimerase deficiency, glycogen storage diseases, lysosomal storage

25 diseases, fructosuria, pentosuria, and inherited abnormalities of pyruvate metabolism; a disorder of lipid metabolism such as fatty liver, cholestasis, primary biliary cirrhosis, carnitine deficiency, carnitine palmitoyltransferase deficiency, myoadenylate deaminase deficiency, hypertriglyceridemia, lipid storage disorders such as Fabry's disease, Gaucher's disease, Niemann-Pick's disease, metachromatic leukodystrophy, adrenoleukodystrophy, GM<sub>2</sub> gangliosidosis, and ceroid

30 lipofuscinosis, abetalipoproteinemia, Tangier disease, hyperlipoproteinemia, diabetes mellitus, lipodystrophy, lipomatosis, acute panniculitis, disseminated fat necrosis, adiposis dolorosa, lipoid adrenal hyperplasia, minimal change disease, lipomas, atherosclerosis, hypercholesterolemia, hypercholesterolemia with hypertriglyceridemia, primary hypoalphalipoproteinemia, hypothyroidism, renal disease, liver disease, lecithin:cholesterol acyltransferase deficiency, cerebrotendinous

35 xanthomatosis, sitosterolemia, hypocholesterolemia, Tay-Sachs disease, Sandhoff's disease,

hyperlipidemia, hyperlipemia, lipid myopathies, and obesity; and a disorder of copper metabolism such as Menke's disease, Wilson's disease, and Ehlers-Danlos syndrome type IX; a pancreatic disorder such as Type I or Type II diabetes mellitus and associated complications; a disorder associated with the adrenals such as hyperplasia, carcinoma, or adenoma of the adrenal cortex,

5 hypertension associated with alkalosis, amyloidosis, hypokalemia, Cushing's disease, Liddle's syndrome, and Arnold-Healy-Gordon syndrome, pheochromocytoma tumors, and Addison's disease; a disorder associated with gonadal steroid hormones such as: in women, abnormal prolactin production, infertility, endometriosis, perturbation of the menstrual cycle, polycystic ovarian disease, hyperprolactinemia, isolated gonadotropin deficiency, amenorrhea, galactorrhea, hermaphroditism,

10 hirsutism and virilization, breast cancer, and, in post-menopausal women, osteoporosis; and, in men, Leydig cell deficiency, male climacteric phase, and germinal cell aplasia, a hypergonadal disorder associated with Leydig cell tumors, androgen resistance associated with absence of androgen receptors, syndrome of 5  $\alpha$ -reductase, and gynecomastia; a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal

15 gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an infection caused by a viral agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus,

20 reovirus, retrovirus, rhabdovirus, or togavirus; an infection caused by a bacterial agent classified as pneumococcus, staphylococcus, streptococcus, bacillus, corynebacterium, clostridium, meningococcus, gonococcus, listeria, moraxella, kingella, haemophilus, legionella, bordetella, gram-negative enterobacterium including shigella, salmonella, or campylobacter, pseudomonas, vibrio, brucella, francisella, yersinia, bartonella, norcardium, actinomyces, mycobacterium, spirochaetale,

25 rickettsia, chlamydia, or mycoplasma; an infection caused by a fungal agent classified as aspergillus, blastomyces, dermatophytes, cryptococcus, coccidioides, malassezia, histoplasma, or other mycosis-causing fungal agent; and an infection caused by a parasite classified as plasmodium or malaria-causing, parasitic entamoeba, leishmania, trypanosoma, toxoplasma, pneumocystis carinii, intestinal protozoa such as giardia, trichomonas, tissue nematode such as trichinella, intestinal nematode such

30 as ascaris, lymphatic filarial nematode, trematode such as schistosoma, and cestode such as tapeworm.

In another embodiment, a vector capable of expressing PMMM or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMMM including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified PMMM in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMMM including, but not limited to, those provided above.

5 In still another embodiment, an agonist which modulates the activity of PMMM may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PMMM including, but not limited to, those listed above.

In a further embodiment, an antagonist of PMMM may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PMMM. Examples of such  
10 disorders include, but are not limited to, those gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, reproductive, endocrine, metabolic, pancreatic disorders, disorders associated with the adrenals, disorders associated with gonadal steroid hormones, cancers, and infections described above. In one aspect, an antibody which specifically binds PMMM may be used directly as an antagonist or indirectly as a  
15 targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express PMMM.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding PMMM may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PMMM including, but not limited to, those described above.

20 In other embodiments, any protein, agonist, antagonist, antibody, complementary sequence, or vector embodiments may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various  
25 disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of PMMM may be produced using methods which are generally known in the art. In particular, purified PMMM may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind PMMM. Antibodies to PMMM may  
30 also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. In an embodiment, neutralizing antibodies (i.e., those which inhibit dimer formation) can be used therapeutically. Single chain antibodies (e.g., from camels or llamas) may be potent enzyme inhibitors and may have application in the design of peptide

mimetics, and in the development of immuno-adsorbents and biosensors (Muyltermans, S. (2001) J. Biotechnol. 74:277-302).

For the production of antibodies, various hosts including goats, rabbits, rats, mice, camels, dromedaries, llamas, humans, and others may be immunized by injection with PMMM or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol.

Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PMMM have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are substantially identical to a portion of the amino acid sequence of the natural protein.

Short stretches of PMMM amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to PMMM may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120).

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used (Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; Takeda, S. et al. (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce PMMM-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137).

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299).

Antibody fragments which contain specific binding sites for PMMM may also be generated. For example, such fragments include, but are not limited to,  $F(ab')_2$  fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the  $F(ab')_2$  fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W.D. et al. (1989) Science 246:1275-1281).

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between PMMM and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering PMMM epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for PMMM. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of PMMM-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple PMMM epitopes, represents the average affinity, or avidity, of the antibodies for PMMM. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular PMMM epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the PMMM-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of PMMM, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of PMMM-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available (Catty, *supra*; Coligan et al., *supra*).



In another embodiment of the invention, polynucleotides encoding PMMM, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding PMMM. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PMMM (Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press, Totawa NJ).

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein (Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102:469-475; Scanlon, K.J. et al. (1995) 9:1288-1296). Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors (Miller, A.D. (1990) *Blood* 76:271; Ausubel et al., *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63:323-347). Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art (Rossi, J.J. (1995) *Br. Med. Bull.* 51:217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87:1308-1315; Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25:2730-2736).

In another embodiment of the invention, polynucleotides encoding PMMM may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as *Candida albicans* and *Paracoccidioides brasiliensis*; and protozoan parasites such as *Plasmodium falciparum* and *Trypanosoma cruzi*). In the case where a genetic deficiency in PMMM expression or regulation causes disease, the expression of

PMMM from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in PMMM are treated by constructing mammalian expression vectors encoding PMMM and introducing these vectors by mechanical means into PMMM-deficient cells. Mechanical transfer technologies for use with cells *in vivo* or *ex vitro* include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J.-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of PMMM include, but are not limited to, the pCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX, PCR2-TOPOTA vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). PMMM may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, *supra*), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding PMMM from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to PMMM expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding PMMM under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences

required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for  
5 receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent No. 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant")  
10 discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716;  
15 Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In an embodiment, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding PMMM to cells which have one or more genetic abnormalities with respect to the expression of PMMM. The construction and packaging of adenovirus-based vectors are well  
20 known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent No. 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999; Annu.  
25 Rev. Nutr. 19:511-544) and Verma, I.M. and N. Somia (1997; Nature 18:389:239-242).

In another embodiment, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding PMMM to target cells which have one or more genetic abnormalities with respect to the expression of PMMM. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing PMMM to cells of the central nervous system, for which HSV has  
30 a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent No. 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby  
35 incorporated by reference. U.S. Patent No. 5,804,413 teaches the use of recombinant HSV d92 which

consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999; J. Virol. 73:519-532) and Xu, H. et al. (1994; Dev. Biol. 163:152-161). The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another embodiment, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding PMMM to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for PMMM into the alphavirus genome in place of the capsid-coding region results in the production of a large number of PMMM-coding RNAs and the synthesis of high levels of PMMM in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of PMMM into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177). A

complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of RNA molecules encoding PMMM.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA molecules encoding PMMM. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

In other embodiments of the invention, the expression of one or more selected polynucleotides of the present invention can be altered, inhibited, decreased, or silenced using RNA interference (RNAi) or post-transcriptional gene silencing (PTGS) methods known in the art. RNAi is a post-transcriptional mode of gene silencing in which double-stranded RNA (dsRNA) introduced into a targeted cell specifically suppresses the expression of the homologous gene (i.e., the gene bearing the sequence complementary to the dsRNA). This effectively knocks out or substantially

reduces the expression of the targeted gene. PTGS can also be accomplished by use of DNA or DNA fragments as well. RNAi methods are described by Fire, A. et al. (1998; Nature 391:806-811) and Gura, T. (2000; Nature 404:804-808). PTGS can also be initiated by introduction of a complementary segment of DNA into the selected tissue using gene delivery and/or viral vector delivery methods described herein or known in the art.

RNAi can be induced in mammalian cells by the use of small interfering RNA also known as siRNA. SiRNA are shorter segments of dsRNA (typically about 21 to 23 nucleotides in length) that result *in vivo* from cleavage of introduced dsRNA by the action of an endogenous ribonuclease. SiRNA appear to be the mediators of the RNAi effect in mammals. The most effective siRNAs appear to be 21 nucleotide dsRNAs with 2 nucleotide 3' overhangs. The use of siRNA for inducing RNAi in mammalian cells is described by Elbashir, S.M. et al. (2001; Nature 411:494-498).

SiRNA can either be generated indirectly by introduction of dsRNA into the targeted cell, or directly by mammalian transfection methods and agents described herein or known in the art (such as liposome-mediated transfection, viral vector methods, or other polynucleotide delivery/introductory methods). Suitable SiRNAs can be selected by examining a transcript of the target polynucleotide (e.g., mRNA) for nucleotide sequences downstream from the AUG start codon and recording the occurrence of each nucleotide and the 3' adjacent 19 to 23 nucleotides as potential siRNA target sites, with sequences having a 21 nucleotide length being preferred. Regions to be avoided for target siRNA sites include the 5' and 3' untranslated regions (UTRs) and regions near the start codon (within 75 bases), as these may be richer in regulatory protein binding sites. UTR-binding proteins and/or translation initiation complexes may interfere with binding of the siRNP endonuclease complex. The selected target sites for siRNA can then be compared to the appropriate genome database (e.g., human, etc.) using BLAST or other sequence comparison algorithms known in the art. Target sequences with significant homology to other coding sequences can be eliminated from consideration. The selected SiRNAs can be produced by chemical synthesis methods known in the art or by *in vitro* transcription using commercially available methods and kits such as the SILENCER siRNA construction kit (Ambion, Austin TX).

In alternative embodiments, long-term gene silencing and/or RNAi effects can be induced in selected tissue using expression vectors that continuously express siRNA. This can be accomplished using expression vectors that are engineered to express hairpin RNAs (shRNAs) using methods known in the art (see, e.g., Brummelkamp, T.R. et al. (2002) Science 296:550-553; and Paddison, P.J. et al. (2002) Genes Dev. 16:948-958). In these and related embodiments, shRNAs can be delivered to target cells using expression vectors known in the art. An example of a suitable expression vector for delivery of siRNA is the PSILENCER1.0-U6 (circular) plasmid (Ambion). Once delivered to the

target tissue, shRNAs are processed *in vivo* into siRNA-like molecules capable of carrying out gene-specific silencing.

In various embodiments, the expression levels of genes targeted by RNAi or PTGS methods can be determined by assays for mRNA and/or protein analysis. Expression levels of the mRNA of a targeted gene, can be determined by northern analysis methods using, for example, the  
5 NORTHERNMAX-GLY kit (Ambion); by microarray methods; by PCR methods; by real time PCR methods; and by other RNA/polynucleotide assays known in the art or described herein. Expression levels of the protein encoded by the targeted gene can be determined by Western analysis using standard techniques known in the art.

10 An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding PMMM. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-  
15 macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased PMMM expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding PMMM may be therapeutically useful, and in the treatment of disorders  
20 associated with decreased PMMM expression or activity, a compound which specifically promotes expression of the polynucleotide encoding PMMM may be therapeutically useful.

In various embodiments, one or more test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in  
25 altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding PMMM is exposed to at least one test compound thus obtained. The sample  
30 may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding PMMM are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding PMMM. The amount of hybridization may be quantified, thus  
35 forming the basis for a comparison of the expression of the polynucleotide both with and without

exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art (Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466).

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of PMMM, antibodies to PMMM, and mimetics, agonists, antagonists, or inhibitors of PMMM.

In various embodiments, the compositions described herein, such as pharmaceutical compositions, may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides



and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery allows administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

5 Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising PMMM or fragments thereof. For example, liposome preparations  
10 containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, PMMM or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

15 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

20 A therapeutically effective dose refers to that amount of active ingredient, for example PMMM or fragments thereof, antibodies of PMMM, and agonists, antagonists or inhibitors of PMMM, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the  
25 dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity.  
30 The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the  
35 severity of the disease state, the general health of the subject, the age, weight, and gender of the

subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## 10 DIAGNOSTICS

In another embodiment, antibodies which specifically bind PMMM may be used for the diagnosis of disorders characterized by expression of PMMM, or in assays to monitor patients being treated with PMMM or agonists, antagonists, or inhibitors of PMMM. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics.

15 Diagnostic assays for PMMM include methods which utilize the antibody and a label to detect PMMM in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

20 A variety of protocols for measuring PMMM, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PMMM expression. Normal or standard values for PMMM expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to PMMM under conditions suitable for complex formation. The amount of standard complex

25 formation may be quantitated by various methods, such as photometric means. Quantities of PMMM expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, polynucleotides encoding PMMM may be used for

30 diagnostic purposes. The polynucleotides which may be used include oligonucleotides, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of PMMM may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of PMMM, and to monitor regulation of PMMM levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotides, including genomic sequences, encoding PMMM or closely related molecules may be used to identify nucleic acid sequences which encode PMMM. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding PMMM, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the PMMM encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:32-62 or from genomic sequences including promoters, enhancers, and introns of the PMMM gene.

Means for producing specific hybridization probes for polynucleotides encoding PMMM include the cloning of polynucleotides encoding PMMM or PMMM-derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotides encoding PMMM may be used for the diagnosis of disorders associated with expression of PMMM. Examples of such disorders include, but are not limited to, a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha<sub>1</sub>-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and

phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; an autoimmune/inflammatory disease, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, atherosclerotic plaque rupture, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, degradation of articular cartilage, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, bone resorption, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, age-related macular degeneration, and sensorineural hearing loss; an epithelial disorder, such as dyshidrotic eczema, allergic contact dermatitis, keratosis pilaris, melasma, vitiligo, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, seborrheic keratosis, folliculitis, herpes simplex, herpes zoster, varicella, candidiasis, dermatophytosis, scabies, insect bites, cherry angioma, keloid, dermatofibroma, acrochordons,

urticaria, transient acantholytic dermatosis, xerosis, eczema, atopic dermatitis, contact dermatitis, hand eczema, nummular eczema, lichen simplex chronicus, asteatotic eczema, stasis dermatitis and stasis ulceration, seborrheic dermatitis, psoriasis, lichen planus, pityriasis rosea, impetigo, ecthyma, dermatophytosis, tinea versicolor, warts, acne vulgaris, acne rosacea, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, bullous pemphigoid, herpes gestationis, dermatitis herpetiformis, linear IgA disease, epidermolysis bullosa acquisita, dermatomyositis, lupus erythematosus, scleroderma and morphea, erythroderma, alopecia, figurate skin lesions, telangiectasias, hypopigmentation, hyperpigmentation, vesicles/bullae, exanthems, cutaneous drug reactions, papulonodular skin lesions, chronic non-healing wounds, photosensitivity diseases, epidermolysis bullosa simplex, epidermolytic hyperkeratosis, epidermolytic and nonepidermolytic palmoplantar keratoderma, ichthyosis bullosa of Siemens, ichthyosis exfoliativa, keratosis palmaris et plantaris, keratosis palmoplantaris, palmoplantar keratoderma, keratosis punctata, Meesmann's corneal dystrophy, pachyonychia congenita, white sponge nevus, steatocystoma multiplex, epidermal nevi/epidermolytic hyperkeratosis type, monilethrix, trichothiodystrophy, chronic hepatitis/cryptogenic cirrhosis, and colorectal hyperplasia; a neurological disorder, such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a reproductive disorder, such as infertility, including tubal disease, ovulatory defects, and endometriosis, a disorder of prolactin production, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation

syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an endocrine disorder such as a disorder of the hypothalamus and/or pituitary resulting from lesions such as a primary brain tumor, adenoma, infarction associated with pregnancy, hypophysectomy, aneurysm, vascular malformation, thrombosis, infection, immunological disorder, and complication due to head trauma; a disorder associated with hypopituitarism including hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease, Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism; a disorder associated with hyperpituitarism including acromegaly, gigantism, and syndrome of inappropriate antidiuretic hormone (ADH) secretion (SIADH) often caused by benign adenoma; a disorder associated with hypothyroidism including goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis (Hashimoto's disease), and cretinism; a disorder associated with hyperthyroidism including thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter, thyroid carcinoma, and Plummer's disease; a disorder associated with hyperparathyroidism including Conn disease (chronic hypercalcemia); a metabolic disorder such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, mannosidosis, neuraminidase deficiency, obesity, pentosuria phenylketonuria, pseudovitamin D-deficiency rickets; a disorder of carbohydrate metabolism such as congenital type II dyserythropoietic anemia, diabetes, insulin-dependent diabetes mellitus, non-insulin-dependent diabetes mellitus, fructose-1,6-diphosphatase deficiency, galactosemia, glucagonoma, hereditary fructose intolerance, hypoglycemia, mannosidosis, neuraminidase deficiency, obesity, galactose epimerase deficiency, glycogen storage diseases, lysosomal storage diseases, fructosuria, pentosuria, and inherited abnormalities of pyruvate metabolism; a disorder of lipid metabolism such as fatty liver, cholestasis, primary biliary cirrhosis, carnitine deficiency, carnitine palmitoyltransferase deficiency, myoadenylate deaminase deficiency, hypertriglyceridemia, lipid storage disorders such as Fabry's disease, Gaucher's disease, Niemann-Pick's disease, metachromatic leukodystrophy, adrenoleukodystrophy, GM<sub>2</sub> gangliosidosis, and ceroid lipofuscinosis, abetalipoproteinemia, Tangier disease, hyperlipoproteinemia, diabetes mellitus,

lipodystrophy, lipomatoses, acute panniculitis, disseminated fat necrosis, adiposis dolorosa, lipoid adrenal hyperplasia, minimal change disease, lipomas, atherosclerosis, hypercholesterolemia, hypercholesterolemia with hypertriglyceridemia, primary hypoalphalipoproteinemia, hypothyroidism, renal disease, liver disease, lecithin:cholesterol acyltransferase deficiency, cerebrotendinous

5 xanthomatosis, sitosterolemia, hypocholesterolemia, Tay-Sachs disease, Sandhoff's disease, hyperlipidemia, hyperlipemia, lipid myopathies, and obesity; and a disorder of copper metabolism such as Menke's disease, Wilson's disease, and Ehlers-Danlos syndrome type IX; a pancreatic disorder such as Type I or Type II diabetes mellitus and associated complications; a disorder associated with the adrenals such as hyperplasia, carcinoma, or adenoma of the adrenal cortex,

10 hypertension associated with alkalosis, amyloidosis, hypokalemia, Cushing's disease, Liddle's syndrome, and Arnold-Healy-Gordon syndrome, pheochromocytoma tumors, and Addison's disease; a disorder associated with gonadal steroid hormones such as: in women, abnormal prolactin production, infertility, endometriosis, perturbation of the menstrual cycle, polycystic ovarian disease, hyperprolactinemia, isolated gonadotropin deficiency, amenorrhea, galactorrhea, hermaphroditism,

15 hirsutism and virilization, breast cancer, and, in post-menopausal women, osteoporosis; and, in men, Leydig cell deficiency, male climacteric phase, and germinal cell aplasia, a hypergonadal disorder associated with Leydig cell tumors, androgen resistance associated with absence of androgen receptors, syndrome of 5  $\alpha$ -reductase, and gynecomastia; a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal

20 gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an infection caused by a viral agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus,

25 reovirus, retrovirus, rhabdovirus, or togavirus; an infection caused by a bacterial agent classified as pneumococcus, staphylococcus, streptococcus, bacillus, corynebacterium, clostridium, meningococcus, gonococcus, listeria, moraxella, kingella, haemophilus, legionella, bordetella, gram-negative enterobacterium including shigella, salmonella, or campylobacter, pseudomonas, vibrio, brucella, francisella, yersinia, bartonella, norcardium, actinomyces, mycobacterium, spirochaetale,

30 rickettsia, chlamydia, or mycoplasma; an infection caused by a fungal agent classified as aspergillus, blastomyces, dermatophytes, cryptococcus, coccidioides, malassezia, histoplasma, or other mycosis-causing fungal agent; and an infection caused by a parasite classified as plasmodium or malaria-causing, parasitic entamoeba, leishmania, trypanosoma, toxoplasma, pneumocystis carinii, intestinal protozoa such as giardia, trichomonas, tissue nematode such as trichinella, intestinal nematode such

35 as ascaris, lymphatic filarial nematode, trematode such as schistosoma, and cestode such as

tapeworm. Polynucleotides encoding PMMM may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered PMMM expression. Such qualitative or quantitative methods are well known in the art.

5 In a particular embodiment, polynucleotides encoding PMMM may be used in assays that detect the presence of associated disorders, particularly those mentioned above. Polynucleotides complementary to sequences encoding PMMM may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and  
10 compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of polynucleotides encoding PMMM in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

15 In order to provide a basis for the diagnosis of a disorder associated with expression of PMMM, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PMMM, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from  
20 normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated,  
25 hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or  
30 overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier, thereby preventing the development or further progression of the cancer.



Additional diagnostic uses for oligonucleotides designed from the sequences encoding PMMM may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding PMMM, or a fragment of a polynucleotide complementary to the polynucleotide encoding PMMM, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from polynucleotides encoding PMMM may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from polynucleotides encoding PMMM are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

SNPs may be used to study the genetic basis of human disease. For example, at least 16 common SNPs have been associated with non-insulin-dependent diabetes mellitus. SNPs are also useful for examining differences in disease outcomes in monogenic disorders, such as cystic fibrosis, sickle cell anemia, or chronic granulomatous disease. For example, variants in the mannose-binding lectin, MBL2, have been shown to be correlated with deleterious pulmonary outcomes in cystic fibrosis. SNPs also have utility in pharmacogenomics, the identification of genetic variants that influence a patient's response to a drug, such as life-threatening toxicity. For example, a variation in N-acetyl transferase is associated with a high incidence of peripheral neuropathy in response to the anti-tuberculosis drug isoniazid, while a variation in the core promoter of the ALOX5 gene results in diminished clinical response to treatment with an anti-asthma drug that targets the 5-lipoxygenase

pathway. Analysis of the distribution of SNPs in different populations is useful for investigating genetic drift, mutation, recombination, and selection, as well as for tracing the origins of populations and their migrations (Taylor, J.G. et al. (2001) *Trends Mol. Med.* 7:507-512; Kwok, P.-Y. and Z. Gu (1999) *Mol. Med. Today* 5:538-543; Nowotny, P. et al. (2001) *Curr. Opin. Neurobiol.* 11:637-641).

5       Methods which may also be used to quantify the expression of PMMM include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves (Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is  
10       presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

      In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotides described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers  
15       of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used  
20       to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

      In another embodiment, PMMM, fragments of PMMM, or antibodies specific for PMMM  
25       may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

      A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of  
30       gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time (Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484; hereby expressly incorporated by reference herein). Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of  
35       transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the

hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression *in vivo*, as in the case of a tissue or biopsy sample, or *in vitro*, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with *in vitro* model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity (see, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>). Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In an embodiment, the toxicity of a test compound can be assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another embodiment relates to the use of the polypeptides disclosed herein to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected

individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using  
5 two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot  
10 is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed  
15 by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of interest. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for PMMM to quantify the levels of PMMM expression. In one embodiment, the antibodies are used as elements on a  
20 microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoz, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at  
25 each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be  
30 useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological  
35 sample containing proteins with the test compound. Proteins that are expressed in the treated

biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art (Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662). Various types of microarrays are well known and thoroughly described in Schena, M., ed. (1999; DNA Microarrays: A Practical Approach, Oxford University Press, London).

In another embodiment of the invention, nucleic acid sequences encoding PMMM may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries (Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; Trask, B.J. (1991) Trends Genet. 7:149-154). Once mapped, the nucleic acid sequences may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP) (Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357).

Fluorescent *in situ* hybridization (FISH) may be correlated with other physical and genetic map data (Heinz-Ulrich, et al. (1995) in Meyers, *supra*, pp. 965-968). Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding PMMM on a physical  
5 map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

*In situ* hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse,  
10 may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further  
15 investigation (Gatti, R.A. et al. (1988) Nature 336:577-580). The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, PMMM, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug  
20 screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between PMMM and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest (Geysen, et al. (1984) PCT application  
25 WO84/03564). In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with PMMM, or fragments thereof, and washed. Bound PMMM is then detected by methods well known in the art. Purified PMMM can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

30 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PMMM specifically compete with a test compound for binding PMMM. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PMMM.

In additional embodiments, the nucleotide sequences which encode PMMM may be used in  
35 any molecular biology techniques that have yet to be developed, provided the new techniques rely on

properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, including U.S. Ser. No. 60/322,196, U.S. Ser. No. 60/324,134, U.S. Ser. No. 60/327,233, U.S. Ser. No. 60/332,423, U.S. Ser. No. 60/334,145, U.S. Ser. No. 60/334,229, U.S. Ser. No. 60/337,451, U.S. Ser. No. 60/343,980, U.S. Ser. No. 60/346,198, U.S. Ser. No. 60/348,887, U.S. Ser. No. 60/351,928, and U.S. Ser. No. 60/366,837, are hereby expressly incorporated by reference.

## EXAMPLES

### I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA). Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Invitrogen), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Invitrogen), using the recommended procedures or similar methods known in the art (Ausubel et al., *supra*, ch. 5). Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Biosciences) or

preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Invitrogen, Carlsbad CA), PCDNA2.1 plasmid (Invitrogen), PBK-CMV plasmid (Stratagene), PCR2-TOPOTA plasmid (Invitrogen), PCMV-ICIS plasmid (Stratagene), pIGEN (Incyte Genomics, Palo Alto CA), pRARE (Incyte Genomics), or pINCY (Incyte Genomics), or derivatives thereof.

Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Invitrogen.

## II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

## III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Biosciences or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Amersham Biosciences); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (Ausubel et al., *supra*, ch.



7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM; PROTEOME databases with sequences from *Homo sapiens*, *Rattus norvegicus*, *Mus musculus*, *Caenorhabditis elegans*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Candida albicans* (Incyte Genomics, Palo Alto CA); hidden Markov model (HMM)-based protein family databases such as PFAM, INCY, and TIGRFAM (Haft, D.H. et al. (2001) Nucleic Acids Res. 29:41-43); and HMM-based protein domain databases such as SMART (Schultz, J. et al. (1998) Proc. Natl. Acad. Sci. USA 95:5857-5864; Letunic, I. et al. (2002) Nucleic Acids Res. 30:242-244). (HMM is a probabilistic approach which analyzes consensus primary structures of gene families; see, for example, Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, the PROTEOME databases, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, hidden Markov model (HMM)-based protein family databases such as PFAM, INCY, and TIGRFAM; and HMM-based protein domain databases such as SMART. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (MiraiBio, Alameda CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and

threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:32-62. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 2.

#### IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative protein modification and maintenance molecules were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpr and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94; Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode protein modification and maintenance molecules, the encoded polypeptides were analyzed by querying against PFAM models for protein modification and maintenance molecules. Potential protein modification and maintenance molecules were also identified by homology to Incyte cDNA sequences that had been annotated as protein modification and maintenance molecules. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpr public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

#### V. Assembly of Genomic Sequence Data with cDNA Sequence Data

### **"Stitched" Sequences**

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpri public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

### **"Stretched" Sequences**

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

## **VI. Chromosomal Mapping of PMMM Encoding Polynucleotides**

The sequences which were used to assemble SEQ ID NO:32-62 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:32-62 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

## VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook and Russell, *supra*, ch. 7; Ausubel et al., *supra*, ch. 4).

Analogous computer techniques applying BLAST were used to search for identical or related molecules in databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum}\{\text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2})\}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is

calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotides encoding PMMM are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding PMMM. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

#### **VIII. Extension of PMMM Encoding Polynucleotides**

Full length polynucleotides are produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and 2-mercaptoethanol, Taq DNA polymerase (Amersham Biosciences), ELONGASE enzyme (Invitrogen), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Biosciences). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Biosciences), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Biosciences) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries

were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Biosciences) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

- 5 In like manner, full length polynucleotides are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### IX. Identification of Single Nucleotide Polymorphisms in PMMM Encoding Polynucleotides

- Common DNA sequence variants known as single nucleotide polymorphisms (SNPs) were identified in SEQ ID NO:32-62 using the LIFESEQ database (Incyte Genomics). Sequences from the same gene were clustered together and assembled as described in Example III, allowing the identification of all sequence variants in the gene. An algorithm consisting of a series of filters was used to distinguish SNPs from other sequence variants. Preliminary filters removed the majority of basecall errors by requiring a minimum Phred quality score of 15, and removed sequence alignment errors and errors resulting from improper trimming of vector sequences, chimeras, and splice variants. An automated procedure of advanced chromosome analysis analysed the original chromatogram files in the vicinity of the putative SNP. Clone error filters used statistically generated algorithms to identify errors introduced during laboratory processing, such as those caused by reverse transcriptase, polymerase, or somatic mutation. Clustering error filters used statistically generated algorithms to identify errors resulting from clustering of close homologs or pseudogenes, or due to contamination by non-human sequences. A final set of filters removed duplicates and SNPs found in immunoglobulins or T-cell receptors.
- 10  
15  
20

- Certain SNPs were selected for further characterization by mass spectrometry using the high throughput MASSARRAY system (Sequenom, Inc.) to analyze allele frequencies at the SNP sites in four different human populations. The Caucasian population comprised 92 individuals (46 male, 46 female), including 83 from Utah, four French, three Venezuelan, and two Amish individuals. The African population comprised 194 individuals (97 male, 97 female), all African Americans. The Hispanic population comprised 324 individuals (162 male, 162 female), all Mexican Hispanic. The Asian population comprised 126 individuals (64 male, 62 female) with a reported parental breakdown of 43% Chinese, 31% Japanese, 13% Korean, 5% Vietnamese, and 8% other Asian. Allele frequencies were first analyzed in the Caucasian population; in some cases those SNPs which showed no allelic variance in this population were not further tested in the other three populations.
- 25  
30

#### X. Labeling and Use of Individual Hybridization Probes

- Hybridization probes derived from SEQ ID NO:32-62 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base
- 35

pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ - $^{32}$ P] adenosine triphosphate (Amersham Biosciences), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Biosciences). An aliquot containing  $10^7$  counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

#### **XI. Microarrays**

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing; see, e.g., Baldeschweiler et al., *supra*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schena, M., ed. (1999) DNA Microarrays: A Practical Approach, Oxford University Press, London). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements (Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645; Marshall, A. and J. Hodgson (1998) *Nat. Biotechnol.* 16:27-31).

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the



biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### **Tissue or Cell Sample Preparation**

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ $\mu$ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ $\mu$ l RNase inhibitor, 500  $\mu$ M dATP, 500  $\mu$ M dGTP, 500  $\mu$ M dTTP, 40  $\mu$ M dCTP, 40  $\mu$ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Biosciences). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with GEMBRIGHT kits (Incyte Genomics). Specific control poly(A)<sup>+</sup> RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (Clontech, Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14  $\mu$ l 5X SSC/0.2% SDS.

#### **Microarray Preparation**

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5  $\mu$ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Biosciences).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in

distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in U.S. Patent No. 5,807,522, incorporated herein by reference. 1  $\mu$ l of the array element DNA, at an  
5 average concentration of 100 ng/ $\mu$ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water.  
10 Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60° C followed by washes in 0.2% SDS and distilled water as before.

#### Hybridization

Hybridization reactions contain 9  $\mu$ l of sample mixture consisting of 0.2  $\mu$ g each of Cy3  
15 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65° C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140  $\mu$ l of 5X SSC in a corner of the chamber. The chamber  
20 containing the arrays is incubated for about 6.5 hours at 60° C. The arrays are washed for 10-min at 45° C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45° C in a second wash buffer (0.1X SSC), and dried.

#### Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with  
25 an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the  
30 present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two

fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000.

When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte Genomics). Array elements that exhibit at least about a two-fold change in expression, a signal-to-background ratio of at least about 2.5, and an element spot size of at least about 40%, are considered to be differentially expressed.

### Expression

For example, SEQ ID NO:40 showed decreased expression in peripheral blood mononuclear cells (PBMCs) treated with PMA and ionomycin versus untreated PBMCs as determined by microarray analysis. Peripheral blood mononuclear cells (PBMCs) are isolated from freshly obtained peripheral blood. PBMCs are stimulated *in vitro* with soluble PMA and

ionomycin for 1, 2, 4, 8, and 20 hours. These treated cells are compared to untreated PBMCs kept in culture. Therefore, in various embodiments, SEQ ID NO:40 can be used for one or more of the following: i) monitoring treatment of immune disorders and related diseases and conditions, ii) diagnostic assays for immune disorders and related diseases and conditions, and iii) developing therapeutics and/or other treatments for immune disorders and related diseases and conditions.

In another example, SEQ ID NO:43 was differentially expressed in human breast tumor cell lines as compared to a nonmalignant breast epithelial cell line, MCF-10A. Histological and molecular evaluation of breast tumors reveals that the development of breast cancer evolves through a multi-step process whereby pre-malignant mammary epithelial cells undergo a relatively defined sequence of events leading to tumor formation. An early event in tumor development is ductal hyperplasia. Cells undergoing rapid neoplastic growth gradually progress to invasive carcinoma and become metastatic to the lung, bone, and potentially other organs. Several variables that may influence the process of tumor progression and malignant transformation include genetic factors, environmental factors, growth factors, and hormones. Based on the complexity of this process, it is critical to study a population of human mammary epithelial cells undergoing the process of malignant transformation, and to associate specific stages of progression with phenotypic and molecular characteristics. In a cross-comparison study, two cell lines out of nine tested exhibited differential expression as compared to controls. BT-20 is a breast carcinoma cell line derived *in vitro* from cells emigrating out of thin slices of the tumor mass isolated from a 74-year old female. MDA-mb-435S is a spindle shaped strain derived from the pleural effusion of a 31-year old female with metastatic, ductal adenocarcinoma of the breast. In this experiment, the expression of SEQ ID NO:43 was increased by at least two-fold in these breast tumor cell lines. Therefore, in various embodiments, SEQ ID NO:43 can be used for one or more of the following: i) monitoring treatment of breast cancer, ii) diagnostic assays for breast cancer, and iii) developing therapeutics and/or other treatments for breast cancer.

In another example, SEQ ID NO:43-44 were differentially expressed in three separate experiments in which human lung tumor cells were tested in a pair comparison with normal lung from the same donor. Lung cancers are divided into four histopathologically distinct groups. Three groups (squamous cell carcinoma, adenocarcinoma, and large cell carcinoma) are classified as non-small cell lung cancers (NSCLCs). The fourth group of cancers is referred to as small cell lung cancer (SCLC). Collectively, NSCLCs account for approximately 70% of cases while SCLCs account for approximately 18% of cases. The molecular and cellular biology underlying the development and progression of lung cancer are incompletely understood. Deletions on chromosome 3 are common in this disease and are thought to indicate the presence of a tumor suppressor gene in this region. Activating mutations in K-ras are commonly found in lung cancer and are the basis of one of the

mouse models for the disease. Analysis of gene expression patterns associated with the development and progression of the disease will yield tremendous insight into the biology underlying this disease, and will lead to the development of improved diagnostics and therapeutics. In these experiments, the expression of SEQ ID NO:43-44 were increased by at least two-fold in the lung tumor cells as compared to the normal lung tissue cells from the same donor.

These experiments indicate that SEQ ID NO:43 and SEQ ID NO:44 exhibited significant differential expression patterns using microarray techniques. Therefore, in various embodiments, SEQ ID NO:43-44 can be used for one or more of the following: i) monitoring treatment of lung cancer, ii) diagnostic assays for lung cancer, and iii) developing therapeutics and/or other treatments for lung cancer.

In another example, SEQ ID NO:45 was differentially expressed in human breast tumor cell lines compared to nonmalignant breast epithelial cell lines. Histological and molecular evaluation of breast tumors reveals that the development of breast cancer evolves through a multi-step process whereby pre-malignant mammary epithelial cells undergo a relatively defined sequence of events leading to tumor formation. An early event in tumor development is ductal hyperplasia. Cells undergoing rapid neoplastic growth gradually progress to invasive carcinoma and become metastatic to the lung, bone, and potentially other organs. Several variables that may influence the process of tumor progression and malignant transformation include genetic factors, environmental factors, growth factors, and hormones. Based on the complexity of this process, it is critical to study a population of human mammary epithelial cells undergoing the process of malignant transformation.

In one set of experiments, human primary epithelial breast cells (HMECs) isolated from a normal donor were compared to various types of breast cancer cell lines. Of six breast cancer cell lines tested, two of these cell lines, MCF-7 (breast adenocarcinoma) and SK-BR-3 (human breast adenocarcinoma, which is also tumorigenic in nude mice) were underexpressed in SEQ ID NO:45 by at least two-fold as compared to HMEC cells.

SEQ ID NO:45 was also underexpressed by at least two-fold in MCF-7 breast adenocarcinoma cells as compared to nonmalignant MCF10A cells isolated from normal breast epithelial tissue.

These experiments indicate that SEQ ID NO:45 exhibits significant differential expression patterns using microarray techniques. Therefore, in various embodiments, SEQ ID NO:45 can be used for one or more of the following: i) monitoring treatment of breast cancer, ii) diagnostic assays for breast cancer, and iii) developing therapeutics and/or other treatments for breast cancer.

In another example, SEQ ID NO:49 showed differential expression in breast cancer tissue, as determined by microarray analysis. In order to better determine the molecular and phenotypic characteristics associated with different stages of breast cancer, breast carcinoma cell lines at various

stages of tumor progression were compared to primary human breast epithelial cells. The breast carcinoma cell lines include MCF7, a breast adenocarcinoma cell line derived from the pleural effusion of a 69-year-old female; Sk-BR-3, a breast adenocarcinoma cell line isolated from a malignant pleural effusion of a 43-year-old female; and BT-20, a breast adenocarcinoma isolated in vitro from cells emigrating out of thin slices of a tumor mass isolated from a 74-year-old female. The primary mammary epithelial cell line HMEC was derived from normal human mammary tissue (Clonetics, San Diego, CA). All cell cultures were propagated in a chemically-defined medium, according to the supplier's recommendations and grown to 70-80% confluence prior to RNA isolation. The microarray experiments showed that expression of SEQ ID NO:49 was decreased by at least two fold in all three breast carcinoma lines (MCF7, Sk-BR-3, and BT20) relative to primary mammary epithelial cells. Therefore, in various embodiments, SEQ ID NO:49 can be used for one or more of the following: i) monitoring treatment of breast cancer, ii) diagnostic assays for breast cancer, and iii) developing therapeutics and/or other treatments for breast cancer.

SEQ ID NO:49 also showed differential expression, as determined by microarray analysis, in liver C3A cells treated with one of the following steroids: beclomethasone, dexamethasone, progesterone, budesonide. The human C3A cell line is a clonal derivative of HepG2/C3 and has been established as an *in vitro* model of the mature human liver (Mickelson et al. (1995) Hepatology 22:866-875; Nagendra et al. (1997) Am J Physiol 272:G408-G416). SEQ ID NO:5 showed at least a two-fold decrease in expression at a minimum of two out of the three time points in early confluent C3A cells treated with beclomethasone, budesonide, dexamethasone, or betamethasone, for 1, 3, or 6 hours. These experiments indicate that SEQ ID NO:49 is useful in diagnostic assays for liver diseases and as a potential biological marker and therapeutic agent in the treatment of liver diseases and disorders. Therefore, in various embodiments, SEQ ID NO:49 can be used for one or more of the following: i) monitoring treatment of liver diseases and disorders, ii) diagnostic assays for liver diseases and disorders, and iii) developing therapeutics and/or other treatments for liver diseases and disorders.

In another example, SEQ ID NO:51 showed differential expression, as determined by microarray analysis, in Alzheimer's Disease (AD). In a comparison of cerebellum tissue from a 76-year-old male with severe AD to cerebellum tissue from a normal 67-year-old male, the expression of SEQ ID NO:51 was decreased at least two-fold. Therefore, in various embodiments, SEQ ID NO:51 can be used for one or more of the following: i) monitoring treatment of Alzheimer's Disease, ii) diagnostic assays for Alzheimer's Disease, and iii) developing therapeutics and/or other treatments for Alzheimer's Disease.

SEQ ID NO:51 also showed differential expression associated with colon cancer, as determined by microarray analysis. Normal colon tissue was compared to colon tumor tissue from a

67-year-old donor with moderately differentiated adenocarcinoma. The expression of SEQ ID NO:51 was decreased at least two-fold in the tumor tissue as compared to the normal tissue. Therefore, in various embodiments, SEQ ID NO:51 can be used for one or more of the following: i) monitoring treatment of colon cancer, ii) diagnostic assays for colon cancer, and iii) developing therapeutics and/or other treatments for colon cancer.

In another example, the expression of SEQ ID NO:56 in a primary prostate epithelial cell line isolated from a normal donor, PrEC, was compared to that in three prostate carcinoma cell lines. DU 145 is a prostate carcinoma cell line isolated from a metastatic site in the brain of a 69 year old male with widespread metastatic prostate carcinoma. DU 145 has no detectable sensitivity to hormones; forms colonies in semi-solid medium, is only weakly positive for acid phosphatase, and is negative for prostate specific antigen. LNCaP is a prostate carcinoma cell line isolated from a lymph node biopsy of a 50 year old male with metastatic prostate carcinoma. LNCaP cells express prostate specific antigens, produce prostatic acid phosphatase, and express androgen receptors. PC-3 is a prostate adenocarcinoma cell line isolated from a metastatic site in the bone of a 62 year old male with grade IV prostate adenocarcinoma. The expression of SEQ ID NO:56 was increased by at least two-fold in DU 145 cells grown under restrictive conditions as compared to PrEC cells grown under restrictive conditions. Therefore, in various embodiments, SEQ ID NO:56 can be used for one or more of the following: i) monitoring treatment of prostate cancer, ii) diagnostic assays for prostate cancer, and iii) developing therapeutics and/or other treatments for prostate cancer.

In another example, SEQ ID NO:58, SEQ ID NO:59 and SEQ ID NO:60 showed differential expression associated with breast cancer, as determined by microarray analysis. The gene expression profile of a nonmalignant mammary epithelial cell line was compared to the gene expression profiles of breast carcinoma lines at different stages of tumor progression. Cell lines compared included: a) BT-20, a breast carcinoma cell line derived *in vitro* from the cells emigrating out of thin slices of tumor mass isolated from a 74-year-old female, b) BT-474, a breast ductal carcinoma cell line that was isolated from a solid, invasive ductal carcinoma of the breast obtained from a 60-year-old woman, c) BT-483, a breast ductal carcinoma cell line that was isolated from a papillary invasive ductal tumor obtained from a 23-year-old normal, menstruating, parous female with a family history of breast cancer, d) Hs578T, a breast ductal carcinoma cell line isolated from a 74-year-old female with breast carcinoma, e) MCF7, a nonmalignant breast adenocarcinoma cell line isolated from the pleural effusion of a 69-year-old female, f) MCF-10A, a breast mammary gland (luminal ductal characteristics) cell line isolated from a 36-year-old woman with fibrocystic breast disease, g) MDA-mb-435S, a spindle-shaped strain that evolved from the parent line (435) isolated by R. Cailleau from pleural effusion of a 31-year-old female with metastatic, ductal adenocarcinoma of the breast, h) Sk-BR-3, a breast adenocarcinoma cell line isolated from a malignant pleural effusion of a 43-year-old

female, i) T-47D, a breast carcinoma cell line isolated from a pleural effusion obtained from a 54-year-old female with an infiltrating ductal carcinoma of the breast and j) HMEC, a primary breast epithelial cell line isolated from a normal donor. SEQ ID NO:58 expression was reduced by at least two-fold in BT20 and MCF7 cells as compared to HMEC cells. The expression of SEQ ID NO:59 was decreased by at least two-fold in carcinoma cell lines BT20, Sk-BR-3, T-47D, MDA-mb-435S and MCF7 as compared to HMEC cells. SEQ ID NO:60 expression was upregulated by at least two-fold in the carcinoma cell line Hs578T as compared to the HMEC cell line. Therefore, in various embodiments, SEQ ID NO:58, SEQ ID NO:59 and SEQ ID NO:60 can be used for one or more of the following: i) monitoring treatment of breast cancer, ii) diagnostic assays for breast cancer, and iii) developing therapeutics and/or other treatments for breast cancer.

In another example, SEQ ID NO:60 showed differential expression associated with lung cancer, as determined by microarray analysis. Expression was compared in matched samples of normal and lung tumor tissue from individual donors. Tissue samples were provided by the Roy Castle International Centre for Lung Cancer Research. SEQ ID NO:60 expression was upregulated by at least two-fold in lung squamous cell carcinoma tissue derived from a 68-year-old female donor as compared to normal lung tissue from the same donor. Therefore, in various embodiments, SEQ ID NO:60 can be used for one or more of the following: i) monitoring treatment of lung cancer, ii) diagnostic assays for lung cancer, and iii) developing therapeutics and/or other treatments for lung cancer.

In another example, SEQ ID NO:58 and SEQ ID NO:59 showed differential expression associated with ovarian cancer, as determined by microarray analysis. A normal ovary from a 79 year-old female donor was compared to an ovarian tumor from the same donor (Huntsman Cancer Institute, Salt Lake City, UT). The expression of SEQ ID NO:58 and SEQ ID NO:59 was decreased by at least two-fold in the tumor tissue as compared to the normal tissue. Therefore, SEQ ID NO:58 and SEQ ID NO:59 are useful in monitoring treatment of, and diagnostic assays for ovarian cancer. Therefore, in various embodiments, SEQ ID NO:58-59 can be used for one or more of the following: i) monitoring treatment of ovarian cancer, ii) diagnostic assays for ovarian cancer, and iii) developing therapeutics and/or other treatments for ovarian cancer.

In another example, SEQ ID NO:59 showed differential expression associated with steroid hormone responses, as determined by microarray analysis. The human C3A cell line is a clonal derivative of HepG2/C3 (hepatoma cell line, isolated from a 15-year-old male with liver tumor), which was selected for strong contact inhibition of growth. The use of a clonal population enhances the reproducibility of the cells. C3A cells have many characteristics of primary human hepatocytes in culture: i) expression of insulin receptor and insulin-like growth factor II receptor; ii) secretion of a high ratio of serum albumin compared with  $\alpha$ -fetoprotein iii) conversion of ammonia to urea and



glutamine; iv) metabolism of aromatic amino acids; and v) proliferation in glucose-free and insulin-free medium. The C3A cell line is now well established as an in vitro model of the mature human liver (Mickelson et al. (1995) Hepatology 22:866-875; Nagendra et al. (1997) Am J Physiol 272:G408-G416). Early Confluent C3A cells were treated with progesterone or budenoside at 1, 10, and 100  $\mu$ M for 1, 3, and 6 hours. The treated cells were compared to untreated early confluent C3A cells. At each of the time points, the expression of SEQ ID NO:59 was decreased by at least two-fold in C3A cells treated with 10 or 100  $\mu$ M budenoside, and in C3A cells treated with 10  $\mu$ M progesterone. Therefore, SEQ ID NO:59 may be useful in monitoring of, and diagnostic assays for steroid hormone-induced responses. Therefore, in various embodiments, SEQ ID NO:59 can be used for one or more of the following: i) monitoring treatment of steroid hormone-induced responses, ii) diagnostic assays for steroid hormone-induced responses, and iii) developing therapeutics and/or other treatments for steroid hormone-induced responses.

In another example, SEQ ID NO:61 showed differential expression associated with lung cancer, as determined by microarray analysis. Pair comparisons of lung tumor tissue and microscopically-normal tissue from the same donor were made. The expression of SEQ ID NO:61 was increased by at least two-fold in lung squamous cell carcinoma tissue from a 68 year-old female as compared to normal lung tissue from the same donor (Roy Castle International Centre for Lung Cancer Research, Liverpool, UK). Therefore, in various embodiments, SEQ ID NO:61 can be used for one or more of the following: i) monitoring treatment of lung cancer, ii) diagnostic assays for lung cancer, and iii) developing therapeutics and/or other treatments for lung cancer.

## **XII. Complementary Polynucleotides**

Sequences complementary to the PMMM-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring PMMM. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of PMMM. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the PMMM-encoding transcript.

## **XIII. Expression of PMMM**

Expression and purification of PMMM is achieved using bacterial or virus-based expression systems. For expression of PMMM in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid

promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

Antibiotic resistant bacteria express PMMM upon induction with isopropyl beta-D-

thiogalactopyranoside (IPTG). Expression of PMMM in eukaryotic cells is achieved by infecting

5 insect or mammalian cell lines with recombinant *Autographica californica* nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding PMMM by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to  
10 infect *Spodoptera frugiperda* (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus (Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945).

In most expression systems, PMMM is synthesized as a fusion protein with, e.g., glutathione  
15 S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from *Schistosoma japonicum*, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Biosciences). Following purification, the GST moiety can be proteolytically cleaved from PMMM at  
20 specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel et al. (*supra*, ch. 10 and 16). Purified PMMM obtained by these methods can be used directly in the assays shown  
25 in Examples XVII, XVIII, XIX, and XX, where applicable.

#### XIV. Functional Assays

PMMM function is assessed by expressing the sequences encoding PMMM at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA  
30 expression. Vectors of choice include PCMV SPORT plasmid (Invitrogen, Carlsbad CA) and PCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of  
35 a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a

reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994; Flow Cytometry, Oxford, New York NY).

The influence of PMMM on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding PMMM and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding PMMM and other genes of interest can be analyzed by northern analysis or microarray techniques.

#### **XV. Production of PMMM Specific Antibodies**

PMMM substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize animals (e.g., rabbits, mice, etc.) and to produce antibodies using standard protocols.

Alternatively, the PMMM amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art (Ausubel et al., *supra*, ch. 11).

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity (Ausubel et al., *supra*). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide and anti-PMMM

activity by, for example, binding the peptide or PMMM to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

#### **XVI. Purification of Naturally Occurring PMMM Using Specific Antibodies**

Naturally occurring or recombinant PMMM is substantially purified by immunoaffinity chromatography using antibodies specific for PMMM. An immunoaffinity column is constructed by covalently coupling anti-PMMM antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Biosciences). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing PMMM are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PMMM (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/PMMM binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and PMMM is collected.

#### **XVII. Identification of Molecules Which Interact with PMMM**

PMMM, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent (Bolton, A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539). Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled PMMM, washed, and any wells with labeled PMMM complex are assayed. Data obtained using different concentrations of PMMM are used to calculate values for the number, affinity, and association of PMMM with the candidate molecules.

Alternatively, molecules interacting with PMMM are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989; *Nature* 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

PMMM may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

#### **XVIII. Demonstration of PMMM Activity**

PMMM activity can be demonstrated using a generic immunoblotting strategy or through a variety of specific activity assays, some of which are outlined below. As a general approach, cell lines or tissues transformed with a vector containing PMMM coding sequences can be assayed for PMMM activity by immunoblotting. Transformed cells are denatured in SDS in the presence of β-mercaptoethanol, nucleic acids are removed by ethanol precipitation, and proteins are purified by acetone precipitation. Pellets are resuspended in 20 mM Tris buffer at pH 7.5 and incubated with Protein G-Sepharose pre-coated with an antibody specific for PMMM. After washing, the Sepharose

beads are boiled in electrophoresis sample buffer, and the eluted proteins subjected to SDS-PAGE. The SDS-PAGE is transferred to a membrane for immunoblotting, and the PMMM activity is assessed by visualizing and quantifying bands on the blot using the antibody specific for PMMM as the primary antibody and  $^{125}\text{I}$ -labeled IgG specific for the primary antibody as the secondary antibody.

5 PMMM kinase activity is measured by quantifying the phosphorylation of a protein substrate by PMMM in the presence of gamma-labeled  $^{32}\text{P}$ -ATP. PMMM is incubated with the protein substrate,  $^{32}\text{P}$ -ATP, and an appropriate kinase buffer. The  $^{32}\text{P}$  incorporated into the substrate is separated from free  $^{32}\text{P}$ -ATP by electrophoresis and the incorporated  $^{32}\text{P}$  is counted using a radioisotope counter. The amount of incorporated  $^{32}\text{P}$  is proportional to the activity of PMMM. A  
10 determination of the specific amino acid residue phosphorylated is made by phosphoamino acid analysis of the hydrolyzed protein.

In one alternative, PMMM activity is demonstrated by a test for galactosyltransferase activity. This can be determined by measuring the transfer of radiolabeled galactose from UDP-galactose to a GlcNAc-terminated oligosaccharide chain (Kolbinger, F. et al. (1998) J. Biol. Chem.  
15 273:58-65). The sample is incubated with 14  $\mu\text{l}$  of assay stock solution (180 mM sodium cacodylate, pH 6.5, 1 mg/ml bovine serum albumin, 0.26 mM UDP-galactose, 2  $\mu\text{l}$  of UDP- $^3\text{H}$ galactose), 1  $\mu\text{l}$  of  $\text{MnCl}_2$  (500 mM), and 2.5  $\mu\text{l}$  of GlcNAc $\beta\text{O}-(\text{CH}_2)_6\text{CO}_2\text{Me}$  (37 mg/ml in dimethyl sulfoxide) for 60 minutes at 37°C. The reaction is quenched by the addition of 1 ml of water and loaded on a C18 Sep-Pak cartridge (Waters), and the column is washed twice with 5 ml of water to remove unreacted UDP-  
20  $^3\text{H}$ galactose. The  $^3\text{H}$ galactosylated GlcNAc $\beta\text{O}-(\text{CH}_2)_6\text{CO}_2\text{Me}$  remains bound to the column during the water washes and is eluted with 5 ml of methanol. Radioactivity in the eluted material is measured by liquid scintillation counting and is proportional to galactosyltransferase activity in the starting sample.

PMMM phosphatase activity is measured by the hydrolysis of p-nitrophenyl phosphate (PNPP). PMMM is incubated together with PNPP in HEPES buffer, pH 7.5, in the presence of 0.1%  
25  $\beta$ -mercaptoethanol at 37°C for 60 min. The reaction is stopped by the addition of 6 ml of 10 N NaOH and the increase in light absorbance at 410 nm resulting from the hydrolysis of PNPP is measured using a spectrophotometer. The increase in light absorbance is proportional to the activity of PMMM in the assay (Diamond, R.H. et al. (1994) Mol. Cell. Biol. 14:3752-3762).

30 In the alternative, PMMM phosphatase activity is determined by measuring the amount of phosphate removed from a phosphorylated protein substrate. Reactions are performed with 2 or 4 nM enzyme in a final volume of 30  $\mu\text{l}$  containing 60 mM Tris, pH 7.6, 1 mM EDTA, 1 mM EGTA, 0.1% 2-mercaptoethanol and 10  $\mu\text{M}$  substrate,  $^{32}\text{P}$ -labeled on serine/threonine or tyrosine, as appropriate. Reactions are initiated with substrate and incubated at 30°C for 10-15 min. Reactions are quenched  
35 with 450  $\mu\text{l}$  of 4% (w/v) activated charcoal in 0.6 M HCl, 90 mM  $\text{Na}_4\text{P}_2\text{O}_7$ , and 2 mM  $\text{NaH}_2\text{PO}_4$ , then

centrifuged at  $12,000 \times g$  for 5 min. Acid-soluble  $^{32}\text{Pi}$  is quantified by liquid scintillation counting (Sinclair, C. et al. (1999) *J. Biol. Chem.* 274:23666-23672).

PMMP protease activity is measured by the hydrolysis of appropriate synthetic peptide substrates conjugated with various chromogenic molecules in which the degree of hydrolysis is quantified by spectrophotometric (or fluorometric) absorption of the released chromophore (Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York, NY, pp. 25-55). Peptide substrates are designed according to the category of protease activity as endopeptidase (serine, cysteine, aspartic proteases, or metalloproteases), aminopeptidase (leucine aminopeptidase), or carboxypeptidase (carboxypeptidases A and B, procollagen C-proteinase).

Commonly used chromogens are 2-naphthylamine, 4-nitroaniline, and furylacrylic acid. Assays are performed at ambient temperature and contain an aliquot of the enzyme and the appropriate substrate in a suitable buffer. Reactions are carried out in an optical cuvette, and the increase/decrease in absorbance of the chromogen released during hydrolysis of the peptide substrate is measured. The change in absorbance is proportional to the enzyme activity in the assay.

In the alternative, an assay for PMMP protease activity takes advantage of fluorescence resonance energy transfer (FRET) that occurs when one donor and one acceptor fluorophore with an appropriate spectral overlap are in close proximity. A flexible peptide linker containing a cleavage site specific for PMMP is fused between a red-shifted variant (RSGFP4) and a blue variant (BFP5) of Green Fluorescent Protein. This fusion protein has spectral properties that suggest energy transfer is occurring from BFP5 to RSGFP4. When the fusion protein is incubated with PMMP, the substrate is cleaved, and the two fluorescent proteins dissociate. This is accompanied by a marked decrease in energy transfer which is quantified by comparing the emission spectra before and after the addition of PMMP (Mitra, R.D. et al (1996) *Gene* 173:13-17). This assay can also be performed in living cells. In this case the fluorescent substrate protein is expressed constitutively in cells and PMMP is introduced on an inducible vector so that FRET can be monitored in the presence and absence of PMMP (Sagot, I. et al (1999) *FEBS Letters* 447:53-57).

An assay for ubiquitin hydrolase activity measures the hydrolysis of a ubiquitin precursor. The assay is performed at ambient temperature and contains an aliquot of PMMP and the appropriate substrate in a suitable buffer. Chemically synthesized human ubiquitin-valine may be used as substrate. Cleavage of the C-terminal valine residue from the substrate is monitored by capillary electrophoresis (Franklin, K. et al. (1997) *Anal. Biochem.* 247:305-309).

PMMP protease inhibitor activity for alpha 2-HS-glycoprotein (AHSG) can be measured as a decrease in osteogenic activity in dexamethasone-treated rat bone marrow cell cultures (dex-RBMC). Assays are carried out in 96-well culture plates containing minimal essential medium supplemented with 15% fetal bovine serum, ascorbic acid (50 mg/ml), antibiotics (100 mg/ml penicillin G, 50

mg/ml gentamicin, 0.3 mg/ml fungizone), 10 mM B-glycerophosphate, dexamethasone ( $10^{-8}$  M) and various concentrations of PMMM for 12-14 days. Mineralized tissue formation in the cultures is quantified by measuring the absorbance at 525 nm using a 96-well plate reader (Binkert, C. et al. (1999) J. Biol. Chem. 274:28514-28520).

5 PMMM protease inhibitor activity for inter-alpha-trypsin inhibitor (ITI) can be measured by a continuous spectrophotometric rate determination of trypsin activity. The assay is performed at ambient temperature in a quartz cuvette in pH 7.6 assay buffer containing 63 mM sodium phosphate, 0.23 mM N a-benzoyl-L-arginine ethyl ester, 0.06 mM hydrochloric acid, 100 units trypsin, and various concentrations of PMMM. Immediately after mixing by inversion, the increase in  $A_{253\text{ nm}}$  is  
10 recorded for approximately 5 minutes and the enzyme activity is calculated (Bergmeyer, H.U. et al. (1974) Meth. Enzym. Anal. 1:515-516).

PMMM isomerase activity such as peptidyl prolyl *cis/trans* isomerase activity can be assayed by an enzyme assay described by Rahfeld, J.U., et al. (1994; FEBS Lett. 352:180-184). The assay is performed at 10°C in 35 mM HEPES buffer, pH 7.8, containing chymotrypsin (0.5 mg/ml) and  
15 PMMM at a variety of concentrations. Under these assay conditions, the substrate, Suc-Ala-Xaa-Pro-Phe-4-NA, is in equilibrium with respect to the prolyl bond, with 80-95% in *trans* and 5-20% in *cis* conformation. An aliquot (2 ml) of the substrate dissolved in dimethyl sulfoxide (10 mg/ml) is added to the reaction mixture described above. Only the *cis* isomer of the substrate is a substrate for cleavage by chymotrypsin. Thus, as the substrate is isomerized by PMMM, the product is cleaved by  
20 chymotrypsin to produce 4-nitroanilide, which is detected by its absorbance at 390 nm. 4-nitroanilide appears in a time-dependent and a PMMM concentration-dependent manner.

PMMM galactosyltransferase activity can be determined by measuring the transfer of radiolabeled galactose from UDP-galactose to a GlcNAc-terminated oligosaccharide chain (Kolbinger, F. et al. (1998) J. Biol. Chem. 273:58-65). The sample is incubated with 14  $\mu$ l of assay  
25 stock solution (180 mM sodium cacodylate, pH 6.5, 1 mg/ml bovine serum albumin, 0.26 mM UDP-galactose, 2  $\mu$ l of UDP-[ $^3$ H]galactose), 1  $\mu$ l of  $\text{MnCl}_2$  (500 mM), and 2.5  $\mu$ l of  $\text{GlcNAc}\beta\text{O}-(\text{CH}_2)_8\text{-CO}_2\text{Me}$  (37 mg/ml in dimethyl sulfoxide) for 60 minutes at 37°C. The reaction is quenched by the addition of 1 ml of water and loaded on a C18 Sep-Pak cartridge (Waters), and the column is washed twice with 5 ml of water to remove unreacted UDP-[ $^3$ H]galactose. The [ $^3$ H]galactosylated  
30  $\text{GlcNAc}\beta\text{O}-(\text{CH}_2)_8\text{-CO}_2\text{Me}$  remains bound to the column during the water washes and is eluted with 5 ml of methanol. Radioactivity in the eluted material is measured by liquid scintillation counting and is proportional to galactosyltransferase activity in the starting sample.

PMMM induction by heat or toxins may be demonstrated using primary cultures of human fibroblasts or human cell lines such as CCL-13, HEK293, or HEP G2 (ATCC). To heat induce  
35 PMMM expression, aliquots of cells are incubated at 42°C for 15, 30, or 60 minutes. Control aliquots

are incubated at 37°C for the same time periods. To induce PMMM expression by toxins, aliquots of cells are treated with 100  $\mu$ M arsenite or 20 mM azetidine-2-carboxylic acid for 0, 3, 6, or 12 hours.

After exposure to heat, arsenite, or the amino acid analogue, samples of the treated cells are harvested and cell lysates prepared for analysis by western blot. Cells are lysed in lysis buffer containing 1%

- 5 Nonidet P-40, 0.15 M NaCl, 50 mM Tris-HCl, 5 mM EDTA, 2 mM N-ethylmaleimide, 2 mM phenylmethylsulfonyl fluoride, 1 mg/ml leupeptin, and 1 mg/ml pepstatin. Twenty micrograms of the cell lysate is separated on an 8% SDS-PAGE gel and transferred to a membrane. After blocking with 5% nonfat dry milk/phosphate-buffered saline for 1 h, the membrane is incubated overnight at 4°C or at room temperature for 2-4 hours with an appropriate dilution of anti-PMMM serum in 2% nonfat
- 10 dry milk/phosphate-buffered saline. The membrane is then washed and incubated with a 1:1000 dilution of horseradish peroxidase-conjugated goat anti-rabbit IgG in 2% dry milk/phosphate-buffered saline. After washing with 0.1% Tween 20 in phosphate-buffered saline, the PMMM protein is detected and compared to controls using chemiluminescence.

- PMMM lysyl hydroxylase activity is determined by measuring the production of
- 15 hydroxy[ $^{14}$ C]lysine from [ $^{14}$ C]lysine. Radiolabeled procollagen is incubated with PMMM in buffer containing ascorbic acid, iron sulfate, dithiothreitol, bovine serum albumin, and catalase. Following a 30 minute incubation, the reaction is stopped by the addition of acetone, and centrifuged. The sedimented material is dried, and the hydroxy[ $^{14}$ C]lysine is converted to [ $^{14}$ C]formaldehyde by oxidation with periodate, and then extracted into toluene. The amount of  $^{14}$ C extracted into toluene is
- 20 quantified by scintillation counting, and is proportional to the activity of PMMM in the sample (Kivirikko, K., and R. Myllyla (1982) *Methods Enzymol.* 82:245-304).

### **XIX. Identification of PMMM Substrates**

- Phage display libraries can be used to identify optimal substrate sequences for PMMM. A random hexamer followed by a linker and a known antibody epitope is cloned as an N-terminal
- 25 extension of gene III in a filamentous phage library. Gene III codes for a coat protein, and the epitope will be displayed on the surface of each phage particle. The library is incubated with PMMM under proteolytic conditions so that the epitope will be removed if the hexamer codes for a PMMM cleavage site. An antibody that recognizes the epitope is added along with immobilized protein A. Uncleaved phage, which still bear the epitope, are removed by centrifugation. Phage in the
- 30 supernatant are then amplified and undergo several more rounds of screening. Individual phage clones are then isolated and sequenced. Reaction kinetics for these peptide substrates can be studied using an assay in Example XVIII, and an optimal cleavage sequence can be derived (Ke, S.H. et al. (1997) *J. Biol. Chem.* 272:16603-16609).

- To screen for *in vivo* PMMM substrates, this method can be expanded to screen a cDNA
- 35 expression library displayed on the surface of phage particles (T7SELECT10-3 Phage display vector,



Novagen, Madison, WI) or yeast cells (pYD1 yeast display vector kit, Invitrogen, Carlsbad, CA). In this case, entire cDNAs are fused between Gene III and the appropriate epitope.

#### **XX. Identification of PMMM Inhibitors**

Compounds to be tested are arrayed in the wells of a multi-well plate in varying concentrations along with an appropriate buffer and substrate, as described in the assays in Example XVIII. PMMM activity is measured for each well and the ability of each compound to inhibit PMMM activity can be determined, as well as the dose-response kinetics. This assay could also be used to identify molecules which enhance PMMM activity.

In the alternative, phage display libraries can be used to screen for peptide PMMM inhibitors. Candidates are found among peptides which bind tightly to a protease. In this case, multi-well plate wells are coated with PMMM and incubated with a random peptide phage display library or a cyclic peptide library (Koivunen, E. et al. (1999) Nature Biotech 17:768-774). Unbound phage are washed away and selected phage amplified and rescreened for several more rounds. Candidates are tested for PMMM inhibitory activity using an assay described in Example XVIII.

Various modifications and variations of the described compositions, methods, and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. It will be appreciated that the invention provides novel and useful proteins, and their encoding polynucleotides, which can be used in the drug discovery process, as well as methods for using these compositions for the detection, diagnosis, and treatment of diseases and conditions. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Nor should the description of such embodiments be considered exhaustive or limit the invention to the precise forms disclosed. Furthermore, elements from one embodiment can be readily recombined with elements from one or more other embodiments. Such combinations can form a number of embodiments within the scope of the invention. It is intended that the scope of the invention be defined by the following claims and their equivalents.

Table 1

IncYTE Project ID	Polypeptide SEQ ID NO:	IncYTE Polypeptide ID	Polynucleotide SEQ ID NO:	IncYTE Polynucleotide ID	IncYTE Full Length Clones
8268274	1	8268274CD1	32	8268274CB1	
7500515	2	7500515CD1	33	7500515CB1	90020436CA2
2256826	3	2256826CD1	34	2256826CB1	
7686186	4	7686186CD1	35	7686186CB1	
72617436	5	72617436CD1	36	72617436CB1	
7501945	6	7501945CD1	37	7501945CB1	
7500264	7	7500264CD1	38	7500264CB1	
7499935	8	7499935CD1	39	7499935CB1	
7982285	9	7982285CD1	40	7982285CB1	4872051CA2
7758505	10	7758505CD1	41	7758505CB1	
6885756	11	6885756CD1	42	6885756CB1	
7500748	12	7500748CD1	43	7500748CB1	
7500749	13	7500749CD1	44	7500749CB1	
7503401	14	7503401CD1	45	7503401CB1	2774614CA2
7503485	15	7503485CD1	46	7503485CB1	5500371CA2
7504076	16	7504076CD1	47	7504076CB1	90173111CA2, 90173203CA2, 90173227CA2
7500926	17	7500926CD1	48	7500926CB1	90205586CA2
7503216	18	7503216CD1	49	7503216CB1	6440464CA2
7503233	19	7503233CD1	50	7503233CB1	
7726576	20	7726576CD1	51	7726576CB1	
7503507	21	7503507CD1	52	7503507CB1	
7503506	22	7503506CD1	53	7503506CB1	90069502CA2
7503509	23	7503509CD1	54	7503509CB1	90208262CA2
7505800	24	7505800CD1	55	7505800CB1	3475431CA2
7503141	25	7503141CD1	56	7503141CB1	
7500362	26	7500362CD1	57	7500362CB1	
7503328	27	7503328CD1	58	7503328CB1	

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Incyte Full Length Clones
7510464	28	7510464CD1	59	7510464CB1	
7510394	29	7510394CD1	60	7510394CB1	
7500745	30	7500745CD1	61	7500745CB1	
7500929	31	7500929CD1	62	7500929CB1	

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
1	8268274CD1	g10441427	4.10E-126	[Drosophila melanogaster] Partner of Paired Raj, L. et al. (2000) Targeted localized degradation of Paired protein in <i>Drosophila</i> development. Curr. Biol. 10:1265-1272
2	7500515CD1	g4096840	0.0	[Homo sapiens] inter-alpha-trypsin inhibitor family heavy chain-related protein Saguchi K. et al. (1996) Isolation and characterization of the human inter-alpha-trypsin inhibitor family heavy chain-related protein (IHRP) gene (ITIH1). J. Biochem. 119:898-905
3	2256826CD1	g5919219	1.20E-188	[Homo sapiens] leucine-rich repeats containing F-box protein FBL3 Ilyin, G. P. et al. (2000) cDNA cloning and expression analysis of new members of the mammalian F-box protein family. Genomics 67:40-47
4	7686186CD1	g11994498	5.70E-72	[Arabidopsis thaliana] DegP protease precursor Kaneko, T. et al. (2000) Structural analysis of Arabidopsis thaliana chromosome 3. II. Sequence features of the 4,251,695 bp regions covered by 90 P1, TAC and BAC clones. DNA Res. 7:217-221
5	72617436CD1	g2190297	1.10E-45	[Oryzias latipes] choriolysin H Yasumasu, S. et al. (1996) Eur. J. Biochem. 237:752-758 Different exon-intron organizations of the genes for two astacin-like proteases, high choriolytic enzyme (choriolysin H) and low choriolytic enzyme (choriolysin L), the constituents of the fish hatching enzyme.
6	7501945CD1	g213504	5.40E-37	[Oryzias latipes] protease Yasumasu, S. et al. (1992) Dev. Biol. 153:250-258 Isolation of cDNAs for LCE and HCE, two constituent proteases of the hatching enzyme of <i>Oryzias latipes</i> , and concurrent expression of their mRNAs during development

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
7	7500264CD1	g2924601	7.60E-12	[Homo sapiens] hepatocyte growth factor activator inhibitor Shimomura, T. et al. (1997) J. Biol. Chem. 272:6370-6376 Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor.
8	7499935CD1	g189842	7.00E-247	[Homo sapiens] prolidase Endo, F. et al. (1989) J. Biol. Chem. 264:4476-4481 Primary structure and gene localization of human prolidase.
9	7982285CD1	g6567172	9.70E-102	[Mus musculus] mDj10 Ohtsuka K, and Hata M. (2000) Cell Stress Chaperones 5:98-112 Mammalian HSP40/DNAJ homologs: cloning of novel cDNAs and a proposal for their classification and nomenclature.
10	7758505CD1	g1167860	1.10E-51	[Spodoptera frugiperda] Endoprotease FURIN
11	6885756CD1	g14994718	3.20E-103	[Mus musculus] (AF393638) deubiquitinating enzyme 2A Baek, K.-H. et al. (2001) Blood 98: 636-42
12	7500748CD1	g14456615	5.40E-280	[Homo sapiens] phosphatidyl inositol glycan class T Ohishi, K. et al. (2001) PIG-S and PIG-T, essential for GPI-anchor attachment to proteins, form a complex with GAA1 and GPI8. EMBO J. 20:4088-4098
13	7500749CD1	g14456615	4.70E-261	[Homo sapiens] phosphatidyl inositol glycan class T Ohishi, K. et al. (supra)
14	7503401CD1	g292031	1.30E-117	[Homo sapiens] farnesyl-protein transferase alpha-subunit Omer, C.A., et al. (1993) Biochemistry 32:5167-76 Characterization of recombinant human farnesyl-protein transferase: Cloning, expression, farnesyl diphosphate binding and functional homology with yeast prenyl-protein transferases.
	7503401CD1	335360[FNTA]	1.10E-118	[Homo sapiens][Transferase][Cyttoplasmic] Alpha subunit of CAAX farnesyltransferase (FPTase) and geranylgeranyltransferase type-I (GGTase-I), transfers farnesyl and geranylgeranyl groups to proteins

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
15	7503485CD1	g11036950	3.30E-60	[Homo sapiens] ubiquitin-conjugating enzyme HRGB
		338766 UBE2A	2.90E-61	[Homo sapiens][Ligase; Protein conjugation factor] Human homolog of S. cerevisiae Rad6p, a member of the ubiquitin-conjugating enzyme family that catalyzes the ubiquitination of cellular proteins and marks them for degradation, also plays a role in DNA repair
16	7504076CD1	g7677403	6.60E-105	[Homo sapiens] F-box protein FBG2 Ilyin, G.P., et al. (2000) Genomics 67: 40-47 cDNA cloning and expression analysis of new members of the mammalian F-box protein family
		598228 FBXO6	5.80E-106	[Homo sapiens][Ligase; Protein conjugation factor] Member of a family of F-box containing proteins, a putative subunit of the SCF ubiquitin ligase involved in protein degradation
17	7500926CD1	g3868871	2.60E-16	[Clostridium histolyticum] Orf2u Matsushita, O., et al. (1999) Gene duplication and multiplicity of collagenases in Clostridium histolyticum. J. Bacteriol. 181:923-933
		377422 pi053	6.90E-16	[Schizosaccharomyces pombe] Conserved protein containing a DUF28 domain
18	7503216CD1	g8489879	0.0	[Homo sapiens] (AF272981) cytosolic aminopeptidase P Cottrell, G.S., et al. (2000) Cloning, expression, and characterization of human cytosolic aminopeptidase P: a single manganese(II)-dependent enzyme. Biochemistry 39:15121-15128
		739810 XPNPEP1	0.0	[Homo sapiens] X-prolyl aminopeptidase (aminopeptidase P)1, soluble
		567960 XPNPEP1L	0.0	[Homo sapiens][Hydrolase; Protease (other than proteasomal)] X-prolyl aminopeptidase-like (aminopeptidase P-like), a putative X-prolyl aminopeptidase, ubiquitously expressed

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
18		344930 XPNPEP2	9.4E-123	[Homo sapiens][Hydrolase; Protease (other than proteasomal)] X-prolyl aminopeptidase (aminopeptidase P) 2 (membrane-bound), metalloproteinase which catalyzes removal of N-terminal amino acids from peptides with N-terminal Xaa-Pro sequences; inhibited by apstatin; may be associated with premature ovarian failure
19	7503233CD1	g29664	0.0	[Homo sapiens] CANP, large subunit (aa 1-714) Aoki, K., et al. (1986) Complete amino acid sequence of the large subunit of the low-Ca2+-requiring form of human Ca2+-activated neutral protease (muCANP) deduced from its cDNA sequence. FEBS Lett. 205:313-317
		661158 CAPN1	0.0	[Homo sapiens][Hydrolase; Protease (other than proteasomal)][Plasma membrane] Calpain I, catalytic subunit of mu-calpain, a calcium-dependent cysteine (thiol) protease that requires micromolar concentrations of calcium in vitro
		334452 CAPN2	1.6E-227	[Homo sapiens][Hydrolase; Protease (other than proteasomal)] Calpain 2, large subunit of the cysteine-type protease m-calpain which may regulate the cell cycle, apoptosis, and cellular differentiation, upregulated in muscle from progressive muscular dystrophy and amyotrophic lateral sclerosis patients
20	7726576CD1	g4079809	2.8E-55	[Homo sapiens] HERC2 Ji, Y., et al. (1999) The ancestral gene for transcribed, low-copy repeats in the Prader-Willi/Angelman region encodes a large protein implicated in protein trafficking, which is deficient in mice with neuromuscular and spermiogenic abnormalities. Hum. Mol. Genet. 8:533-542
		691012 FLJ21156	3.4E-110	[Homo sapiens] has moderate similarity to a region of human HERC1, which is a guanine-nucleotide exchange factor that interacts with ARF1 and binds to Hsp70 and clathrin heavy chain (CLTC)

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
20		345082 HERC2	2.4E-56	[Homo sapiens][Guanine-nucleotide exchange factor] Homolog of murine Mm.20929, which is a guanine-nucleotide exchange factor involved in intracellular protein transport; duplicated and truncated copies of the corresponding gene are associated with deletion breakpoints in Prader-Willi and Angelman syndromes
	7726576CD1	341506 HERC1	4.4E-44	[Homo sapiens][Guanine-nucleotide exchange factor][Golgi; Cytoplasmic] HECT (homologous to E6-AP (UBE3A) carboxy terminus) domain and RCC1 (CHC1)-like domain (RLD) 1, functions as a guanine-nucleotide exchange factor for Rab related proteins and ARF1, may be involved in membrane transport processes
21	7503507CD1	g2924601	3.40E-30	[Homo sapiens] hepatocyte growth factor activator inhibitor. Shimomura, T. et al. (1997) Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor. J. Biol. Chem. 272 (10), 6370-6376
22	7503506CD1	341496 SPINT1	5.9E-31	[Homo sapiens][Inhibitor or repressor][Extracellular (excluding cell wall); Unspecified membrane; Plasma membrane] Serine protease inhibitor (Kunitz type 1), a Kunitz type serine protease inhibitor that inhibits hepatocyte growth factor activator and is found in both membrane-associated and secreted forms Shimomura, T. et al. J. Biol. Chem. 272:6370-6 (1997) Kataoka, H. et al. Cancer Res. 60:6148-59 (2000)
		588049 Spint1	7.7E-27	[Mus musculus][Inhibitor or repressor; Small molecule-binding protein][Unspecified membrane; Extracellular (excluding cell wall)] Serine protease inhibitor (Kunitz type 1), a Kunitz type serine protease inhibitor that inhibits hepatocyte growth factor activator; contains a transmembrane domain
23	7503509CD1	g13278723	2.40E-14	[Homo sapiens] serine protease inhibitor, Kunitz type 1



Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
23		341496 SPINT1	2.10E-15	[Homo sapiens][Inhibitor or repressor][Extracellular (excluding cell wall); Plasma membrane] Serine protease inhibitor (Kunitz type 1), a Kunitz type serine protease inhibitor that inhibits hepatocyte growth factor activator and is found in both membrane-associated and secreted forms
		588049 Spint1	4.60E-13	[Mus musculus][Inhibitor or repressor; Small molecule-binding protein] [Extracellular (excluding cell wall)] Serine protease inhibitor (Kunitz type 1), a Kunitz type serine protease inhibitor that inhibits hepatocyte growth factor activator, contains a transmembrane domain
24	7505800CD1	g292031	2.30E-134	[Homo sapiens] farnesyl-protein transferase alpha-subunit
		335360 FNNTA	2.00E-135	[Homo sapiens][Transferase][Cyttoplasmic] Farnesyl transferase alpha subunit, transfers farnesyl and geranyl-geranyl groups to proteins, implicated to be involved in TGF-beta and activin signaling
				Zhang, F. L. et al. (1994) cDNA cloning and expression of rat and human protein geranylgeranyltransferase type-I. J. Biol. Chem. 269:3175-3180
		720247 lqbg_A	2.60E-128	[Protein Data Bank] Fpt Alpha-Subunit
		328782 Fnta	1.10E-127	[Rattus norvegicus][Transferase] Farnesyl transferase alpha subunit, transfers farnesyl groups to proteins
				Chen, W. J. et al. (1991) Cloning and expression of a cDNA encoding the alpha subunit of rat p21ras protein farnesyltransferase. Proc. Natl. Acad. Sci. U S A 88:11368-11372
		725889 l88d_A	1.10E-127	[Protein Data Bank] Farnesyltransferase (Alpha Subunit)
		582901 Fnta	3.80E-127	[Mus musculus][Transferase] Protein with strong similarity to human FNNTA, which is the alpha subunit of CAAX farnesyl transferase (FPTase) and geranyl-geranyl transferase type-I (GGTase-I), and that transfers farnesyl and geranyl-geranyl groups to proteins
25	7503141CD1	g15929143	3.9E-248	[Homo sapiens] Peptidase D

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
25		339574 PEPD	1.1E-241	[Homo sapiens][Hydrolase; Protease (other than proteasomal)] Peptidase D (prolidase), catalyzes hydrolysis of dipeptides having a C-terminal proline, functions in proline recycling during collagen synthesis, deficiency causes iminodipeptiduria, mental retardation, collagenous tissue defects, and skin lesions
		429336 Pep4	2.9E-221	[Mus musculus][Hydrolase; Protease (other than proteasomal)] Peptidase D (prolidase), putative dipeptidase that catalyzes hydrolysis of substrates having a C-terminal proline, may function in collagen metabolism; deficiency of human PEPD causes mental retardation, collagenous tissue defects, and skin lesions
		687787 K12C11.1	3.3E-118	[Caenorhabditis elegans][Hydrolase; Protease (other than proteasomal)] Small protein with a region of strong similarity to peptidases
		9721 YFR006W	2.6E-54	[Saccharomyces cerevisiae][Unknown] Protein with weak similarity to human X-pro dipeptidase
		646084 orf6.8163	2.9E-53	[Candida albicans][Unknown] Member of the metallopeptidase family M24, has high similarity to S. cerevisiae Yfr006p, which is a protein with weak similarity to human X-pro dipeptidase
26	7500362CD1	g20271451	0.0	[Homo sapiens] peptidase D
		339574 PEPD	5.0E-250	[Homo sapiens] [Hydrolase; Protease (other than proteasomal)] Peptidase D (prolidase), catalyzes hydrolysis of dipeptides having a C-terminal proline, functions in proline recycling during collagen synthesis, deficiency causes iminodipeptiduria, mental retardation, collagenous tissue defects, and skin lesions Endo, F. et al. J Biol Chem 264, 4476-81 (1989). Tanoue, A. et al. J Biol Chem 265, 11306-11 (1990).

Table 2

Polyptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
26		429336 Pep4	8.9E-231	[Mus musculus] [Hydrolase; Protease (other than proteasomal)] Peptidase D (prolidase), putative dipeptidase that catalyzes hydrolysis of substrates having a C-terminal proline, may function in collagen metabolism; deficiency of human PEPD causes mental retardation, collagenous tissue defects, and skin lesions Ishii, T. et al. Biochim Biophys Acta 1308, 15-6 (1996).
27	7503328CD1	g8489879	7.2E-210	[Homo sapiens] cytosolic aminopeptidase P Cottrell, G. S. et al. Biochemistry 39, 15121-15128 (2000)
		567960 XPNPEP L	3.2E-210	[Homo sapiens] [Hydrolase; Protease (other than proteasomal)] X-prolyl aminopeptidase-like (aminopeptidase P-like), a putative X-prolyl aminopeptidase, ubiquitously expressed Vanhoof, G. et al. Isolation and sequence analysis of a human cDNA clone (XPNPEPL) homologous to X-prolyl aminopeptidase (aminopeptidase P). Cytogenet Cell Genet 78, 275-80 (1997).
		332548 Rn.25763	7.5E-202	[Rattus norvegicus] [Hydrolase; Protease (other than proteasomal)] [Cytoplasmic] X-prolyl aminopeptidase (aminopeptidase P)-1 (soluble), catalyzes removal of the N-terminal amino acid from peptides with N terminal Xaa-Pro sequences, has activity against bradykinin, substance P and other bioactive peptides Czirjak, G. et al. Cloning and functional expression of the cytoplasmic form of rat aminopeptidase P. Biochim Biophys Acta 1444, 326-36 (1999).
28	7510464CD1	g13477305	0.0	[Homo sapiens] X-prolyl aminopeptidase (aminopeptidase P) 1, soluble
29	7510394CD1	g14456615	3.60E-64	[Homo sapiens] phosphatidyl inositol glycan class T Ohishi, K. et al. (supra)

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO: or PROTEOME ID NO:	Probability Score	Annotation
30	7500745CD1	g14456615	1.30E-55	[Homo sapiens] phosphatidyl inositol glycan class T Ohishi, K. et al. (supra)

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	8268274CD1	404	S77 S118 S178 S196 S219 S254 S278 S282 S287 S401 S402 T208 T344 T359 T370 T395	N96 N102 N123 N155 N234 N265	F-box domain: T3-A50	HMMER_PFAM
					PROTEIN GRR1 REPEAT SIMILAR C02F5.7 CAENORHA GLUCOSE METABOLISM LEUCINE- REPEAT T13E15.9 PD003743: K128-I266, N102- C238	BLAST_PRODROM
2	7500515CD1	900	S45 S185 S286 S305 S366 S407 S510 S525 S535 S562 S564 S636 S672 S731 S751 S780 S808 T26 T54 T63 T121 T231 T384 T389 T594 T756 T834 T866 Y146	N81 N207 N517 N577	signal_cleavage: M1-A28	SPSCAN
					Signal Peptides: M1-H23, M1-Q24, M1-A28, M1-T27 von Willebrand factor type A domain: N274-V457	HMMER
					Inosine-uridine preferring nucleoside hydrolase family signature BL01247: N353-S397, A531-N542, I298-L312	HMMER_PFAM
						BLIMPS_BLOCKS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
2					INHIBITOR HEAVY CHAIN PD01101: Q65-K98, N256-D308, G348-N367, R439-V493, W548-L557	BLIMPS_PRODROM
					HEAVY CHAIN H4 PRECURSOR INTER-ALPHA-TRYPSIN INHIBITOR ITI FAMILY CHAIN RELATED PROTEIN PD017446: P669-L900	BLAST_PRODROM
					HEAVY CHAIN PRECURSOR INTER-ALPHA-TRYPSIN INHIBITOR ITI SERINE PROTEASE REPEAT SIGNAL PD004379: Q24-K273 PD004369: A430-S620	BLAST_PRODROM
					INTER-ALPHA-TRYPSIN INHIBITOR HEAVY CHAIN H4 PRECURSOR ITI FAMILY CHAIN-RELATED PLASMA KALLIKREIN SENSITIVE GLYCOPROTEIN 120 SERINE PROTEASE PD120343: A621-P668	BLAST_PRODROM
					INTER-ALPHA-TRYPSIN INHIBITOR COMPLEX COMPONENT II DM03009  JX0368 372-855: L372-E738, K607-G826  S30350 378-841: L372-E610, P702-Q850  P19823 408-896: L372-S613, P645-E851	BLAST_DOMO
					HUMAN INTER-ALPHA-TRYPSIN INHIBITOR HEAVY CHAIN-RELATED PROTEIN PRECURSOR DM03690 JX0368 96-278: K96-I279	BLAST_DOMO
					ATP/GTP-binding site motif A (P-loop): A107-S114	MOTIFS

Table 3

SEQ ID NO:	Incye Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	2256826CD1	436	S11 S17 S38 S86 S97 S160 S177 S333 T206	N84 N158 N175 N259	signal_cleavage: M1-A54	SPSCAN
					F-box domain: V23-L70, R99-R146	HMMER_PFAM
					PROTEIN GRR1 REPEAT SIMILAR CAENORHA GLUCOSE METABOLISM LEUCINE REPEAT PD003743: S155-M304, L208-E336, V104-L252, N259-L375, L286-E385	BLAST_PRODOR
					HYPOTHETICAL 76.5 KD PROTEIN EEED8.10 IN CHROMOSOME II PD135828: S17-R344	BLAST_PRODOR
					DNA REPAIR PROTEIN PUTATIVE EXCISION RAD7 PD135808: V104-L386	BLAST_PRODOR
					P45; CYCLIN; CDK2 DM08625 P34284 62-155; K30-L124	BLAST_DOMO
					Cytochrome c family heme-binding site signature: C243-K248	MOTIFS
4	7686186CD1	356	S48 S201 T6 T58 T69 T131 T172 T278 Y140		signal_cleavage: M1-A45	SPSCAN
					Trypsin: H116-V272	HMMER_PFAM
					Cytosolic domain: M1-R18 Transmembrane domain: F19-P41 Non-cytosolic domain: A42-C356	TMHMMER
					HtrA/DegQ protease family signature PR00834: S238-A255, G332-G344, G115-A127, D136- L156, V178-A202, I216-G233	BLIMPS_PRINTS
					V8 serine protease family signature PR00839: V178- L191, I220-L236, D237-I249	BLIMPS_PRINTS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4					Alpha-lytic endopeptidase serine protease (S2A) signature PR00861: H116-A130, L195-S212, I216-A239	BLIMPS_PRINTS
					PROTEASE SERINE PROTEIN PERIPLASMIC SIGNAL PRECURSOR HTRA HYDROLASE PD001397: S173-T302	BLAST_PRODOM
					PROTEASE DEGS CHAIN DM01722[P45129]3-373: R103-A330 DM01722[P09376]1-383: G106-V328 DM01722[P54925]11-395: P102-A346 DM01722[P39099]1-361: G104-V334	BLAST_DOMO
5	72617436CD1	432	S5 S10 S38 S92 S110 S111 S245 S269 S356 S360 S370 S393 T138 T196 T229 T299	N267	signal_cleavage: M1-G33	SPSCAN
					Signal Peptide: M1-A34, M1-G24, M1-S30	HMMER
					CUB domain proteins profile BL01180: G190-D201	BLIMPS_BLOCKS
					Astacin (Peptidase family M12A): N93-P284	HMMER_PFAM
					Astacin family signature PR00480: P230-S245, L268-G281, I121-Y139, Q175-H193, E194-I211	BLIMPS_PRINTS
					PROTEIN GLYCOPROTEIN EGFLIKE DOMAIN HYDROLASE METALLOPROTEASE ZINC PRECURSOR SIGNAL ZYMOGEN PD000834: S92-C282	BLAST_PRODOM



Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
5					ASTACIN DM00570 P31580 5-269: E46-C282 P31579 6-271: L74-C282 P42662 1-183: G101-Y280 P55112 37-308: L87-C282 Neutral zinc metalloproteinases, zinc-binding region signature: I180-L189	BLAST_DOMO
6	7501945CD1	248	S5 S10 S38 S92 S110 S111 T138 T196 T229		signal_cleavage: M1-G33	SPSCAN
					Signal Peptide: M1-A34, M1-G24, M1-S30 CUB domain proteins profile BL01180: G190-D201	HMMER BLIMPS_BLOCKS
					Astacin (Peptidase family M12A): N93-Q246 Neutral zinc metalloproteinases, zinc-binding region signature: S161-N207	HMMER_PFAM PROFILES SCAN
					Astacin family signature PR00480: E194-I211, P230-P245, I121-Y139, Q175-H193	BLIMPS_PRINTS
					PROTEIN GLYCOPROTEIN EGFLIKE DOMAIN HYDROLASE METALLOPROTEASE ZINC PRECURSOR SIGNAL ZYMOGEN PD000834: S92-R241	BLAST_PROD OM
					ASTACIN DM00570 P31580 5-269: E46-R241 P31579 6-271: L74-R241 P42662 1-183: G101-G240 P55112 37-308: L87-G240 Neutral zinc metalloproteinases, zinc-binding region signature: I180-L189	BLAST_DOMO
						MOTIFS

Table 3

SEQ ID NO:	Incye Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7	7500264CD1	388	S11 S34 S53 S201 S234 S261 S265 S270 S283 S310 S323 T166 T197 T285 T375 Y319	N164 N289	signal_cleavage: M1-A37	SPSCAN
					Signal Peptide: M1-A37	HMIMER
					Cytosolic domain: R362-L388 Transmembrane domain: A339-L361 Non-cytosolic domain: M1-G338	TMHMMER
					HEPATOCYTE GROWTH FACTOR ACTIVATOR INHIBITOR GLYCOPROTEIN PD120361: G84-V254, S11-P44	BLAST_PRODROM
					Leucine zipper pattern: L45-L66, L347-L368	MOTIFS
8	7499935CD1	467	S138 S167 S224 S312 S434 T15 T54 T90 T146 T188 T198 T445 Y128	N13 N172	Metallopeptidase family M24: V185-E447	HMIMER_PFAM
					Aminopeptidase P and proline dipeptidase proteins BL00491: A319-H330, H366-D378, G422-G435	BLIMPS_BLOCKS
					PROLIDASE HYDROLASE XAAPRO DIPEPTIDASE XPRO PROLINE IMDODIPEPTIDASE ACETYLATION MANGANESE PEPTIDASE PD013444: A2-V185	BLAST_PRODROM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
8					AMINOPEPTIDASE HYDROLASE METHIONINE PEPTIDASE PROTEIN COBAL T M DIPEPTIDASE XPRO MAP PD000555: V185-T440	BLAST_PRODROM
					AMINOPEPTIDASE P AND PROLINE DIPEPTIDASE DM00816 P12955 179-467: I180-I389, I389-P443 P43590 225-509: E182-P383, Y390-T440 P15034 169-423: E182-T440 P44881 163-417: E182-P443	BLAST_DOMO
					Aminopeptidase P and proline dipeptidase signature: H366-D378	MOTIFS
9	7982285CDI	379	S42 S67 S160 S234 S319 T171 T197 T286 Y324	N53 N76	DnaJ domain: N108-G172	HMMER_PFAM
					Nt-dnaJ domain proteins BL00636: D123-K139, F149-D169	BLIMPS_BLOCKS
					DnaJ domains signatures and profile: R129-N187	PROFILES CAN
					DnaJ protein family signature PR00625: A119-D138, F149-D169, S55-K74	BLIMPS_PRINTS
					PROTEIN HLJ1 B0035.14 CHROMOSOME IV CHAPERONE PD036881: P198-L372	BLAST_PRODROM
					PROTEIN CHAPERONE DNAJ HEAT SHOCK DNA REPLICATION REPEAT ANTIGEN T PD000231: N108-D169	BLAST_PRODROM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
9					NT-DNAJ DOMAIN DM00098 P35515 I-101: K107-G208 P30725 I-102: K107-G20 P25685 I-107: K107-G208 S34632 I-99: Y109-G209 Cell attachment sequence: R242-D244 Ni-dnaJ domain signature: F149-Y168 N-6 Adenine-specific DNA methylases signature: M269-Y275	BLAST_DOMO     MOTIFS MOTIFS MOTIFS
10	7758505CD1	737	S48 S217 S301 S477 S559 S572 S733 T390 T498 T524 T577 T579 T602 Y646	N361	signal_cleavage: M1-G26  Signal Peptide: M1-A15; M1-H23; M1-G26; M1-P21  von Willebrand factor type C domain: C159-C216, C285-C342, C28-C87, K366-C416, C95-C152, C221-C278  PRECURSOR SIGNAL RECEPTOR GLYCOPROTEIN TRANSMEMBRANE KINASE TRANSFERASE TYROSINE PROTEIN ATP-BINDING PHOSPHORYLATION PD000495: S395-C662  PROTEIN PRECURSOR REPEAT GLYCOPROTEIN SIGNAL NEL EGF-LIKE DOMAIN CHORDIN B0024.14 PD015143: C84-C152	SPSCAN  HMMER  HMMER_PFAM  BLAST_PRODOM  BLAST_PRODOM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
10					VON WILLEBRAND FACTOR TYPE C REPEAT DM00551 A38963 649-756: W41-V153 Cytochrome c family heme-binding site signature: C510-E515, C605-N610, C652-S657 VWFC domain signature: C46-C87, C113-C152, C177-C216, C239-C278, C303-C342, C379-C416	BLAST_DOMO  MOTIFS  MOTIFS
11	6885756CD1	530	S22 S23 S36 S38 S71 S72 S110 S232 S272 S284 S376 S432 S491 S511 S522 T47 T399 T404 T446 T495 T512 Y306 Y347	N92 N444 N488	Ubiquitin carboxyl-terminal hydrolases family: A80-R111  Ubiquitin carboxyl-terminal hydrolase family: G313-Q374 Ubiquitin carboxyl-terminal hydrolases family 2 proteins BL00972: G81-L98, G156-L165, I193-Y207, Y317-A341, E343-S364 PROTEASE UBIQUITIN HYDROLASE UBIQUITIN SPECIFIC ENZYME DEUBIQUITINATING CARBOXYL TERMINAL THIOLESTERASE PROCESSING CONJUGATION PD017412: F217-Q309 UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2 DM00659  P40818 782-1103: H180-L370, I85-Y107  P50102 141-420: Q158-G327	HMMER_PFAM  BLIMPS_BLOCKS  BLAST_PRODROM  BLAST_DOMO

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
11					Ubiquitin carboxyl-terminal hydrolases family 2 signature 2: Y317-Y335 signal_cleavage: M1-A23	MOTIFS
12	7500748CD1	511	S29 S81 S108 S126 S156 S196 T52 T119 T121 T267 T303 T311 Y287 Y346	N164 N291 N327		SPSCAN
					Signal Peptide: M5-C21; M5-A23; M1-A23; M1-P25;	HMMER
					Cytosolic domain: N478-L511 Transmembrane domain: F455-Y477 Non-cytosolic domain: M1-D454	TMHMMER
					PROTEIN FI7C11.7 IKI1ERG9 INTERGENIC REGION TRANSMEMBRANE PD043343; E32-V245	BLAST_PRODOM
					PROTEIN FI7C11.7 IKI1ERG9 INTERGENIC REGION TRANSMEMBRANE PD043344; H348-V399, L441-G474	BLAST_PRODOM
13	7500749CD1	476	S29 S94 T52 T165 T201 T209 T301 Y67 Y185	N189 N225	signal_cleavage: M1-A23	SPSCAN
					Signal Peptide: M5-C21; M5-A23; M1-C21; M1-A23; M1-P25	HMMER
					Cytosolic domain: N443-L476 Transmembrane domain: F420-Y442 Non-cytosolic domain: M1-D419	TMHMMER
					PROTEIN FI7C11.7 IKI1ERG9 INTERGENIC REGION TRANSMEMBRANE PD043344; H313-V364	BLAST_PRODOM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
14	7503401CD1	344	S49 S93 S113 T315 T334 Y102	N130 N198 N211 N238	Protein prenyltransferase alpha subunit repe: Q149-K179, R115-R145, R223-G253, N183-T213 Protein prenyltransferases alpha subunit repeat proteins BL00904: R115-S148, Q149-D189, R223-G247, I248-E285, Q291-T315, G14-Y63	HMMER_PFAM BLIMPS_BLOCKS
					TRANSFERASE SUBUNIT ALPHA PROTEIN PRENYLTRANSFERASE REPEAT FARNESYLTRANSFERASE PROTEINS CAAX RAS PD005875: V99-Q343 TRANSFERASE SUBUNIT FARNESYLTRANSFERASE PROTEIN ALPHA CAAX RAS PROTEINS PRENYLTRANSFERASE FTASE ALPHA PD011572: V64-V106	BLAST_PRODROM BLAST_PRODROM
					FARNESYLTRANSFERASE; ALPHA; DM07118 P49354 257-378: E222-Q344 PROTEIN PRENYLTRANSFERASES ALPHA SUBUNIT REPEAT DM01356 P49354 174-212: V139-F178	BLAST_DOMO BLAST_DOMO
					Protein prenyltransferases alpha subunit repeat signature: A128-R137, A162-R171, V196-R205, P236-L245	MOTIFS
15	7503485CD1	122	S81 S112 S118 T46		Ubiquitin-conjugating enzyme: M1-Q117	HMMER_PFAM
					signal_cleavage: M34-D84	SPSCAN
					Ubiquitin-conjugating enzymes active site: P21-E85	PROFILESAN

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
15					UBIQUITIN LIGASE ENZYME PROTEIN UBIQUITIN-CONJUGATING CONJUGATION CARRIER UBIQUITIN PROTEIN MULTIGENE FAMILY PD000461: A53-R110, M1-E49 UBIQUITIN-CONJUGATING ENZYMES DM00225 A41222 2-149: D50-R120, S20-D50 P52478 2-149: D50-W119, S20-D50 P25865 2-149: D50-W119, S2-E99 P23566 2-149: D50-W119, S20-D50	BLAST_PRODOM
16	7504076CD1	255	T153 T154	N245	F-box domain: G4-L52	HMIMER_PFAM
17	7500926CD1	166	S35 S120 S157	N155	signal_cleavage: M1-A19 Signal Peptide: M1-A26	SPSCAN HMIMER
					PROTEIN INTERGENIC REGION CONSERVED OF SECTION COAT MG332 VMA7RPS31A VMA7RPS25A PD004323: K62-K127	BLAST_PRODOM
18	7503216CD1	591	S54 S182 S214 S251 S259 S329 S375 S447 S473 S544 T102 T156 T370 T405 T476 T546 T559	N89 N371	metallopeptidase family M24: A288-N520	HMIMER_PFAM
					Aminopeptidase P and proline dipeptidase proteins BL00491: I352-H363, H453-V465, L482-E496, G501-K514	BLIMPS_BLOCKS
					Proteasome A-type and B-type PF00227: I31-Y42	BLIMPS_PFAM



Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
18					AMINOPEPTIDASE AMINOACYL PROLINE HYDROLASE XAA PRO XPRO PROLINE P PROTEIN PRECURSOR P-LIKE PD009635: K514-S588	BLAST_PRODROM
					AMINOPEPTIDASE HYDROLASE AMINOACYL PROLINE XAA PRO P XPRO PROTEIN PUTATIVE PROLINE APP PD004954: L10-D120	BLAST_PRODROM
					AMINOPEPTIDASE HYDROLASE METHIONINE PEPTIDASE PROTEIN COBAL T M DIPEPTIDASE XPRO MAP PD000555: E369-V510, V289-G493	BLAST_PRODROM
					AMINOPEPTIDASE AMINOACYL PROLINE HYDROLASE P XAA PRO XPRO PROLINE PRECURSOR PROTEIN P-LIKE PD008419: E148-A288	BLAST_PRODROM
					AMINOPEPTIDASE P AND PROLINE DIPEPTIDASE DM00816 S64780 449-697: I283-P534 P54518 121-347: K322-V510 P46545 131-358: E319-N506 Q10698 140-366: K287-V510	BLAST_DOMO
19	7503233CD1	652	S77 S147 S194 S248 S275 S317 S422 S437 S453 S477 S500 S527 S574 S602 T202 T284 T300 T307 T312 T632	N76 N137 N305 N406	Calpain large subunit, domain III: K303-V460	HMMER_PFAM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
19					Calpain family cysteine protease: L55-T292	HMME PFAM
					EF hand: N557-A585, S527-I555, T622-T649, N484-I511	HMME PFAM
					EF-hand calcium-binding domain proteins BL00018: D536-F548	BLIMPS_BLOCKS
					Calpain cysteine protease (C2) family signature PR00704: Y83-I108, L113-V136, G138-L165, E268-C289, T318-F335, R426-E454	BLIMPS_PRINTS
					PROTEASE CALPAIN HYDROLASE SUBUNIT NEUTRAL THIOL LARGE CALCIUM- ACTIVATED PROTEINASE CANP PD001545: Y83-T292, L55-G82	BLAST_PRODROM
					PROTEASE CALPAIN HYDROLASE SUBUNIT LARGE NEUTRAL THIOL CALCIUM- ACTIVATED PROTEINASE CANP PD001874: K303-T459	BLAST_PRODROM
					CALPAIN SUBUNIT CALCIUM-BINDING NEUTRAL PROTEASE CALCIUM-ACTIVATED PROTEINASE CANP HYDROLASE LARGE PD003609: E480-F548	BLAST_PRODROM
					CALPAIN SUBUNIT PROTEASE NEUTRAL CALCIUM-BINDING CALCIUM-ACTIVATED PROTEINASE CANP HYDROLASE LARGE PD002827: N549-V614	BLAST_PRODROM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
19					CALPAIN CATALYTIC DOMAIN DM01305 P07384 I1-517: G82-S456, C11-G82 P00789 3-507: Y83-K455, G13-G82 P17655 I-505: N76-K455, G13-G82, V434-Q491 A48764 I-507: Q66-K455, V14-G82, Y432-Q491, K521-R554 EF-hand calcium-binding domain: D536-F548, D566-M578	BLAST_DOMO      MOTIFS
20	7726576CD1	861	S12 S29 S187 S239 S251 S264 S324 S327 S372 S465 S509 S547 S558 S591 S628 S834 S857 T106 T107 T268 T284 T289 T300 T399 T435 T735 T758 T781 T793 Y844	N257 N577 N779	HECT-domain (ubiquitin-transferase): P565-S857	HMMER_PFAM
					Pyrokinins proteins BL00539: F470-L474 HECT-domain (ubiquitin-transferase) PF00632: V680-G686, Y772-P799, L817-N848	BLIMPS_BLOCKS BLIMPS_PFAM
					PROTEIN LIGASE UBIQUITIN CONJUGATION REPEAT UBIQUITIN PROTEIN DNA-BINDING PROBABLE ONCOGENIC PD002225: V587-A845	BLAST_PRODUM
					HERC2 RELEASING UBIQUITIN FACTOR REPEAT CONJUGATION GUANINE- NUCLEOTIDE KIAA0076 HA0936 PD155960: E245-I370	BLAST_PRODUM

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
20					HECT DOMAIN DM01690 P51593 9-306: R583-C849 P40985 578-891: Y598-A845 P39940 513-808: N589-A845 A38919 785-1082: N574-A851 Leucine zipper pattern: L404-L425 signal_cleavage: M1-A37	BLAST_DOMO
21	7503507CD1	447	S11 S34 S53 S201 S234 S339 S377 S382 S395 S406 T166 T197 T263 T279 T397	N164 N291	Signal Peptide: M1-A37 HEPATOCYTE GROWTH FACTOR ACTIVATOR INHIBITOR GLYCOPROTEIN PD120361: G84-L297 signal_cleavage: M1-A37	HMMER BLAST_PRODUM
22	7503506CD1	468	S11 S34 S53 S201 S234 S339 S377 S382 S395 T166 T197 T263 T279 T397 T455	N164 N291 N401	signal_cleavage: M1-A37	SPSCAN
					Signal Peptide: M1-A37	HMMER
					Low-density lipoprotein receptor domain: H308-N346	HMMER_PFAM
					Cytosolic domain: A438-L468 Transmembrane domain: V415-V437 Non-cytosolic domain: M1-P414	TMHMMER
					LDL-receptor class A (LDLRA) domain proteins BL01209: C329-E341	BLIMPS_BLOCKS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
22					HEPATOCYTE GROWTH FACTOR ACTIVATOR INHIBITOR GLYCOPROTEIN PD120361: G84-L297, S11-P44 LDL-receptor class A (LDLRA) domain signature: C322-C344 Leucine zipper pattern: L45-L66, L427-L448 signal_cleavage: M1-A37 Signal Peptide: M1-A37	BLAST_PRODROM MOTIFS MOTIFS SPSCAN HMMER
23	7503509CD1	236	S11 S34 S53 S201 S207 T166 T197	N164	HEPATOCYTE GROWTH FACTOR ACTIVATOR INHIBITOR GLYCOPROTEIN PD120361: G84-R188, S11-P44 Leucine zipper pattern: L45-L66 Protein prenyltransferase alpha subunit repe: Q117-K147, E83-R113, R191-G221, N151-T181 Protein prenyltransferases alpha subunit repeat proteins BL00904: E83-S116, Q117-D157, R191-G215, I216-E253, Q259-T283 TRANSFERASE SUBUNIT ALPHA PROTEIN PRENYL TRANSFERASE REPEAT FARNESYL TRANSFERASE PROTEINS CAAX RAS PD005875: A40-Q236, D59-Q311 FARNESYL TRANSFERASE; ALPHA; DM07118 P49354 257-378: E190-Q312 PROTEIN PRENYL TRANSFERASES ALPHA SUBUNIT REPEAT DM01356 P49354 174-212: V107-F146	BLAST_PRODROM MOTIFS HMMER_PFAM BLIMPS_BLOCKS BLAST_PRODROM BLAST_DOMO BLAST_DOMO
24	7505800CD1	312	S49 T283 T302 Y87	N166 N179 N206		

Table 3

SEQ ID NO:	Incye Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
					Protein prenyltransferases alpha subunit repeat signature: P96-R105, A130-R139, V164-R173, P204-L213	MOTIFS
25	7503141CD1	452	S138 S167 S271 S355 S419 T15 T54 T90 T146 T358 T430 Y128	N13 N172	metallopeptidase family M24: V143-E432	HMME PFAM
					Aminopeptidase P and proline dipeptidase proteins BL00491: A278-H289, H325-D337, L362-F376, G407-G420	BLIMPS_BLOCKS
					PROLIDASE HYDROLASE XAA PRO DPEPTIDASE XPRO PROLINE IMIDODIPEPTIDASE ACETYLATION MANGANESE PEPTIDASE PD013444: A2-C183	BLAST_PRODOME
					AMINOPEPTIDASE HYDROLASE METHIONINE PEPTIDASE PROTEIN COBAL T M DPEPTIDASE XPRO MAP PD000555: E160-Y375; G404-T425	BLAST_PRODOME
					AMINOPEPTIDASE P AND PROLINE DPEPTIDASE DM00816  P12955 179-467: V181-P428  P43590 225-509: V181-T425  P15034 169-423: H187-Y375; E397-T425  P4488 163-417: K168-Y375; E397-P428	BLAST_DOMO
					Aminopeptidase P and proline dipeptidase signature: H325-D337	MOTIFS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
26	7500362CD1	471	S116 S145 S202 S290 S374 S438 T15 T68 T124 T166 T176 T377 T449 Y106	N13 N150	signal_cleavage: M1-T56	SPSCAN
					metallopeptidase family M24: V163-E451	HMMEP_FAM
					Aminopeptidase P and proline dipeptidase proteins BL00491: A297-H308, H344-D356, L381-F395, G426-G439	BLIMPS_BLOCKS
					Pyrokinins proteins BL00539: F74-L78	BLIMPS_BLOCKS
					PROLIDASE HYDROLASE XAAPRO	BLAST_PRODUM
					DIPEPTIDASE XPRO PROLINE IMIDO- DIPEPTIDASE ACETYLATION MANGANESE PEPTIDASE	
					PD013444: A2-F48, F48-V163	
					AMINOPEPTIDASE HYDROLASE METHIONINE PEPTIDASE PROTEIN COBAL T M DIPEPTIDASE XPRO MAP	BLAST_PRODUM
					PD000555: V163-Y394, G423-T444	
					AMINOPEPTIDASE P AND PROLINE DIPEPTIDASE DM00816	BLAST_DOMO
					P12955 179-467: I158-P447 P43590 225-509: E160-T444 P15034 169-423: E160-Y394, E416-T444 P4488 163-417: E160-Y394, E416-P447	
					Aminopeptidase P and proline dipeptidase signature: H344-D356	MOTIFS

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
27	7503328CD1	458	S4 S97 S220 S257 S289 S326 S334 S404 S431 T36 T145 T199 T243	N132	AMINOPEPTIDASE AMINOACYL-PROLINE HYDROLASE P XAA PRO XPRO PROLINE PRECURSOR PROTEIN P-LIKE PD008419: G183-A363	BLAST_PRODROM
					AMINOPEPTIDASE HYDROLASE AMINOACYLPROLINE XAAPRO P XPRO PROTEIN PUTATIVE PROLINE APP PD004954: L53-D163	BLAST_PRODROM
28	7510464CD1	695	S4 S97 S220 S257 S289 S326 S334 S404 S450 S522 S548 S619 S670 T36 T145 T199 T243 T445 T480 T551 T621 T643	N132 N446	metallopeptidase family M24: A363-N595	HMMER_PFAM
					Aminopeptidase P and proline dipeptidase proteins BL00491: I427-H438, H528-V540, L557-E571, G576-K589	BLIMPS_BLOCKS
					Proteasome A-type and B-type PF00227: I74-Y85	BLIMPS_PFAM
					AMINOPEPTIDASE AMINOACYL PROLINE HYDROLASE P XAA PRO XPRO PROLINE PRECURSOR PROTEIN P-LIKE PD008419: G183-A363	BLAST_PRODROM
					AMINOPEPTIDASE HYDROLASE AMINOACYL PROLINE XAA PRO P XPRO PROTEIN PUTATIVE PROLINE APP PD004954: L53-D163	BLAST_PRODROM



Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
28					AMINOPEPTIDASE HYDROLASE METHIONINE PEPTIDASE PROTEIN COBAL T M DIPEPTIDASE XPRO MAP PD000555: V364-G568, E444-V585	BLAST_PRODUM
					AMINOPEPTIDASE P AND PROLINE DIPEPTIDASE DM00816 S64780 449-697: I358-P609 DM00816 P54518 121-347: K397-V585 DM00816 P46545 131-358: E394-N581 DM00816 Q10698 140-366: K362-V585 signal_cleavage: M1-A23	BLAST_DOMO
29	7510394CD1	140	S29 S81 S108 T52 T119 T129		signal_cleavage: M1-A23	SPSCAN
					Signal Peptide: M5-C21, M5-A23, M1-C21, M1-A23, M1-P25	HMMER
					DEVELOPMENT-ASSOCIATED NEURONAL T1B9.20 DJ453C12.7 CGI-06 PD043343: E32-A134 signal_cleavage: M1-A23	BLAST_PRODUM
30	7500745CD1	191	S29 S81 T52		Signal Peptide: M1-C21, M1-A23, M1-P25, M5-C21, M5-A23 signal_cleavage: M1-A19	SPSCAN
					Signal Peptide: M1-C21, M1-A23, M1-P25, M5-C21, M5-A23	HMMER
31	7500929CD1	145	S35		Signal Peptide: M1-A26	SPSCAN
					Signal Peptide: M1-A26	HMMER

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
32/8268274CB1/ 2129	1-530, 53-231, 53-626, 81-845, 127-387, 174-746, 337-654, 369-766, 377-1014, 420-716, 443-887, 478-761, 505-745, 505-921, 505-1101, 523-1000, 545-838, 605-913, 645-864, 665-721, 671-962, 671-967, 714-922, 758-1019, 778-1249, 796-1046, 846-1358, 868-1104, 1146-1296, 1159-1613, 1236-1748, 1284-1319, 1316-1860, 1317-1714, 1361-1926, 1362-1597, 1421-1794, 1450-1747, 1471-1748, 1616-1913, 1616-2097, 1616-2119, 1618-1887, 1666-2129, 1699-1952, 1715-1960
33/7500515CB1/ 3489	1-799, 1-845, 1-851, 1-870, 1-880, 1-894, 1-907, 1-942, 1-2859, 8-171, 8-193, 11-229, 14-736, 108-845, 199-790, 217-577, 233-886, 288-898, 299-804, 302-942, 339-790, 350-859, 360-629, 391-890, 396-988, 461-739, 566-1183, 587-1301, 589-1360, 595-738, 621-1216, 632-981, 736-1009, 736-1288, 741-1363, 790-1019, 814-1581, 853-1339, 854-1114, 863-1536, 864-1206, 868-1340, 872-1479, 874-1195, 897-1596, 898-1165, 904-1173, 912-1626, 948-1626, 991-1220, 1001-1631, 1037-1250, 1047-1803, 1093-1751, 1101-1809, 1103-1738, 1127-1664, 1160-1409, 1189-1873, 1190-1874, 1235-1576, 1241-1506, 1297-1463, 1302-1880, 1370-1491, 1370-1624, 1390-1779, 1401-1574, 1404-1868, 1419-1651, 1437-1875, 1442-1854, 1456-1756, 1474-1880, 1495-1759, 1530-1810, 1542-1880, 1544-1797, 1609-2049, 1660-1933, 1676-1926, 1704-1866, 1710-1880, 1751-2098, 1835-2412, 1852-2366, 1866-2472, 1916-2859, 1940-2861, 1963-2861, 1985-2858, 1991-2859, 1995-2608, 2007-2861, 2008-2859, 2014-2375, 2032-2557, 2083-2690, 2104-2667, 2104-2838, 2106-2617, 2107-2389, 2111-2606, 2138-2687, 2141-2775, 2148-2744, 2158-2692, 2160-2467, 2176-2822, 2188-2857, 2199-2383, 2221-2798, 2260-2483, 2265-2827, 2267-2789, 2276-2536, 2281-2603, 2282-2526, 2294-2876, 2307-2845, 2310-2429, 2325-2848, 2340-2591, 2342-2845, 2349-2582, 2365-2788, 2367-2702, 2374-2632, 2386-2518, 2388-2654, 2388-2847, 2388-2873, 2397-2826, 2401-2849, 2428-2646, 2435-2862, 2435-2876, 2435-2877, 2454-2863, 2456-2673, 2481-2865, 2493-2858, 2497-2753, 2498-2751, 2525-3489, 2533-2698, 2536-2839, 2542-2792, 2543-2785, 2550-2790, 2550-2846, 2550-2857, 2550-2871, 2576-2824, 2608-2858, 2662-2856, 2669-2857, 2735-2855, 2737-2898

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
34/2256826CB1/ 2996	1-686, 424-2842, 443-1082, 452-978, 461-1074, 511-800, 511-867, 511-924, 530-597, 530-881, 567-987, 688-1062, 721-837, 721-956, 721-970, 721-1144, 721-1155, 721-1188, 721-1222, 721-1232, 721-1236, 721-1270, 721-1273, 721-1279, 721-1281, 721-1294, 721-1312, 721-1322, 722-1351, 725-1342, 785-1330, 851-1473, 919-1600, 971-1410, 982-1572, 985-1574, 998-1581, 1001-1478, 1055-1705, 1072-1724, 1105-1645, 1123-1745, 1142-1774, 1187-1828, 1217-1619, 1274-1827, 1280-1914, 1296-1936, 1341-1907, 1345-1836, 1363-1735, 1371-1814, 1379-1950, 1392-1852, 1440-1996, 1456-1935, 1458-1944, 1464-2017, 1469-1984, 1481-1987, 1488-1997, 1491-1967, 1491-1988, 1494-1988, 1504-1980, 1504-2035, 1517-2051, 1538-2209, 1547-2109, 1555-2183, 1580-2238, 1591-2236, 1610-2261, 1619-2099, 1657-2220, 1666-2180, 1671-2114, 1672-2315, 1675-2370, 1677-2222, 1686-2149, 1691-2250, 1697-2189, 1736-2292, 1755-2235, 1775-2375, 1789-2266, 1794-2363, 1828-2395, 1852-2368, 1859-2317, 1890-2529, 1911-2493, 1946-2352, 1953-2498, 1962-2435, 1977-2617, 2072-2610, 2093-2602, 2102-2498, 2113-2565, 2137-2498, 2141-2546, 2158-2706, 2173-2663, 2177-2803, 2191-2842, 2207-2719, 2208-2813, 2316-2799, 2330-2996, 2356-2817, 2375-2805, 2398-2822, 2452-2828 1-936, 4-820, 32-1002, 118-955, 135-957, 141-970, 142-951, 150-1094, 178-800, 208-1045, 223-1245, 235-1130, 244-1046, 254-1067, 271-1007, 314-1137, 318-946, 340-1181, 377-1000, 377-1005, 399-1218, 420-1190, 444-1017, 473-1086, 473-1321, 476-1219, 502-1002, 508-924, 542-1375, 580-1054, 594-1461, 620-1526, 625-1457, 649-1078, 732-1432, 735-1244, 744-1312, 769-945, 777-1648, 781-1314, 817-1629, 818-1618, 818-1626, 825-1721, 830-1667, 832-1814, 837-1381, 839-1216, 839-1389, 881-1668, 924-1756, 930-1696, 939-1844, 958-1844, 965-1830, 970-1814, 973-1457, 980-1814, 980-1826, 985-1814, 988-1737, 991-1814, 995-1834, 998-1814, 999-1770, 999-1846, 1003-1836, 1018-1400, 1022-1830, 1022-1860, 1024-1814, 1029-1842, 1035-1837, 1036-1830, 1037-1814, 1038-1845, 1039-1814, 1040-1833, 1042-1830, 1049-1828, 1055-1811, 1056-1814, 1060-1814, 1061-1814, 1064-1557, 1066-1814, 1069-1700, 1074-1844, 1082-1818, 1103-1841, 1111-1814, 1122-1814, 1129-1814, 1132-1840, 1141-1814, 1147-1814, 1153-1812, 1154-1814, 1160-1814, 1168-1814, 1176-1814, 1176-1823, 1186-1814, 1193-1842, 1196-1814, 1205-1814, 1222-1814, 1226-1814, 1257-1834, 1263-1814, 1266-1814, 1270-1814, 1271-1814, 1276-1814, 1277-1816, 1280-1814, 1288-1830, 1296-1814, 1310-1814, 1311-1600, 1311-1814, 1328-1842, 1336-1814, 1359-1814, 1446-1835, 1461-1844, 1462-1815, 1469-1814, 1536-1738
35/7686186CB1/ 1860	

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
36/72617436CB1/ 1334	1-281, 1-1334, 77-432, 77-477, 78-375, 78-407, 78-456, 78-459, 78-485, 85-375, 94-317, 94-379, 94-380, 94-1334, 114-576, 159-622, 282-600, 282-917, 282-966, 282-1035, 376-758, 581-1334, 859-1334
37/7501945CB1/ 2070	1-655, 502-984, 792-2070, 892-1172, 969-1266, 969-1298, 985-1207, 985-1208, 985-1270, 1173-1832, 1173-1841, 1173-1867, 1173-1890, 1173-1918, 1173-1931, 1173-1966, 1173-1983, 1173-1995, 1173-1996, 1173-2044, 1173-2054, 1177-1979, 1233-2070, 1234-2069, 1236-1708, 1242-2070, 1277-2070, 1336-2070, 1417-1800, 1514-1801, 1524-2070, 1579-1899
38/7500264CB1/ 2265	1-471, 1-2265, 46-537, 113-743, 113-800, 113-850, 113-871, 114-589, 114-856, 117-857, 124-619, 649-1118, 649-1129, 653-785, 653-1234, 657-931, 657-1159, 657-1162, 660-1185, 665-1449, 667-1000, 672-908, 677-1137, 691-826, 691-880, 691-901, 693-891, 759-906, 915-1449, 937-1449, 948-1449, 1012-1290, 1025-1699, 1033-1457, 1034-1290, 1066-1363, 1144-1419, 1164-1396, 1238-1542, 1242-1471, 1242-1491, 1246-1496, 1251-1856, 1282-1539, 1282-1540, 1282-1807, 1282-1874, 1322-1813, 1328-1797, 1334-1847, 1334-1882, 1347-1808, 1381-1640, 1416-1707, 1417-1704, 1420-1641, 1420-2200, 1475-1950, 1486-1963, 1529-1832, 1551-1814, 1551-2113, 1589-1896, 1597-1876, 1603-1829, 1618-1858, 1638-1918, 1666-1914, 1666-1961, 1667-1931, 1673-1955, 1702-1968, 1716-2006, 1718-1905, 1747-2170, 1808-1952, 1808-2126, 1819-2035, 1819-2149, 1853-2064, 1853-2244, 1859-2149, 1859-2265, 1885-2167, 1892-2117, 1909-2176, 1941-2156, 1941-2161, 1954-2192, 1956-2222, 1956-2223, 2004-2251, 2012-2253, 2031-2197, 2031-2265, 2039-2265, 2075-2211
39/7499935CB1/ 1834	1-369, 1-654, 1-715, 1-1174, 2-285, 2-300, 2-429, 2-458, 2-514, 2-524, 2-583, 2-594, 2-608, 2-609, 2-637, 2-642, 2-646, 2-663, 2-691, 2-735, 2-772, 2-838, 3-290, 3-308, 3-362, 3-583, 5-146, 5-308, 5-343, 5-484, 5-536, 5-541, 6-609, 7-267, 7-285, 7-297, 8-199, 8-206, 8-236, 8-273, 8-547, 8-655, 9-633, 11-253, 11-585, 12-227, 12-243, 12-254, 12-283, 12-319, 14-524, 14-1834, 15-270, 15-272, 15-287, 16-247, 16-252, 16-277, 16-296, 16-297, 16-529, 16-652, 16-775, 17-282, 17-290, 17-294, 18-133, 18-260, 18-290, 18-311, 18-623, 25-260, 25-313, 25-320, 26-133, 26-147, 26-180, 26-259, 26-295, 26-296, 26-304, 26-305, 26-450, 26-532, 26-578, 26-593, 26-648, 28-497, 28-600, 31-373, 33-295, 34-646, 34-657, 37-452, 37-637, 37-676, 43-344, 51-305, 53-850, 89-583, 103-644, 105-576, 124-394, 143-305, 160-653, 164-470, 233-462, 234-848, 249-1031, 255-476, 255-541, 258-638, 278-490, 278-495, 289-925, 300-592, 306-700,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
39	306-719, 306-816, 306-879, 306-887, 306-888, 306-900, 306-945, 306-947, 306-961, 306-986, 307-789, 307-824, 307-831, 307-840, 307-841, 307-855, 307-858, 307-865, 307-891, 307-899, 307-932, 307-944, 307-945, 307-952, 307-955, 307-978, 307-981, 307-1005, 308-884, 309-479, 309-977, 318-568, 322-563, 325-548, 328-641, 329-546, 346-933, 347-1045, 350-1032, 353-830, 354-1015, 364-607, 364-639, 376-1135, 413-673, 413-1012, 415-703, 415-980, 416-895, 422-1121, 423-833, 427-824, 427-933, 452-722, 465-1170, 507-757, 521-1242, 542-803, 547-1187, 563-891, 580-1181, 591-881, 592-889, 627-1000, 630-844, 653-874, 655-935, 663-886, 673-914, 683-953, 690-949, 693-925, 693-927, 693-1319, 693-1433, 693-1461, 856-1576, 884-1164, 884-1169, 896-1150, 939-1167, 947-1229, 947-1414, 947-1415, 947-1438, 947-1456, 947-1457, 947-1523, 947-1589, 947-1591, 947-1611, 947-1635, 947-1645, 952-1179, 953-1444, 1130-1667, 1170-1526, 1275-1830, 1352-1419, 1377-1830, 1450-1830, 1515-1831, 1534-1788
40/7982285CB1/ 1524	1-292, 2-421, 44-805, 57-109, 59-624, 61-272, 61-656, 61-697, 61-698, 67-216, 67-247, 68-749, 74-437, 74-544, 74-573, 74-579, 74-676, 74-679, 77-756, 81-388, 82-340, 83-388, 84-658, 89-352, 90-309, 90-530, 91-306, 91-313, 91-674, 93-216, 95-356, 95-432, 95-435, 95-500, 95-501, 95-534, 95-545, 95-548, 96-767, 97-919, 98-505, 98-754, 100-346, 101-635, 102-960, 118-368, 123-544, 123-617, 151-787, 154-409, 162-780, 203-880, 224-388, 239-815, 305-388, 369-1003, 392-853, 463-1070, 500-1100, 501-761, 519-1155, 536-565, 539-672, 608-1261, 628-1186, 678-1175, 697-1241, 804-1446, 839-1111, 845-1478, 916-1524, 929-1088, 938-1524, 957-1226, 957-1473, 972-1418, 1036-1085
41/7758505CB1/ 2973	1-635, 119-364, 119-742, 119-752, 244-735, 244-753, 244-755, 244-869, 666-755, 750-1300, 754-2973, 961-1142, 1033-1562, 1033-1938, 1034-1689, 1185-2973, 1327-1934, 1335-1847, 1353-1947, 1796-2305, 1937-1966
42/6885756CB1/ 2126	1-1789, 401-1789, 402-1789, 1674-2126

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
43/7500748CB1/ 1973	1-482, 1-657, 1-671, 1-686, 2-597, 2-657, 8-256, 9-240, 11-361, 14-257, 14-266, 14-278, 15-276, 15-321, 15-529, 15-579, 19-271, 19-281, 19-284, 19-286, 20-273, 20-282, 20-285, 20-287, 20-288, 20-315, 22-242, 22-267, 22-277, 23-255, 23-294, 23-304, 23-321, 34-284, 36-642, 37-649, 42-273, 60-299, 93-228, 240-392, 255-489, 283-828, 313-1049, 361-923, 367-524, 367-540, 367-559, 405-557, 502-842, 520-766, 701-934, 726-953, 732-974, 755-1182, 758-920, 758-1015, 777-1038, 779-1029, 781-1035, 805-1049, 838-977, 863-1049, 897-1080, 934-1239, 940-1208, 1070-1360, 1073-1872, 1077-1708, 1080-1629, 1086-1472, 1090-1305, 1102-1966, 1113-1395, 1115-1314, 1119-1351, 1119-1784, 1126-1765, 1133-1525, 1136-1539, 1138-1426, 1138-1912, 1146-1724, 1166-1438, 1166-1598, 1186-1444, 1191-1437, 1205-1541, 1210-1930, 1210-1944, 1220-1552, 1221-1973, 1226-1479, 1235-1488, 1235-1840, 1236-1822, 1241-1502, 1242-1536, 1242-1861, 1256-1494, 1256-1523, 1256-1813, 1259-1460, 1260-1854, 1262-1460, 1265-1722, 1279-1589, 1282-1909, 1287-1929, 1296-1834, 1299-1876, 1300-1927, 1303-1605, 1305-1787, 1307-1565, 1309-1555, 1316-1788, 1316-1953, 1317-1805, 1318-1718, 1319-1929, 1327-1594, 1333-1583, 1333-1737, 1334-1554, 1335-1590, 1336-1576, 1336-1588, 1339-1562, 1342-1604, 1345-1966, 1346-1723, 1349-1444, 1353-1808, 1356-1851, 1357-1958, 1357-1968, 1359-1631, 1361-1939, 1362-1956, 1363-1643, 1366-1973, 1367-1638, 1375-1637, 1387-1967, 1394-1679, 1394-1695, 1395-1631, 1397-1665, 1404-1621, 1404-1641, 1404-1801, 1406-1550, 1406-1661, 1406-1682, 1406-1967, 1411-1617, 1418-1669, 1418-1852, 1422-1953, 1431-1968, 1435-1681, 1441-1845, 1454-1968, 1465-1949, 1467-1952, 1467-1973, 1471-1775, 1475-1726, 1481-1938, 1493-1763, 1499-1781, 1504-1769, 1529-1624, 1529-1933, 1531-1794, 1539-1934, 1540-1930, 1555-1814, 1561-1898, 1628-1829, 1664-1901 1-190, 1-1846, 22-670, 158-358, 190-408, 190-416, 190-456, 190-507, 190-523, 190-661, 190-662, 190-697, 190-724, 190-775, 190-794, 194-626, 195-801, 199-772, 201-447, 201-598, 208-729, 218-472, 221-813, 223-780, 224-818, 229-767, 235-834, 236-764, 239-830, 242-511, 246-866, 253-944, 259-915, 261-492, 264-506, 266-411, 266-526, 267-566, 271-542, 271-883, 278-873, 281-773, 284-651, 285-916, 297-704, 309-579, 316-859, 335-581, 339-702, 339-733, 340-835, 342-611, 342-889, 347-702, 353-952, 359-932,
44/7500749CB1/ 1884	

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
44	<p>365-629, 365-640, 368-616, 370-859, 372-955, 374-1118, 376-668, 381-793, 382-615, 382-640, 382-652, 385-670, 385-784, 386-1210, 387-589, 387-883, 390-975, 392-513, 395-651, 398-944, 407-634, 407-702, 413-655, 413-1003, 433-1091, 436-730, 439-601, 439-696, 441-1065, 452-913, 452-961, 452-985, 452-1193, 452-1197, 452-1218, 452-1274, 452-1303, 455-1162, 458-719, 459-639, 460-710, 462-716, 469-937, 474-1042, 477-1074, 486-734, 489-777, 491-1090, 493-1023, 493-1066, 496-1158, 497-959, 509-1018, 519-658, 522-1167, 523-1090, 532-1061, 542-1079, 544-772, 549-811, 551-838, 561-794, 563-986, 564-1169, 565-814, 565-1227, 571-875, 573-1271, 574-1256, 578-727, 583-1204, 588-824, 588-831,</p> <p>592-1213, 593-887, 603-1111, 606-887, 611-950, 611-1245, 612-895, 614-859, 620-1228, 625-870, 629-755, 629-817, 639-1345, 642-922, 644-884, 649-1310, 650-1225, 652-920, 653-1191, 654-899, 656-995, 661-933, 665-934, 665-942, 670-902, 671-1164, 673-925, 675-948, 675-952, 675-963, 675-1208, 678-908, 680-1319, 683-935, 683-1262, 683-1279, 693-988, 697-858, 697-943, 700-908, 708-1363, 710-978, 710-1142, 713-1331, 715-1291, 719-978, 720-972, 720-974, 720-1018, 723-1035, 726-926, 726-941, 726-988, 728-887, 734-1365, 736-983, 742-973, 744-978, 744-1316, 745-1332, 746-923, 746-1032, 770-1381, 773-1076, 783-1090, 785-1051, 788-977, 790-1013, 793-1122, 793-1329, 798-1051, 798-1080, 798-1093, 800-1072, 800-1421, 802-1451, 803-893, 803-1116, 814-1344, 819-1405, 822-1523, 823-1095, 831-1088, 831-1328, 831-1345, 833-1264, 837-1341, 837-1346, 840-965, 842-1126, 842-1480, 843-1430, 848-1415, 848-1492,</p> <p>852-1388, 861-1095, 862-1115, 876-1054, 886-1130, 895-1387, 901-1158, 906-1155, 908-1191, 911-1614, 912-1114, 913-1604, 915-1053, 922-1572, 925-1189, 925-1387, 926-1618, 932-1045, 932-1178, 939-1207, 942-1458, 943-1512, 952-1242, 953-1195, 953-1638, 955-1754, 959-1590, 962-1511, 965-1177, 966-1213, 968-1354, 969-1254, 971-1258, 972-1187, 984-1848, 993-1798, 995-1277, 997-1196, 1001-1233, 1001-1666, 1003-1235, 1003-1402, 1007-1601, 1008-1647, 1015-1407, 1018-1421, 1020-1281, 1020-1308, 1020-1515, 1020-1794, 1021-1254, 1028-1606, 1040-1316, 1042-1288, 1043-1349, 1045-1574, 1045-1669, 1048-1320, 1048-1480, 1055-1316, 1055-1318, 1060-1282, 1068-1326, 1073-1319, 1080-1629, 1084-1340, 1087-1423, 1087-1801, 1088-1145, 1092-1812, 1092-1826, 1097-1338, 1102-1330, 1102-1434, 1103-1855, 1105-1322, 1105-1436, 1108-1361, 1110-1663, 1112-1369, 1117-1370, 1117-1413, 1117-1722, 1118-1704, 1122-1403, 1122-1416, 1122-1785, 1123-1384, 1124-1418, 1124-1743,</p>

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
44	<p>1137-1376, 1138-1371, 1138-1376, 1138-1405, 1138-1695, 1141-1342, 1142-1736, 1144-1342, 1146-1820, 1147-1604, 1153-1721, 1159-1803, 1161-1471, 1163-1794, 1164-1791, 1169-1811, 1172-1806, 1178-1716, 1180-1837, 1181-1758, 1182-1777, 1182-1809, 1185-1487, 1187-1669, 1189-1447, 1191-1437, 1195-1761, 1198-1670, 1198-1835, 1199-1687, 1200-1600, 1200-1881, 1201-1811, 1201-1879, 1208-1446, 1209-1476, 1215-1465, 1215-1503, 1215-1619, 1216-1436, 1217-1472, 1218-1458, 1218-1470, 1220-1806, 1221-1444, 1222-1849, 1224-1486, 1224-1487, 1224-1492, 1227-1848, 1228-1605, 1231-1326, 1235-1690, 1238-1733, 1239-1840, 1239-1850, 1241-1513, 1243-1821, 1244-1838, 1245-1525, 1248-1863, 1249-1520, 1253-1847, 1255-1860, 1257-1519, 1262-1778, 1265-1863, 1266-1560, 1269-1849, 1276-1561, 1276-1577, 1276-1876, 1277-1513, 1279-1547, 1286-1503, 1286-1523, 1286-1586,</p> <p>1286-1683, 1288-1432, 1288-1485, 1288-1543, 1288-1564, 1288-1849, 1288-1883, 1293-1499, 1295-1882, 1300-1551, 1300-1734, 1304-1835, 1310-1850, 1313-1881, 1317-1563, 1318-1882, 1323-1727, 1324-1470, 1334-1583, 1336-1850, 1338-1581, 1341-1622, 1341-1880, 1343-1809, 1347-1666, 1347-1831, 1348-1565, 1349-1572, 1349-1834, 1349-1864, 1353-1657, 1354-1608, 1357-1608, 1359-1880, 1363-1820, 1375-1645, 1381-1663, 1382-1821, 1382-1850, 1385-1847, 1386-1651, 1388-1884, 1392-1851, 1394-1636, 1396-1850, 1399-1883, 1402-1873, 1404-1848, 1409-1853, 1411-1506, 1411-1815, 1413-1676, 1414-1848, 1421-1816, 1422-1812, 1424-1858, 1426-1853, 1432-1850, 1433-1807, 1433-1849, 1433-1853, 1436-1849, 1436-1855, 1437-1696, 1438-1722, 1439-1849, 1441-1871, 1442-1849, 1442-1871, 1443-1780, 1443-1850, 1443-1862, 1443-1863, 1446-1849, 1448-1858, 1450-1856, 1452-1847, 1452-1855, 1453-1849, 1454-1753, 1455-1822, 1456-1734, 1456-1849, 1458-1827, 1459-1849, 1465-1859, 1468-1638, 1471-1883, 1472-1756, 1481-1696, 1481-1849, 1485-1850, 1486-1717, 1486-1867, 1488-1850, 1492-1883, 1495-1849, 1502-1524, 1502-1821, 1506-1849, 1510-1711, 1510-1853, 1515-1735, 1515-1850, 1524-1849, 1526-1805, 1526-1849, 1528-1812, 1528-1850, 1531-1850, 1532-1849, 1533-1851, 1537-1737, 1537-1796, 1544-1833, 1545-1760, 1545-1770, 1545-1790, 1545-1853, 1546-1783, 1547-1831, 1547-1844, 1547-1878, 1549-1830, 1550-1830, 1558-1849, 1560-1855, 1594-1793, 1596-1884, 1602-1855, 1604-1853, 1605-1849, 1648-1865, 1649-1883, 1654-1883, 1668-1872, 1682-1849, 1689-1883, 1692-1850, 1699-1868, 1711-1877, 1713-1849, 1734-1849, 1735-1863, 1736-1850, 1738-1883, 1753-1812, 1753-1848, 1753-1849, 1753-1850, 1758-1850, 1760-1864, 1765-1849</p>



Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
45/7503401CB1/ 1581	1-255, 1-328, 1-359, 1-436, 4-332, 10-295, 12-362, 13-317, 20-641, 21-272, 21-281, 21-436, 22-239, 22-616, 22-644, 26-341, 27-286, 27-287, 31-265, 34-311, 34-328, 35-299, 35-307, 37-327, 39-154, 46-289, 81-395, 85-380, 112-366, 143-413, 144-411, 146-399, 149-436, 178-436, 179-436, 185-947, 189-436, 237-429, 427-668, 432-755, 433-1284, 458-918, 487-650, 487-712, 489-995, 490-1360, 493-743, 494-991, 494-1134, 503-783, 507-910, 516-772, 517-753, 518-731, 520-763, 522-814, 522-1149, 524-703, 532-1024, 541-699, 543-807, 557-688, 559-818, 568-788, 569-1290, 586-849, 596-874, 599-882, 623-735, 642-904, 665-1282, 670-904, 676-922, 676-931, 678-1531, 681-1532, 693-1512, 694-899, 694-1059, 699-867, 701-941, 704-865, 704-997, 716-878, 716-1207, 717-1264, 725-1155, 734-1061, 737-1514, 741-1355, 743-1241, 748-1563, 759-1194, 770-1224, 775-1547, 781-1254, 788-1085, 788-1278, 794-1341, 800-1563, 801-1236, 807-1450, 820-1581, 822-1502, 835-1528, 836-1563, 839-1565, 841-1388, 841-1569, 846-1138, 846-1221, 850-1579, 861-1427, 863-1171, 865-1568, 865-1575, 874-1568, 875-1518, 878-1569, 882-1101, 883-1569, 884-1508, 885-1123, 888-1568, 897-1569, 898-1234, 901-1569, 906-1577, 919-1569, 925-1568, 927-1579, 932-1578, 933-1187, 933-1519, 937-1554, 938-1569, 948-1182, 948-1223, 950-1261, 950-1553, 952-1487, 954-1431, 954-1569, 955-1235, 960-1548, 964-1564, 967-1151, 970-1210, 973-1489, 975-1232, 980-1575, 985-1249, 989-1576, 991-1497, 992-1248, 992-1577, 992-1579, 993-1291, 996-1173, 996-1579, 998-1429, 1004-1224, 1010-1569, 1014-1261, 1019-1242, 1020-1248, 1028-1579, 1036-1579, 1042-1579, 1043-1523, 1050-1574, 1050-1579, 1055-1526, 1065-1579, 1068-1271, 1069-1579, 1076-1557, 1076-1575, 1078-1499, 1079-1350, 1081-1354, 1091-1306, 1092-1364, 1092-1565, 1094-1280, 1094-1579, 1096-1579, 1101-1566, 1102-1564, 1102-1579, 1104-1579, 1106-1569, 1109-1416, 1109-1568, 1109-1569, 1115-1567, 1115-1569, 1116-1566, 1121-1565, 1121-1567, 1126-1579, 1131-1567, 1132-1407, 1133-1581, 1134-1399, 1134-1420, 1136-1566, 1136-1579, 1137-1579, 1139-1564, 1139-1569, 1143-1567, 1143-1569, 1143-1579, 1144-1569, 1145-1427, 1146-1564, 1146-1579, 1148-1570, 1149-1564, 1150-1565, 1151-1562, 1152-1563, 1152-1565, 1154-1564, 1157-1569, 1157-1579, 1158-1487, 1158-1579.

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
45	1159-1567, 1164-1579, 1167-1564, 1168-1445, 1170-1564, 1173-1569, 1180-1451, 1181-1579, 1182-1569, 1183-1560, 1184-1564, 1184-1565, 1186-1579, 1186-1581, 1187-1564, 1187-1579, 1188-1563, 1188-1564, 1188-1570, 1189-1565, 1193-1579, 1195-1570, 1198-1562, 1206-1499, 1206-1569, 1209-1560, 1209-1579, 1214-1564, 1216-1473, 1220-1422, 1220-1567, 1221-1579, 1223-1493, 1223-1501, 1225-1579, 1226-1507, 1240-1565, 1246-1565, 1247-1564, 1254-1552, 1260-1567, 1261-1529, 1262-1567, 1268-1571, 1269-1564, 1272-1564, 1272-1565, 1277-1547, 1277-1562, 1278-1567, 1279-1485, 1288-1530, 1289-1564, 1289-1565, 1305-1581, 1308-1567, 1316-1572, 1316-1577, 1320-1564, 1322-1579, 1334-1574, 1355-1564, 1355-1565, 1374-1562, 1374-1564, 1376-1579, 1387-1569, 1392-1578, 1397-1564, 1414-1567, 1414-1571, 1427-1579, 1434-1564, 1448-1564, 1449-1564, 1469-1564
46/7503485CBI/ 1996	1-282, 9-1724, 29-300, 30-292, 30-293, 34-305, 34-326, 52-266, 52-299, 52-325, 53-330, 53-338, 54-317, 54-328, 54-330, 54-335, 55-314, 57-303, 57-321, 57-323, 57-325, 57-328, 59-314, 60-268, 60-285, 60-305, 60-315, 60-316, 60-318, 60-321, 60-338, 60-342, 61-342, 62-272, 62-319, 63-308, 65-337, 66-298, 70-323, 72-330, 73-325, 106-287, 142-429, 176-668, 176-826, 238-520, 242-503, 242-532, 242-782, 242-873, 242-912, 325-958, 340-911, 348-649, 365-597, 371-936, 376-549, 377-641, 378-640, 387-704, 394-671, 397-586, 443-720, 443-1010, 447-705, 452-897, 455-966, 466-807, 472-1026, 486-610, 486-817, 505-1117, 506-651, 506-803, 528-980, 528-981, 529-1048, 537-1122, 548-776, 549-810, 550-859, 551-807, 556-1132, 560-778, 562-724, 593-858, 610-669, 611-875, 618-902, 619-921, 620-1249, 633-868, 641-876, 643-898, 645-911, 646-842, 646-900, 647-895, 656-864, 668-881, 676-1065, 676-1223, 677-916, 677-1012, 684-881, 684-965, 687-1135, 688-865, 688-968, 695-938, 702-1312, 708-970, 715-1343, 721-992, 735-949, 745-1336, 750-1362, 755-992, 757-1263, 777-1057, 777-1352, 788-999, 799-1062, 820-1271, 830-1087, 855-1092, 855-1298, 856-986, 856-1288, 869-1113, 871-1342, 873-1159, 879-1274, 880-1147, 883-1443, 885-1486, 890-1696, 902-1529, 908-1135, 910-1197, 918-1068, 918-1084, 918-1105, 918-1145, 918-1168, 918-1169, 920-1454, 921-1719, 924-1674, 939-1191, 941-1086, 943-1103, 944-1135, 944-1174, 944-1184, 944-1198, 955-1431, 962-1215, 965-1226, 974-1224, 975-1600, 979-1679, 987-1260, 987-1536, 992-1308, 993-1268, 993-1314, 995-1254, 997-1732, 999-1275, 1004-1209, 1007-1458, 1007-1651, 1021-1279, 1022-1715, 1034-1256, 1037-1293, 1045-1249, 1046-1752, 1054-1661, 1055-1228,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
46	1055-1262, 1055-1322, 1055-1324, 1056-1333, 1059-1655, 1066-1511, 1068-1718, 1069-1740, 1084-1302, 1086-1720, 1086-1748, 1093-1336, 1093-1714, 1095-1379, 1097-1380, 1100-1707, 1111-1344, 1118-1252, 1119-1696, 1123-1747, 1125-1418, 1125-1432, 1126-1386, 1126-1666, 1132-1645, 1132-1723, 1135-1399, 1139-1471, 1142-1716, 1148-1742, 1158-1388, 1160-1403, 1160-1721, 1164-1748, 1168-1410, 1172-1434, 1177-1746, 1184-1770, 1187-1724, 1197-1458, 1197-1469, 1211-1439, 1211-1468, 1211-1486, 1212-1742, 1234-1747, 1235-1742, 1240-1600, 1244-1749, 1247-1747, 1248-1745, 1249-1730, 1250-1724, 1253-1648, 1253-1745, 1254-1728, 1254-1745, 1258-1720, 1260-1730, 1260-1744, 1261-1729, 1265-1703, 1267-1482, 1268-1758, 1269-1745, 1271-1692, 1272-1522, 1272-1728, 1275-1728, 1277-1730, 1277-1731, 1278-1728, 1281-1519, 1283-1508, 1288-1727, 1290-1723,
	1292-1720, 1293-1719, 1294-1720, 1296-1720, 1296-1725, 1296-1728, 1297-1720, 1297-1732, 1298-1673, 1298-1725, 1298-1752, 1300-1727, 1301-1724, 1301-1728, 1303-1729, 1303-1731, 1304-1720, 1305-1728, 1307-1729, 1308-1706, 1308-1727, 1308-1741, 1308-1747, 1309-1765, 1316-1720, 1319-1720, 1323-1720, 1325-1741, 1328-1727, 1330-1735, 1331-1720, 1333-1736, 1336-1732, 1337-1747, 1341-1677, 1346-1722, 1354-1727, 1356-1727, 1358-1500, 1361-1646, 1361-1732, 1367-1672, 1370-1503, 1370-1720, 1376-1718, 1377-1720, 1378-1720, 1382-1519, 1391-1734, 1409-1720, 1415-1663, 1421-1650, 1421-1672, 1421-1676, 1421-1759, 1422-1808, 1424-1727, 1433-1720, 1438-1709, 1439-1721, 1440-1725, 1453-1721, 1456-1714, 1457-1996, 1459-1738, 1459-1741, 1466-1715, 1468-1727, 1473-1727, 1475-1714, 1476-1728, 1479-1705, 1483-1729, 1512-1729, 1519-1729, 1524-1729, 1540-1729, 1550-1721, 1550-1783, 1560-1727, 1598-1721, 1610-1750, 1624-1724, 1642-1727
47/7504076CB1/ 1232	1-695, 16-646, 17-583, 27-487, 27-561, 46-668, 61-611, 83-448, 91-611, 159-614, 258-632, 258-693, 259-679, 259-845, 260-626, 260-632, 285-698, 285-848, 288-745, 298-590, 323-790, 361-791, 368-805, 373-832, 387-815, 393-873, 403-639, 429-774, 488-655, 497-746, 497-833, 497-1043, 506-813, 533-663, 543-783, 578-773, 583-1232
48/7500926CB1/ 810	1-537, 11-467, 11-475, 22-190, 22-285, 22-295, 22-299, 22-400, 22-458, 22-468, 22-475, 22-478, 22-484, 23-386, 23-482, 23-484, 26-379, 27-276, 27-434, 38-468, 41-810, 69-355, 164-471, 257-549, 280-469, 299-481, 299-546, 391-558, 405-809

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
49/7503216CB1/ 2625	1-586, 1-2598, 310-913, 335-913, 391-1023, 393-919, 393-928, 404-660, 580-884, 588-793, 928-1381, 930-1254, 930-1349, 930-1392, 931-1349, 937-1589, 953-1492, 953-1568, 955-1604, 959-1482, 972-1519, 975-1620, 976-1402, 996-1594, 1006-1227, 1009-1465, 1036-1732, 1043-1473, 1044-1640, 1052-1327, 1078-1345, 1100-1597, 1118-1603, 1136-1418, 1160-1749, 1167-1601, 1173-1548, 1184-1722, 1188-1417, 1193-1665, 1193-1736, 1196-1640, 1204-1489, 1222-1653, 1238-1506, 1246-2182, 1261-1676, 1267-2183, 1281-1753, 1291-1482, 1313-1789, 1318-1965, 1325-1792, 1337-1988, 1340-1678, 1341-1912, 1350-1747, 1352-2183, 1356-1633, 1358-1913, 1401-1969, 1408-1968, 1413-2183, 1418-1891, 1422-1975, 1434-1831, 1434-1904, 1434-2183, 1440-1995, 1451-1947, 1451-2183, 1452-2183, 1459-2108, 1479-1921, 1507-2183, 1536-1694, 1551-2075, 1579-2155, 1580-1809, 1580-1878, 1580-2183, 1584-1890, 1610-1875, 1614-1839, 1614-1891, 1614-1899, 1614-1929, 1614-1935, 1622-2156, 1625-1870, 1626-1918, 1640-2083, 1655-1962, 1657-1893, 1665-1935, 1675-1919, 1678-2244, 1678-2302, 1680-2112, 1719-2031, 1719-2341, 1721-2314, 1726-1975, 1735-2131, 1744-1989, 1746-1965, 1748-2130, 1748-2307, 1751-2240, 1758-1962, 1763-1977, 1763-2349, 1778-1935, 1797-2249, 1799-2327, 1816-2358, 1847-2137, 1866-2113, 1872-2592, 1879-2136, 1879-2472, 1881-2091, 1888-2475, 1902-2551, 1903-2134, 1911-2149, 1926-2222, 1926-2578, 1942-2124, 1943-2154, 1946-2559, 1946-2580, 1953-2218, 1953-2369, 1958-2581, 1970-2239, 1972-2207, 1982-2208, 2190-2612, 2275-2625, 2455-2598
50/7503233CB1/ 2432	1-253, 1-272, 1-2432, 15-311, 18-288, 19-257, 21-311, 23-244, 23-275, 24-311, 24-335, 27-335, 28-285, 33-294, 33-299, 37-271, 37-279, 38-305, 38-313, 44-313, 44-336, 48-304, 48-315, 51-335, 58-335, 79-319, 144-688, 292-778, 331-590, 331-787, 331-820, 331-927, 331-958, 331-959, 331-961, 331-965, 333-562, 334-961, 334-963, 335-786, 340-589, 342-945, 349-625, 349-630, 349-631, 353-852, 362-980, 366-747, 379-830, 381-975, 410-981, 427-1006, 442-1051, 448-948, 455-715, 456-705, 462-961, 467-813, 469-1146, 475-666, 480-989, 480-1063, 482-924, 488-683, 489-747, 494-719, 494-762, 522-738, 523-1028, 547-864, 547-1184, 556-1349, 564-1190, 571-745, 577-1154, 581-848, 593-1148, 609-1033, 615-915, 618-826, 618-1183, 621-1098, 623-1174, 623-1261, 631-1167, 632-881, 635-831, 635-1107, 636-986, 647-1178, 650-1371,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
50	<p>653-989, 656-1107, 665-899, 669-923, 669-1347, 682-1275, 692-939, 693-953, 694-1456, 706-1037, 712-984, 714-902, 719-1121, 723-1367, 733-985, 736-1012, 747-1034, 747-1273, 750-1452, 752-1001, 754-1015, 754-1314, 756-1139, 756-1140, 758-1299, 771-1333, 776-1065, 777-1360, 778-1260, 780-1413, 782-1076, 786-999, 798-1081, 805-1423, 806-1254, 835-1249, 838-1336, 851-1470, 852-1455, 854-1356, 854-1407, 855-1499, 856-1510, 857-1373, 860-1415, 865-1106, 865-1118, 868-1517, 884-1131, 886-1068, 891-1602, 894-1193, 906-1383, 916-1360, 917-1396, 918-1523, 925-1345, 930-1104, 938-1231, 946-1425, 948-1425, 954-1183, 966-1412, 971-1534, 973-1407, 976-1257, 977-1219, 987-1728, 992-1548, 993-1245, 993-1250, 994-1346, 995-1297, 997-1598, 1005-1278, 1006-1236, 1006-1916, 1027-1414, 1028-1414, 1030-1506, 1034-1491, 1035-1597, 1053-1253, 1054-1345, 1059-1448,</p> <p>1063-1261, 1069-1423, 1070-1485, 1073-1231, 1074-1499, 1083-1683, 1099-1750, 1102-1401, 1103-1382, 1109-1284, 1114-1698, 1129-1414, 1141-1232, 1148-1635, 1154-1610, 1155-1696, 1157-1660, 1157-1785, 1158-1692, 1166-1839, 1174-1616, 1176-1492, 1186-1372, 1186-1661, 1190-1800, 1191-2008, 1192-1430, 1193-1765, 1202-1781, 1209-1735, 1209-1809, 1210-1451, 1213-1720, 1217-1431, 1219-1507, 1223-1331, 1224-1454, 1224-1467, 1230-1680, 1236-1936, 1241-1489, 1245-1808, 1258-1603, 1259-1517, 1264-1757, 1277-1907, 1289-1627, 1289-1864, 1292-1527, 1293-1702, 1295-1585, 1298-1621, 1299-1746, 1300-1584, 1308-1548, 1309-1833, 1309-2098, 1310-1851, 1312-1589, 1313-1902, 1314-1882, 1315-1809, 1317-1426, 1319-1765, 1326-1523, 1328-1447, 1351-1864, 1351-1977, 1354-1921, 1356-1941, 1366-2008, 1369-1844, 1377-1928, 1383-1792, 1389-1999, 1390-1890, 1394-2056,</p> <p>1399-2162, 1402-1934, 1402-2041, 1403-2087, 1404-1907, 1415-1684, 1419-1907, 1421-2365, 1422-1968, 1422-2036, 1422-2049, 1423-1632, 1423-2066, 1426-1846, 1426-1974, 1432-2046, 1435-1953, 1437-2065, 1456-1902, 1456-1904, 1458-1587, 1459-1692, 1465-1736, 1467-1724, 1468-1769, 1469-1763, 1472-1701, 1472-1758, 1472-1839, 1472-2106, 1475-1768, 1478-1988, 1479-1747, 1482-1704, 1482-1758, 1483-2082, 1489-1930, 1493-2172, 1496-2024, 1507-1804, 1513-1748, 1514-2272, 1521-2025, 1525-1773, 1526-1747, 1528-1744, 1528-2007, 1528-2125, 1531-2113, 1535-2158, 1544-2377, 1553-1844, 1557-2110, 1559-2187, 1565-2229, 1566-1837, 1573-1824, 1574-1757, 1579-1816, 1579-1854, 1579-1870, 1579-2156, 1580-1902, 1583-2032, 1589-1987, 1590-2159, 1591-2160, 1591-2206, 1598-1869, 1599-2077, 1600-1851, 1600-2173, 1604-1991, 1607-1889, 1608-2053, 1615-1872, 1624-1869, 1624-1874, 1625-1874, 1626-1840, 1626-1864, 1626-1866, 1626-1882, 1628-1869, 1632-1924,</p>

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
50	1638-1879, 1645-2136, 1647-2103, 1647-2247, 1649-1908, 1651-1922, 1651-2191, 1651-2206, 1652-2236, 1653- 1968, 1656-2190, 1657-1902, 1673-1976, 1673-2245, 1674-1930, 1678-1987, 1678-1988, 1682-1883, 1691-1899, 1694-1841, 1696-1975, 1696-1983, 1696-2331, 1702-1922, 1702-2291, 1704-1996, 1705-1946, 1708-1954, 1708- 1977, 1709-1926, 1709-1961, 1710-2280, 1712-1904, 1712-2118, 1715-2189, 1721-2129, 1727-2251, 1742-2182, 1748-1990, 1749-2028, 1755-1941, 1756-2115, 1763-1988, 1768-2272, 1768-2373, 1773-2339, 1777-2054, 1778- 2020, 1781-2313, 1788-2288, 1789-2000, 1790-2045, 1798-2052, 1804-2086, 1807-2366, 1812-2097, 1813-2039, 1813-2404, 1825-2053, 1828-1993, 1829-2122, 1832-2069, 1832-2077, 1835-2353, 1838-2086, 1838-2126, 1840- 2024, 1843-2120, 1844-2398, 1849-2362, 1868-2144, 1872-2160, 1873-2150, 1874-2119, 1877-2127, 1885-2420, 1886-2145, 1887-2170, 1894-2149, 1902-2174, 1907-2151, 1907-2157, 1907-2197, 1910-2123, 1912-2231, 1917-2098, 1917- 2127, 1917-2169, 1917-2173, 1917-2198, 1917-2359, 1917-2398, 1919-2146, 1935-2132, 1935-2192, 1940-2122, 1940-2147, 1942-2190, 1944-2181, 1953-2266, 1959-2266, 1962-2202, 1962-2276, 1962-2279, 1963-2079, 1963- 2116, 1970-2214, 1976-2430, 1977-2088, 1982-2214, 1982-2217, 1984-2225, 1984-2229, 1984-2232, 1984-2240, 1984-2255, 1984-2260, 1987-2254, 1995-2123, 1998-2266, 2001-2100, 2005-2216, 2021-2296, 2029-2313, 2030- 2127, 2033-2151, 2035-2282, 2036-2243, 2044-2265, 2045-2265, 2080-2267, 2095-2353, 2099-2337, 2099-2349, 2102-2337, 2104-2337, 2107-2321, 2119-2398, 2121-2278, 2121-2285, 2121-2336, 2126-2349, 2127-2321, 2132- 2323, 2132-2379, 2133-2381, 2133-2400, 2191-2389, 2205-2343, 2260-2398 1-473, 1-666, 27-604, 48-658, 151-666, 151-688, 151-800, 173-594, 184-666, 216-765, 298-910, 331-945, 378-666, 408-668, 408-678, 506-1234, 539-868, 578-1173, 656-1233, 681-1294, 714-1281, 725-1335, 727-1393, 749-1390, 759-1390, 762-1360, 787-1155, 802-1329, 803-869, 803-885, 822-1245, 879-1266, 883-1138, 883-1221, 883-1334, 883-1359, 883-1381, 883-1447, 883-1462, 883-1514, 883-1560, 929-1343, 929-1344, 929-1355, 929-1490, 931- 1063, 932-1303, 952-1819, 955-1115, 955-1329, 973-1630, 994-1629, 1032-1710, 1049-1638, 1064-1723, 1068- 1677, 1081-1686, 1114-1263, 1121-2063, 1126-1744, 1182-1724, 1255-1894, 1266-1907, 1274-1720, 1278-1573, 1298-1891, 1313-1615, 1322-1546, 1322-1601,

51/7726576CBI/  
3969

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
51	1342-1997, 1351-1581, 1352-1977, 1356-1816, 1366-1653, 1368-1877, 1379-1996, 1387-1989, 1409-1769, 1429- 1917, 1460-1989, 1463-2071, 1468-1904, 1468-1919, 1477-1884, 1478-1919, 1486-2070, 1500-2011, 1552-2282, 1561-1786, 1561-1987, 1597-1665, 1639-2303, 1646-2253, 1699-1964, 1712-2264, 1714-2013, 1714-2269, 1724- 2379, 1766-2368, 1775-2324, 1829-2379, 1864-2459, 1864-2521, 1946-2275, 1978-2445, 2096-2303, 2096-2643, 2127-2401, 2183-2525, 2244-2496, 2244-2497, 2344-2601, 2426-2952, 2466-2716, 2468-3032, 2472-3052, 2537- 2659, 2537-3139, 2540-2777, 2540-2782, 2556-3171, 2560-2797, 2623-3268, 2626-3258, 2630-3136, 2638-2875, 2648-3269, 2731-2866, 2731-3006, 2731-3213, 2738-3283, 2774-3015, 2774-3059, 2774-3398, 2779-3380, 2807- 3043, 2832-2958, 2835-3100, 2835-3130, 2854-3116, 2868-3256, 2881-3100, 2881-3114, 2881-3241, 2885-3071, 2885-3134, 2887- 3465, 2894-3139, 2896-3165, 2896-3172, 2897-3465, 2904-3165, 2911-3074, 2933-3199, 2933-3465, 2939-3156, 2939-3366, 2942-3318, 2947-3373, 2957-3465, 2965-3465, 2990-3258, 3008-3465, 3012-3292, 3044-3300, 3047- 3350, 3057-3276, 3087-3340, 3127-3464, 3139-3402, 3153-3411, 3190-3440, 3278-3969, 3283-3465, 3309-3465, 3413-3465, 3442-3465, 3585-3672, 3586-3656, 3586-3667, 3586-3678, 3586-3685, 3586-3706, 3586-3727, 3586- 3771, 3586-3782, 3586-3804, 3586-3848, 3586-3862, 3586-3924, 3586-3928, 3647-3899, 3647-3900, 3747-3928, 3829-3928, 3869-3928, 3875-3928, 11-471, 1-2516, 46-537, 113-743, 113-800, 113-850, 113-871, 114-589, 114-856, 117-857, 124-619, 648-1320, 649- 984, 649-1030, 649-1036, 649-1040, 649-1067, 649-1092, 649-1121, 649-1128, 649-1130, 649-1131, 649-1132, 649- 1135, 649-1147, 649-1149, 649-1167, 649-1174, 651-978, 653-785, 653-1030, 655-1197, 655-1300, 657-1053, 657- 1086, 657-1154, 657-1195, 657-1244, 657-1262, 657-1297, 657-1300, 657-1320, 658-1069, 658-1249, 667-1001, 667-1128, 667-1135, 672-908, 676-1014, 684-1142, 691-826, 691-880, 691-901, 691-1025, 691-1029, 691-1149, 691-1156, 691-1173, 693-891, 700-1103, 710-981, 719-967, 759-906, 762-1255, 766-1309, 767-1000, 767-1133, 793-1322, 839-1255, 840-1345, 864-1328, 954-1265, 956-1265, 1055-1339, 1091-1338, 1178-1348, 1335-1700, 1335-1950, 1336-1614, 1395-1670.
52/7503507CBI/ 2537	

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
52	1415-1647, 1489-1793, 1493-1722, 1493-1742, 1497-1747, 1502-2107, 1533-1784, 1533-1790, 1533-1791, 1533-2058, 1533-2125, 1573-2064, 1579-2048, 1585-2098, 1585-2133, 1598-2059, 1599-2008, 1599-2022, 1632-1891, 1655-1711, 1667-1958, 1668-1955, 1671-1892, 1671-2412, 1726-2201, 1737-2214, 1762-1936, 1780-2083, 1802-2065, 1802-2364, 1840-2147, 1848-2127, 1854-2080, 1869-2109, 1889-1989, 1889-2169, 1917-2165, 1917-2212, 1918-2182, 1924-2206, 1953-2219, 1967-2257, 1969-2156, 1998-2421, 2059-2203, 2059-2377, 2070-2286, 2070-2400, 2104-2315, 2104-2495, 2110-2400, 2110-2534, 2136-2418, 2143-2368, 2160-2427, 2192-2407, 2192-2412, 2205-2443, 2207-2473, 2207-2474, 2255-2502, 2263-2504, 2282-2448, 2282-2516, 2290-2537, 2326-2462 1-471, 1-2505, 46-537, 113-743, 113-743, 113-800, 113-850, 113-871, 114-589, 114-856, 117-857, 124-619, 648-1320, 649-984, 649-1030, 649-1036, 649-1040, 649-1067, 649-1092, 649-1121, 649-1128, 649-1130, 649-1131, 649-1132, 649-1135, 649-1147, 649-1149, 649-1167, 649-1174, 651-978, 653-785, 653-1030, 655-1197, 655-1300, 657-1053, 657-1086, 657-1154, 657-1195, 657-1244, 657-1262, 657-1297, 657-1300, 657-1320, 658-1069, 658-1249, 667-1001, 667-1128, 667-1135, 672-908, 676-1014, 684-1142, 691-826, 691-880, 691-901, 691-1025, 691-1029, 691-1149, 691-1156, 691-1173, 693-891, 700-1103, 710-981, 719-967, 759-906, 762-1255, 766-1309, 767-1000, 767-1113, 793-1322, 839-1255, 840-1395, 864-1328, 954-1265, 956-1265, 968-1603, 1055-1339, 1091-1338, 1151-1385, 1178-1398,
53/7503506CBI/ 2526	1281-1705, 1396-1659, 1396-1689, 1404-1636, 1478-1782, 1482-1711, 1482-1731, 1486-1736, 1491-2096, 1522-1773, 1522-1779, 1522-1780, 1522-2047, 1522-2114, 1562-2053, 1568-2037, 1574-2087, 1574-2122, 1587-2048, 1588-1997, 1588-2011, 1621-1880, 1644-1700, 1656-1947, 1657-1944, 1660-1881, 1660-2401, 1715-2190, 1726-2203, 1751-1925, 1769-2072, 1791-2054, 1791-2353, 1829-2136, 1837-2116, 1843-2069, 1858-2098, 1878-1978, 1878-2158, 1906-2154, 1906-2201, 1907-2171, 1913-2195, 1942-2208, 1956-2246, 1958-2145, 1987-2410, 2048-2192, 2048-2366, 2059-2275, 2059-2389, 2093-2304, 2093-2484, 2099-2389, 2099-2523, 2125-2407, 2132-2357, 2149-2416, 2181-2396, 2181-2401, 2194-2432, 2196-2462, 2196-2463, 2244-2491, 2252-2493, 2271-2437, 2271-2505, 2279-2526, 2315-2451



Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
54/7503509CB1/ 2464	1-471, 1-2443, 46-537, 113-743, 114-589, 124-619, 649-1123, 796-1107, 798-1107, 820-1312, 897-1181, 910-1554, 933-1180, 933-1280, 987-1528, 993-1227, 1002-1302, 1017-1266, 1020-1438, 1106-1627, 1109-1627, 1115-1627, 1126-1627, 1190-1468, 1203-1877, 1211-1635, 1212-1468, 1244-1541, 1322-1597, 1342-1574, 1416-1720, 1420-1649, 1420-1669, 1424-1674, 1429-2034, 1460-1711, 1460-1717, 1460-1718, 1460-1985, 1460-2052, 1500-1991, 1506-1975, 1512-2025, 1512-2060, 1525-1986, 1526-1935, 1526-1949, 1559-1818, 1582-1638, 1594-1885, 1595-1882, 1598-1819, 1598-2339, 1653-2128, 1664-2141, 1689-1863, 1707-2010, 1729-1992, 1729-2291, 1767-2074, 1775-2054, 1781-2007, 1796-2036, 1816-1916, 1816-2096, 1844-2092, 1844-2139, 1845-2109, 1851-2133, 1880-2146, 1894-2184, 1896-2083, 1925-2348, 1986-2130, 1986-2304, 1997-2213, 1997-2327, 2031-2242, 2031-2422, 2037-2327, 2037-2461, 2063-2345, 2070-2295, 2087-2354, 2119-2334, 2119-2339, 2132-2370, 2134-2400, 2134-2401, 2182-2429, 2190-2431, 2209-2375, 2209-2443, 2217-2464, 2253-2389
55/7505800CB1/ 1452	1-211, 1-1447, 3-294, 3-369, 3-401, 3-402, 3-447, 3-517, 3-537, 3-577, 3-603, 3-607, 3-637, 3-639, 3-694, 3-721, 3-846, 6-728, 18-133, 39-503, 40-619, 40-767, 50-736, 51-327, 60-778, 80-285, 99-469, 129-866, 210-438, 210-463, 210-576, 210-735, 227-485, 241-590, 244-516, 248-551, 260-567, 263-396, 265-500, 265-1060, 268-520, 268-835, 271-567, 278-716, 298-551, 304-1167, 315-638, 341-801, 370-533, 370-595, 372-878, 373-1243, 376-626, 377-874, 377-1017, 386-666, 390-793, 399-655, 400-636, 401-614, 403-646, 405-697, 405-1032, 407-586, 415-907, 424-582, 426-690, 440-571, 442-701, 451-671, 452-1173, 469-732, 479-757, 482-765, 506-618, 524-1314, 525-787, 548-1165, 553-787, 559-805, 559-814, 561-1414, 564-1415, 576-1395, 577-782, 577-942, 582-750, 584-824, 587-748, 587-880, 592-882, 592-894, 599-807, 599-916, 599-1090, 600-1147, 601-814, 608-872, 608-1038, 617-944, 620-1397, 624-1238, 626-1124, 627-899, 628-889, 629-941, 630-806, 631-1446, 640-828, 641-918, 642-1077, 643-856, 653-1107, 655-881, 658-1430, 664-1137, 671-968, 671-1161, 677-1224, 683-1446, 684-1119, 690-1333, 703-1443, 705-1385, 709-1451, 715-1452, 718-1411, 719-1446, 722-1448, 722-1451, 724-1271, 724-1452, 725-1002, 729-1021, 729-1104, 733-1452, 734-1449, 743-1452, 744-1310, 746-1054, 748-1450, 748-1451, 757-1451, 758-1401, 759-1446, 761-1452, 765-984, 766-1452, 767-1391, 768-1006, 771-1451, 775-905, 780-1452, 781-1117, 784-1452, 789-1452, 791-1450, 802-1452, 808-1451, 810-1451, 815-1452, 816-1070, 816-1402,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
55	820-1437, 821-1452, 831-1065, 831-1106, 833-1144, 833-1436, 835-1370, 837-1314, 837-1452, 838-1118, 843-1431, 847-1447, 850-1034, 853-1093, 856-1372, 858-1115, 863-1450, 868-1132, 872-1452, 874-1380, 875-1131, 875-1452, 876-1174, 879-1056, 879-1452, 881-1312, 887-1107, 893-1442, 893-1452, 896-1452, 897-1144, 902-1125, 903-1131, 911-1452, 919-1452, 925-1452, 926-1406, 933-1450, 933-1452, 938-1409, 943-1247, 948-1452, 951-1154, 952-1452, 959-1440, 959-1450, 961-1382, 962-1233, 964-1237, 974-1189, 975-1247, 975-1448, 977-1163, 977-1452, 979-1452, 984-1449, 985-1447, 985-1452, 987-1452, 989-1452, 992-1299, 992-1451, 992-1452, 998-1450, 998-1452, 999-1449, 1004-1448, 1004-1450, 1009-1452, 1014-1450, 1015-1290, 1016-1452, 1017-1282, 1017-1303, 1019-1449, 1019-1452, 1020-1452, 1022-1447, 1022-1452, 1024-1284, 1026-1450, 1026-1452, 1027-1452, 1028-1310, 1029-1447, 1029-1452, 1030-1289, 1031-1452, 1032-1447, 1033-1448, 1034-1445, 1035-1446, 1035-1448, 1037-1447, 1040-1452, 1041-1370, 1041-1451, 1042-1450, 1045-1452, 1047-1452, 1050-1447, 1051-1328, 1053-1447, 1056-1452, 1063-1334, 1064-1452, 1065-1452, 1066-1443, 1067-1447, 1067-1448, 1069-1452, 1070-1443, 1070-1447, 1070-1452, 1071-1446, 1071-1447, 1071-1452, 1072-1448, 1076-1452, 1078-1452, 1081-1445, 1082-1318, 1083-1450, 1089-1382, 1089-1437, 1089-1452, 1092-1443, 1092-1452, 1097-1447, 1099-1356, 1103-1305, 1103-1450, 1104-1448, 1106-1376, 1106-1384, 1108-1452, 1109-1390, 1123-1448, 1129-1448, 1130-1447, 1137-1435, 1143-1450, 1144-1412, 1145-1450, 1150-1452, 1151-1452, 1152-1447, 1155-1447, 1155-1448, 1155-1452, 1160-1430, 1160-1445, 1160-1447, 1161-1450, 1162-1368, 1171-1413, 1172-1447, 1172-1448, 1188-1452, 1191-1450, 1199-1452, 1200-1448, 1201-1452, 1203-1447, 1205-1452, 1217-1452, 1238-1447, 1238-1448, 1238-1449, 1242-1452, 1257-1445, 1257-1447, 1259-1452, 1270-1452, 1275-1452, 1280-1447, 1297-1450, 1297-1452, 1310-1452, 1316-1448, 1317-1447, 1319-1445, 1331-1447, 1332-1447, 1352-1447

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
56/750314ICB1/ 1802	1-142, 1-281, 1-286, 1-296, 1-304, 1-339, 1-358, 1-365, 1-369, 1-425, 1-454, 1-480, 1-510, 1-520, 1-532, 1-537, 1-565, 1-568, 3-194, 3-263, 3-281, 3-293, 4-195, 4-202, 4-232, 4-246, 4-264, 4-269, 4-302, 4-484, 4-543, 6-567, 7-249, 7-568, 8-223, 8-239, 8-250, 8-279, 8-315, 10-244, 10-446, 10-501, 10-520, 10-528, 10-1789, 11-266, 11-268, 11-283, 12-243, 12-248, 12-273, 12-292, 12-293, 12-525, 13-278, 13-286, 13-290, 14-129, 14-256, 14-286, 14-307, 19-481, 21-252, 21-256, 21-309, 21-316, 21-365, 21-369, 22-129, 22-143, 22-176, 22-255, 22-291, 22-292, 22-300, 22-301, 22-369, 22-413, 22-446, 22-528, 22-559, 22-568, 24-493, 24-568, 27-369, 29-291, 32-250, 33-448, 39-340, 47-301, 49-369, 51-458, 51-563, 53-252, 64-680, 82-504, 85-568, 101-568, 120-390, 139-301, 160-466, 229-458, 251-472, 251-537, 254-568, 274-486, 274-491, 296-568, 302-761, 303-379, 303-844, 305-475, 314-564, 318-559, 321-544, 325-542, 359-568, 419-568, 566-798, 575-1270, 630-1325, 637-1271, 638-1133, 642-892, 645-1222, 645-1349, 646-887, 652-1026, 670-1392, 679-971, 701-1264, 705-1394, 709-1340, 713-969, 726-1325, 736-1308, 739-1002, 743-1376, 744-1386, 757-1037, 757-1042, 757-1065, 758-1394, 769-1023, 780-1332, 781-1027, 781-1281, 782-1063, 783-1440, 788-1638, 792-1061, 792-1081, 812-1040, 814-1044, 814-1214, 816-919, 817-1093, 822-1348, 825-1052, 828-1399, 831-1111, 846-1071, 849-1399, 860-1058, 860-1070, 860-1399, 866-1452, 866-1555, 874-1650, 875-1170, 875-1248, 875-1263, 875-1304, 875-1329, 875-1334, 875-1341, 875-1358, 875-1371, 875-1375, 875-1381, 875-1389, 875-1395, 875-1396, 875-1403, 875-1405, 875-1412, 875-1418, 875-1431, 875-1469, 875-1472, 875-1474, 875-1491, 875-1506, 875-1509, 875-1530, 875-1533, 875-1540, 875-1542, 875-1550, 875-1615, 875-1629, 875-1641, 875-1670, 882-1393, 886-1162, 886-1733, 893-1171, 893-1183, 894-1126, 896-1156, 900-1549, 904-1156, 904-1514, 907-1395, 921-1768, 923-1696.

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
56	927-1171, 931-1396, 941-1742, 945-1134, 952-1218, 953-1477, 959-1170, 959-1186, 969-1216, 976-1240, 989-1430, 992-1269, 995-1269, 1003-1343, 1010-1235, 1017-1727, 1023-1279, 1023-1300, 1023-1704, 1026-1304, 1027-1241, 1027-1325, 1027-1636, 1030-1744, 1040-1755, 1042-1787, 1043-1205, 1043-1308, 1045-1226, 1047-1671, 1064-1618, 1075-1748, 1076-1761, 1079-1351, 1084-1705, 1086-1719, 1092-1769, 1114-1393, 1121-1630, 1124-1706, 1124-1755, 1125-1799, 1127-1785, 1133-1395, 1134-1787, 1136-1758, 1138-1622, 1141-1765, 1141-1802, 1144-1477, 1145-1704, 1148-1786, 1156-1772, 1167-1333, 1168-1441, 1168-1701, 1168-1762, 1168-1786, 1176-1474, 1180-1423, 1186-1456, 1191-1798, 1200-1787, 1205-1515, 1212-1753, 1224-1761, 1224-1787, 1226-1785, 1234-1515, 1234-1787, 1236-1433, 1236-1752, 1236-1772, 1236-1785, 1237-1786, 1264-1500, 1264-1790, 1264-1793, 1275-1796, 1282-1604, 1282-1772, 1285-1795, 1286-1794, 1301-1796, 1302-1588, 1304-1772, 1308-1772, 1312-1766, 1315-1772, 1321-1772, 1322-1772, 1325-1796, 1327-1772, 1328-1766, 1328-1785, 1337-1772, 1337-1799, 1340-1780, 1342-1766, 1342-1772, 1343-1772, 1344-1772, 1345-1588, 1345-1772, 1348-1775, 1349-1781, 1356-1772, 1358-1770, 1359-1773, 1363-1772, 1366-1772, 1368-1802, 1369-1772, 1374-1772, 1375-1766, 1375-1772, 1379-1765, 1380-1802, 1386-1701, 1386-1714, 1393-1772, 1401-1785, 1401-1802, 1407-1772, 1408-1772, 1409-1703, 1415-1786, 1415-1787, 1423-1704, 1425-1772, 1433-1787, 1434-1782, 1437-1668, 1441-1692, 1448-1772, 1464-1772, 1466-1786, 1472-1757, 1477-1611, 1478-1789, 1479-1750, 1484-1759, 1485-1743, 1485-1772, 1491-1772, 1498-1748, 1498-1772, 1503-1772, 1512-1777, 1515-1786, 1530-1779, 1534-1772, 1553-1772, 1559-1772, 1568-1786, 1579-1785, 1599-1802, 1611-1802, 1612-1784, 1621-1802, 1629-1802, 1637-1767, 1637-1797, 1654-1775, 1673-1799, 1682-1767, 1711-1787 1-133, 1-147, 1-151, 1-1833, 5-120, 13-120, 13-134, 13-147, 89-761, 90-308, 90-309, 90-469, 90-508, 90-545, 90-682, 90-767, 154-383, 154-682, 155-769, 170-952, 176-397, 176-462, 179-559, 199-411, 199-416, 210-846, 214-834, 219-742, 219-809, 221-513, 227-621, 227-640, 227-700, 227-737, 227-762, 227-800, 227-808, 227-809, 227-821, 227-866, 227-868, 227-882, 227-907, 228-304, 228-710, 228-745, 228-752, 228-761, 228-762, 228-776, 228-779, 228-786, 228-812, 228-820, 228-853, 228-865, 228-866, 228-873, 228-876, 228-899, 228-902, 228-926, 229-805, 230-400, 230-898, 239-489, 243-484, 246-469, 249-562.
57/7500362CB1/ 1833	

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
57	250-467, 267-854, 268-966, 271-953, 274-751, 275-936, 284-510, 285-528, 285-560, 297-1056, 334-594, 334-933, 336-624, 336-901, 337-816, 343-1042, 344-754, 348-745, 348-854, 373-643, 386-1091, 390-607, 420-676, 428-678, 444-659, 463-724, 468-1108, 474-730, 477-712, 480-723, 484-761, 484-776, 484-812, 491-758, 493-747, 494-1121, 498-725, 498-738, 498-773, 501-1102, 512-802, 513-810, 519-764, 528-996, 530-1180, 537-1184, 548-921, 551- 765, 552-1073, 565-1212, 567-1203, 574-795, 576-856, 584-807, 590-1146, 594-835, 604-874, 611-870, 614-846, 614-848, 623-1318, 650-843, 678-1373, 685-1319, 686-1181, 690-940, 693-1270, 693-1397, 694-935, 698-986, 699-953, 700-976, 700-1074, 718-976, 718-1440, 727-1019, 749- 1312, 753-1442, 757-1388, 761-1017, 774-1373, 784-1356, 787-1050, 791-1424, 792-1434, 805-1085, 805-1090, 805-1113, 806-1442, 817-1071, 828-1380, 829-1075, 829-1329, 830-1111, 831-1488, 836-1682, 840-1109, 840- 1129, 860-1088, 862-1092, 862-1262, 864-967, 865-1141, 870-1396, 873-1100, 876-1447, 879-1159, 894-1119, 897- 1447, 908-1447, 914-1500, 914-1634, 922-1694, 923-1218, 923-1296, 923-1311, 923-1352, 923-1377, 923-1382, 923-1389, 923-1406, 923-1419, 923-1423, 923-1429, 923-1437, 923-1443, 923-1444, 923-1451, 923-1453, 923- 1460, 923-1466, 923-1479, 923-1517, 923-1520, 923-1522, 923-1539, 923-1554, 923-1557, 923-1578, 923-1581, 923-1588, 923-1590, 923-1598, 923-1659, 923-1673, 923-1685, 923-1714, 930-1441, 934-1210, 934-1777, 941- 1219, 941-1231, 942-1174, 944-1204, 948-1609, 952-1204, 952-1562, 955-1443, 969-1812, 971-1740, 975-1219, 979-1444, 989-1786, 993-1182, 1001-1525, 1007-1234, 1037-1478, 1040-1317, 1043-1317, 1051-1391, 1058-1283, 1065-1771, 1071-1327, 1071- 1348, 1071-1748, 1074-1352, 1075-1289, 1075-1373, 1075-1680, 1078-1788, 1088-1799, 1091-1253, 1091-1356, 1095-1715, 1112-1662, 1123-1792, 1124-1805, 1127-1399, 1132-1749, 1134-1763, 1140-1813, 1162-1441, 1169- 1674, 1172-1750, 1172-1799, 1173-1833, 1175-1829, 1181-1443, 1184-1802, 1186-1666, 1189-1809, 1189-1833, 1192-1525, 1193-1748, 1196-1830, 1204-1816, 1216-1489, 1216-1745, 1216-1806, 1224-1522, 1228-1471, 1234- 1504, 1239-1833, 1253-1563, 1260-1797, 1272-1805, 1274-1814, 1282-1563, 1282-1831, 1284-1481, 1284-1796, 1284-1829, 1285-1830, 1312-1548, 1323-1833, 1330-1648, 1330-1825, 1334-1833, 1352-1820, 1356-1818, 1360- 1810, 1372-1652, 1373-1833, 1376-1814, 1385-1833, 1406-1814, 1455-1822, 1612-1830, 1643-1833, 1655-1833, 1656-1828, 1665-1833, 1673-1833, 1681-1833, 1717-1833, 1726-1811, 1737-1819

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
58/7503328CB1/ 2465	1-262, 1-286, 1-308, 1-317, 1-325, 1-496, 1-501, 1-512, 1-621, 2-201, 4-296, 8-233, 8-246, 8-524, 8-535, 8-2463, 15- 699, 15-774, 15-783, 15-809, 15-854, 15-903, 16-771, 19-265, 22-187, 22-235, 27-457, 27-631, 38-731, 39-627, 44- 243, 44-592, 44-642, 49-645, 50-181, 50-325, 56-175, 56-543, 58-170, 58-329, 60-352, 62-302, 63-206, 64-335, 67- 330, 69-294, 69-360, 69-548, 69-598, 69-613, 71-630, 71-747, 73-560, 73-730, 76-545, 84-685, 88-392, 89-636, 90- 312, 90-681, 97-377, 98-662, 100-375, 100-736, 103-368, 106-369, 109-691, 109-764, 134-386, 139-616, 139-767, 147-731, 182-675, 182-692, 184-710, 184-719, 190-758, 190-836, 195-451, 195-720, 195-747, 208-611, 232-780, 282-531, 290-520, 342-798, 345-790, 351-914, 366-614, 371-675, 407-880, 407-984, 420-914, 439-883, 459-1000, 472-961, 473-810, 480-1017, 488-797, 491-1048, 510-1060, 524-1084, 549-1100, 574-1139, 578-856, 581-1139, 584-845, 586-789, 587-815, 593-1014, 593-1136, 593-1139, 593-1141, 600-850, 608-888, 608-1040, 615-1024, 615-1079, 615-1082, 617-1036, 636-884, 636-1106, 636-1153, 640-1270, 644-1278, 648-1192, 660-1105, 669-956, 669-1192, 669-1305, 677-956, 681-1325, 688-1010, 688-1075, 688-1279, 689-1265, 698-971, 698-1289, 705-1168, 705-1169, 705-1192, 709-1079, 709-1189, 719-1188, 726-1291, 730-1279, 734-1293, 747-1075, 748-1023, 764-1192, 764-1352, 765-1079, 765-1192, 767- 1278, 770-1244, 794-1352, 799-1268, 802-1304, 804-1344, 816-1329, 816-1344, 816-1352, 817-1079, 817-1141, 817-1153, 817-1236, 817-1279, 817-1280, 817-1350, 818-1236, 829-1111, 846-1062, 849-1222, 863-1289, 872-1093, 893-1114, 896-1352, 900- 1157, 926-1351, 930-1352, 939-1214, 965-1232, 1008-1351, 1023-1305, 1059-1233, 1075-1304, 1125-1352, 1125- 1418, 1125-1466, 1128-1410, 1178-1352, 1355-1904, 1360-2036, 1366-1862, 1376-1725, 1380-1645, 1384-1862, 1384-1883, 1404-1928, 1419-1639, 1426-1864, 1432-2008, 1433-1662, 1433-2036, 1437-1743, 1467-1752, 1467- 1782, 1467-1788, 1475-2009, 1479-1771, 1490-1904, 1493-1936, 1498-2090, 1508-1815, 1518-1788, 1528-1772, 1572-2194, 1574-2167, 1579-1828, 1588-1984, 1597-1842, 1599-2178, 1601-1977, 1601-2159, 1601-2175, 1601- 2189, 1610-1884, 1611-1815, 1616-1830, 1616-2202, 1631-1788, 1636-1860, 1650-2102, 1652-2180, 1659-2431, 1667-2405, 1668-1859, 1669-2211, 1681-2369, 1700-1990, 1719-1966, 1725-2445, 1727-2314, 1732-1989, 1732- 2325, 1735-2434, 1741-2328, 1753-2381, 1755-2404, 1763-2434, 1779-2075, 1779-2431, 1783-2404, 1783-2442,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
58	1791-2400, 1798-2019, 1798-2398, 1799-2412, 1799-2433, 1806-2222, 1808-2437, 1810-2465, 1811-2434, 1823-2092, 1825-2060, 1830-2465, 1832-2325, 1835-2061, 1835-2104, 1838-2073, 1839-2435, 1843-2130, 1845-2102, 1850-2098, 1852-2410, 1859-2325, 1869-2142, 1873-2451, 1878-2382, 1902-2443, 1903-2172, 1903-2438, 1903-2445, 1903-2459, 1904-2156, 1905-2201, 1906-2378, 1912-2145, 1915-2325, 1919-2266, 1922-2440, 1923-2219, 1924-2406, 1925-2442, 1929-2169, 1929-2465, 1941-2240, 1943-2465, 1944-2465, 1952-2130, 1952-2199, 1952-2451, 1956-2189, 1964-2160, 1964-2464, 1964-2465, 1967-2231, 1970-2455, 1971-2167, 1974-2437, 1974-2465, 1984-2401, 1984-2464, 1984-2465, 1985-2453, 1991-2349, 1992-2449, 1992-2451, 1994-2451, 1995-2242, 1996-2451, 1996-2465, 2002-2216, 2002-2453, 2004-2250, 2007-2228, 2007-2457, 2011-2453, 2011-2465, 2015-2401, 2015-2454, 2015-2457, 2016-2401, 2019-2401, 2021-2401, 2021-2450, 2022-2465, 2023-2465, 2024-2465, 2025-2281, 2025-2465, 2030-2465, 2031-2238, 2033-2450, 2036-2450, 2037-2401, 2038-2401, 2039-2401, 2041-2401, 2042-2387, 2043-2401, 2043-2451, 2043-2465, 2045-2401, 2047-2455, 2048-2291, 2049-2465, 2052-2401, 2053-2401, 2055-2401, 2056-2318, 2057-2401, 2058-2318, 2058-2319, 2059-2307, 2061-2465, 2062-2401, 2063-2394, 2068-2401, 2070-2465, 2079-2291, 2082-2312, 2082-2338, 2086-2451, 2092-2349, 2093-2368, 2097-2352, 2097-2449, 2097-2451, 2098-2346, 2100-2401, 2101-2450, 2101-2465, 2103-2449, 2103-2451, 2103-2465, 2108-2401, 2109-2452, 2118-2401, 2121-2401, 2128-2465, 2136-2356, 2136-2427, 2143-2386, 2151-2401, 2158-2465, 2160-2401, 2166-2434, 2167-2455, 2169-2401, 2178-2401, 2180-2454, 2181-2463, 2188-2401, 2205-2465, 2214-2451, 2216-2443, 2219-2401, 2219-2451, 2234-2465, 2238-2451, 2239-2465, 2242-2376, 2243-2452, 2253-2464, 2272-2401, 2279-2401, 2286-2454, 2308-2451, 2323-2401, 2349-2465, 2396-2465
59/7510464CBI/ 2560	1-308, 1-325, 1-374, 1-496, 1-501, 1-512, 2-201, 4-296, 6-249, 6-317, 8-233, 8-246, 8-480, 8-535, 8-2555, 15-699, 15-760, 15-783, 15-809, 16-771, 19-265, 22-187, 22-235, 27-457, 30-165, 38-731, 43-286, 44-243, 49-645, 50-181, 50-325, 50-729, 56-175, 56-543, 56-674, 57-639, 57-641, 57-772, 57-774, 58-170, 58-329, 60-352, 62-302, 67-330, 69-294, 69-360, 69-391, 69-548, 69-613, 70-335, 71-630, 71-747, 73-455, 75-545, 77-917, 80-557, 80-614, 84-685, 88-165, 88-392, 89-422, 89-730, 90-312, 90-642, 90-681, 96-631, 97-200, 100-375, 100-736, 101-704, 103-368, 106-369, 109-691, 109-764, 110-304, 126-704, 134-386, 139-616, 139-767, 147-731, 173-781, 182-675, 182-692, 184-710, 184-719, 187-373, 188-542, 190-758, 190-836, 195-451, 195-677, 195-747, 208-611, 212-921, 232-780, 261-797, 281-911, 281-934, 282-531, 285-662, 290-520, 303-852, 307-816, 329-583, 342-798, 345-790, 347-816, 347-861, 366-614, 371-675, 371-921, 372-898, 379-584, 402-932, 404-882, 407-969, 407-984, 443-833, 447-559, 452-838, 464-1045,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
59	581-1139, 593-1139, 600-850, 615-1024, 640-1270, 681-1325, 688-1087, 698-1289, 709-1189, 719-1188, 726-1291, 734-1293, 764-1192, 765-1192, 770-1244, 799-1268, 804-1344, 817-1083, 817-1153, 817-1236, 817-1279, 817- 1280, 829-1111, 840-1379, 840-1455, 846-1062, 846-1369, 857-1404, 859-1406, 862-1507, 863-1289, 872-1093, 872-1415, 883-1481, 896-1352, 903-1507, 923-1619, 930-1360, 931-1374, 948-1507, 958-1506, 965-1232, 970- 1458, 987-1484, 993-1535, 1005-1490, 1008-1351, 1019-1563, 1023-1305, 1030-1507, 1047-1442, 1054-1488, 1071- 1609, 1076-1507, 1080-1473, 1080-1507, 1080-1552, 1083-1527, 1091-1376, 1091-1445, 1101-1617, 1105-1555, 1109-1540, 1111-1540, 1119-1650, 1125-1393, 1153-1552, 1154-1507, 1162-1417, 1168-1640, 1187-1467, 1197- 1746, 1200-1676, 1205-1852, 1212-1468, 1212-1517, 1214-1564, 1216-1484, 1222-1507, 1224-1875, 1227-1565, 1227-1657, 1235-1664, 1237-1575, 1237-1634, 1239-1446, 1239-2027, 1243-1520, 1245-1800, 1253-1590, 1268-1531, 1274-1609, 1278- 1562, 1280-1731, 1288-1856, 1295-1855, 1305-1778, 1321-1718, 1321-1791, 1327-1882, 1338-1834, 1348-1799, 1349-1727, 1360-1452, 1365-1720, 1384-2025, 1389-1950, 1396-2128, 1400-1896, 1406-1609, 1406-1692, 1416- 1970, 1418-1896, 1418-1917, 1423-1581, 1436-1854, 1437-2119, 1460-1898, 1471-1777, 1489-2004, 1501-1726, 1501-1778, 1501-1822, 1512-1757, 1513-1805, 1522-2121, 1524-1959, 1534-2108, 1542-1849, 1544-1780, 1560- 2128, 1562-1806, 1606-1918, 1619-2239, 1631-1876, 1644-1918, 1650-1864, 1665-1822, 1702-1893, 1803-2281, 1830-2444, 1860-2132, 1902-2504, 1965-2551, 1992-2508, 1994-2264, 1995-2535, 1999-2470, 2004-2237, 2007- 2417, 2015-2311, 2016-2498, 2021-2261, 2021-2560, 2025-2504, 2035-2558, 2036-2560, 2044-2475, 2048-2281, 2056-2557, 2059-2323, 2062-2547, 2066-2503, 2066-2559, 2076-2493, 2076-2557, 2076-2558, 2077-2545, 2078-2502, 2081-2500, 2084-2551, 2087-2334, 2088-2560, 2091-2509, 2095-2308, 2096-2342, 2099-2320, 2099- 2549, 2103-2545, 2104-2546, 2106-2549, 2107-2493, 2108-2493, 2111-2493, 2113-2542, 2114-2559, 2115-2557, 2117-2373, 2117-2560, 2122-2560, 2123-2327, 2125-2542, 2128-2542, 2129-2493, 2130-2493, 2131-2493, 2133- 2493, 2134-2479, 2135-2493, 2135-2504, 2135-2543, 2135-2557, 2137-2493, 2139-2547, 2140-2383, 2141-2557, 2143-2493, 2144-2493, 2145-2493, 2147-2493, 2149-2493, 2151-2399, 2153-2558, 2155-2486, 2160-2493, 2162- 2560, 2171-2383, 2174-2404, 2174-2430, 2178-2543, 2184-2441, 2185-2460, 2189-2444, 2189-2551, 2190-2438, 2192-2493, 2193-2560, 2195-2543, 2195-2560, 2201-2544, 2210-2493, 2213-2493, 2228-2448, 2228-2519, 2235- 2478, 2243-2493, 2250-2560, 2252-2493, 2258-2526, 2259-2547, 2261-2493, 2270-2493, 2272-2546, 2273-2555, 2280-2493, 2306-2543, 2311-2493, 2311-2543, 2326-2557, 2330-2543, 2331-2560, 2334-2468, 2335-2544, 2345- 2560, 2364-2493, 2367-2560, 2371-2493, 2378-2546, 2415-2493, 2488-2557



Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
60/7510394CB1/ 2254	1-351, 4-299, 4-2254, 5-266, 5-292, 5-311, 6-639, 9-261, 9-274, 9-276, 10-263, 10-277, 10-305, 12-232, 12-257, 13- 245, 13-284, 13-294, 15-296, 24-274, 32-263, 50-289, 52-553, 83-218, 108-281, 450-714, 452-1035, 452- 1108, 455-984, 455-993, 455-1039, 457-1294, 467-963, 474-1083, 482-1086, 483-896, 487-994, 500-693, 500-796, 500-1000, 500-1088, 502-737, 502-1062, 512-777, 517-992, 517-1102, 519-896, 519-1068, 520-650, 524-1118, 529- 1094, 543-841, 554-845, 556-1050, 564-797, 565-1423, 574-1164, 575-1051, 578-1086, 584-1190, 588-1161, 590- 836, 597-1118, 607-861, 610-1202, 612-1169, 613-1207, 618-1156, 635-1255, 642-1333, 648-1304, 650-881, 653- 895, 655-915, 656-955, 660-931, 660-1272, 667-1262, 670-1162, 673-1040, 674-1305, 686-1093, 698-968, 705- 1248, 724-970, 728-1091, 728-1122, 729-1224, 731-1000, 731-1278, 736-1091, 742-1341, 748-1321, 754-1018, 754- 1029, 757-1005, 759-1248, 761-1344, 763-1507, 765-1057, 770-1182, 771-1004, 771-1029, 771-1041, 774-1054, 774-1059, 776-1272, 776-1484, 779-1364, 781-902, 784-1040, 787-1333, 796-1023, 796-1091, 802-1044, 802-1392, 822-1480, 828-1085, 830-1454, 840-1099, 841-1302, 841-1350, 841-1374, 841-1582, 841-1586, 841-1607, 841-1692, 844-1551, 847-1108, 848-1028, 849-1099, 851-1105, 858-1326, 863-1431, 866-1463, 875-1123, 878- 1166, 880-1479, 882-1412, 882-1455, 885-1547, 886-1348, 898-1407, 908-1047, 911-1556, 912-1479, 921-1450, 931-1468, 940-1227, 950-1183, 952-1375, 953-1558, 954-1203, 954-1616, 960-1264, 962-1660, 963-1645, 970- 1582, 972-1593, 977-1213, 977-1220, 981-1602, 982-1276, 992-1500, 995-1276, 1000-1339, 1000-1634, 1001- 1284, 1003-1248, 1009-1617, 1014-1259, 1018-1144, 1018-1206, 1028-1734, 1031-1311, 1033-1273, 1038-1699, 1039-1614, 1041-1309, 1042-1580, 1045-1384, 1050-1322, 1054-1323, 1054-1331, 1059-1291, 1060-1553, 1062- 1314, 1064-1277, 1064-1337, 1064-1341, 1064-1352, 1067-1297, 1072-1324, 1072-1651, 1072-1668, 1082-1359, 1082-1377, 1097-1752, 1099-1531, 1102-1720, 1104-1680, 1108-1367, 1109-1361, 1109-1363, 1109-1407, 1112- 1424, 1115-1315, 1115-1330, 1123-1754, 1124-1371, 1125-1372, 1131-1362, 1133-1367, 1133-1705, 1134-1721, 1135-1312.

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
60	1135-1421, 1159-1770, 1162-1465, 1172-1479, 1174-1440, 1177-1366, 1179-1402, 1182-1511, 1182-1718, 1187- 1440, 1187-1469, 1187-1482, 1189-1461, 1189-1810, 1191-1840, 1192-1282, 1192-1505, 1203-1733, 1208-1794, 1211-1912, 1212-1484, 1220-1477, 1220-1717, 1220-1734, 1221-1488, 1222-1653, 1226-1730, 1229-1354, 1231- 1515, 1231-1869, 1232-1819, 1237-1804, 1237-1881, 1241-1777, 1250-1484, 1251-1504, 1265-1443, 1275-1519, 1278-1918, 1284-1776, 1290-1547, 1295-1544, 1297-1580, 1301-1503, 1302-1993, 1304-1442, 1314-1578, 1314- 1776, 1315-2007, 1321-1567, 1328-1596, 1331-1847, 1332-1901, 1341-1631, 1342-1584, 1342-1961, 1344-2131, 1348-1979, 1351-1900, 1354-1566, 1355-1602, 1357-1743, 1358-1643, 1360-1647, 1361-1576, 1373-2237, 1384- 1666, 1386-1585, 1390-1622, 1390-2055, 1392-1624, 1392-1791, 1396-1990, 1397-2036, 1404-1796, 1407-1810, 1409-1670, 1409-1697, 1409-1904, 1410-1643, 1417-1995, 1428-1626, 1429-1705, 1431-1677, 1432-1738, 1434-1963, 1434-2058, 1437-1709, 1437-1869, 1444-1705, 1444-1707, 1449-1671, 1458-1715, 1462-1708, 1469- 2018, 1473-1729, 1476-2190, 1477-1534, 1481-2201, 1486-1727, 1491-1719, 1491-1823, 1492-2244, 1494-1825, 1497-1750, 1499-2052, 1501-1758, 1506-1759, 1506-2111, 1507-2093, 1511-1792, 1511-1805, 1511-2174, 1512- 1773, 1513-1807, 1513-2132, 1515-2052, 1527-1765, 1527-1794, 1527-2084, 1531-2125, 1533-1731, 1536-1993, 1542-2110, 1546-1796, 1548-2192, 1550-1860, 1558-2200, 1567-2105, 1570-2147, 1571-2052, 1571-2166, 1574- 1876, 1576-2058, 1578-1836, 1580-1826, 1587-2059, 1588-2076, 1589-1989, 1589-2254, 1590-2200, 1590-2238, 1597-1835, 1598-1865, 1604-1854, 1604-1892, 1604-2008, 1604-2052, 1606-1861, 1607-1847, 1607-1859, 1609- 2195, 1610-1833, 1611-2238, 1613-1875, 1613-1876, 1613-1881, 1616-2237, 1617-1994, 1620-1715, 1624-2079, 1627-2122, 1628-1913, 1628-2239, 1630-1902, 1632-2210, 1633-2227, 1634-1914, 1637-2252, 1638-1909, 1642-2236, 1644-2249, 1646-1908, 1654-2252, 1655-1949, 1658-2238, 1665-1950, 1665-1966, 1665-2254, 1666- 1902, 1668-1936, 1675-1892, 1675-1912, 1675-2072, 1677-1874, 1677-1932, 1677-1953, 1677-2238, 1677-2251, 1678-2199, 1682-1888, 1684-2254, 1689-1940, 1689-2047, 1696-2200, 1699-2239, 1702-2254, 1706-1952, 1707- 2254, 1712-2116, 1713-1859, 1723-1972, 1725-2239, 1727-1970, 1736-2055, 1736-2220, 1737-1954, 1738-1961, 1738-2201, 1738-2224, 1738-2227, 1742-2046, 1743-1997, 1746-1997, 1748-2253, 1752-2209, 1764-2034, 1770- 2052, 1771-2210, 1771-2239, 1774-2246, 1775-2040, 1777-2254, 1781-2240, 1783-2025, 1785-2239, 1788-2254, 1791-2199, 1791-2254, 1793-2245, 1798-2242, 1800-1895, 1800-2204, 1802-2065, 1803-2237, 1805-2201, 1810- 2205, 1811-2201, 1813-2247, 1815-2242, 1821-2239, 1822-2196, 1822-2238, 1822-2242, 1825-2238, 1825-2244, 1826-2085, 1827-2111, 1828-2238, 1830-2254, 1831-2238, 1831-2254, 1832-2169, 1832-2239, 1832-2251,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
60	1832-2252, 1835-2238, 1837-2247, 1839-2245, 1841-2244, 1841-2247, 1842-2238, 1843-2142, 1844-2211, 1845-2123, 1845-2238, 1847-2216, 1848-2238, 1854-2248, 1857-2027, 1860-2254, 1861-2145, 1870-2085, 1870-2238, 1874-2239, 1875-2106, 1875-2254, 1877-2239, 1881-2254, 1884-2238, 1891-1913, 1891-2210, 1895-2238, 1899-2100, 1899-2242, 1904-2124, 1904-2239, 1913-2238, 1915-2194, 1915-2238, 1917-2201, 1917-2239, 1918-2238, 1920-2239, 1921-2238, 1922-2240, 1926-2126, 1926-2185, 1933-2222, 1934-2149, 1934-2159, 1934-2179, 1934-2242, 1935-2172, 1936-2213, 1936-2220, 1936-2239, 1938-2219, 1939-2219, 1947-2238, 1949-2244, 1971-2238, 1983-2182, 1985-2249, 1991-2244, 1993-2242, 1994-2238, 2037-2254, 2038-2254, 2043-2254, 2057-2254, 2071-2238, 2078-2254, 2081-2239, 2088-2254, 2100-2254, 2102-2238, 2123-2238, 2124-2252, 2125-2239, 2127-2254, 2142-2201, 2142-2237, 2142-2238, 2145-2254, 2149-2253, 2154-2238
61/7500745CB1/ 2139	1-125, 1-249, 1-290, 1-294, 1-334, 2-233, 4-334, 7-250, 7-255, 7-259, 7-271, 7-279, 7-302, 7-2139, 8-269, 8-290, 8-295, 8-314, 8-324, 10-334, 12-264, 12-274, 12-277, 12-279, 13-266, 13-275, 13-278, 13-280, 13-281, 13-308, 15-235, 15-260, 15-270, 16-248, 16-287, 16-297, 16-314, 18-299, 27-277, 35-266, 53-292, 58-329, 71-254, 86-221, 86-314, 86-334, 111-284, 264-867, 333-597, 335-756, 335-797, 335-820, 335-844, 335-848, 335-918, 337-991, 338-867, 338-876, 338-922, 350-846, 356-816, 357-900, 357-966, 358-510, 358-616, 362-885, 365-934, 365-969, 366-779, 370-877, 377-759, 383-576, 383-620, 383-679, 383-940, 383-971, 385-620, 385-945, 395-660, 400-875, 400-985, 402-779, 402-951, 403-533, 407-1001, 412-977, 426-724, 437-728, 439-933, 447-680, 447-688, 454-996, 455-1066, 458-934, 461-969, 466-898, 467-1073, 471-1044, 473-719, 490-744, 493-1085, 495-1052, 496-1090, 501-1039, 507-1106, 508-1036, 511-1102, 514-783, 518-1138, 525-1216, 531-1187, 533-764, 536-778, 538-683, 538-798, 539-838, 543-814, 550-1145, 553-1045, 556-923, 569-976, 581-851, 588-1131, 607-853, 611-974, 611-1005, 612-1107, 614-883, 614-1161, 619-974, 625-1224, 631-1204, 637-901, 637-912, 640-888, 642-1131, 644-1227, 646-1390, 648-940, 653-1065, 654-887, 654-912, 654-924, 657-942, 657-1056, 658-1482, 659-861, 662-1247, 664-785, 667-923, 670-1216, 679-906, 679-974, 685-927, 685-1275, 705-1363, 711-873, 711-968, 713-1337, 724-1185, 724-1233, 724-1257, 724-1465, 724-1469, 724-1490, 724-1546, 724-1575, 727-1434, 730-991, 731-911, 732-982, 734-988, 741-1209, 746-1314, 758-1006, 761-1049, 763-1362, 765-1295, 765-1338, 768-1430, 769-1231, 781-1290, 791-930, 794-1439, 795-1362, 804-1333, 814-1351, 816-1044,

Table 4

Polynucleotide SEQ ID NO./ Incye ID/ Sequence Length	Sequence Fragments
61	821-1083, 823-1110, 833-1066, 836-1441, 837-1086, 837-1499, 843-1147, 845-1543, 846-1528, 855-1476, 860-1096, 860-1103, 864-1485, 872-1159, 875-1383, 878-1159, 883-1517, 884-1167, 886-1131, 892-1500, 897-1142, 901-1027, 901-1089, 911-1617, 916-1156, 922-1497, 924-1192, 926-1171, 928-1267, 933-1205, 937-1206, 937-1214, 942-1174, 945-1197, 947-1220, 947-1224, 947-1235, 947-1480, 950-1180, 955-1207, 955-1534, 955-1551, 965-1260, 969-1130, 969-1215, 972-1180, 980-1635, 982-1250, 982-1414, 985-1603, 987-1563, 991-1250, 992-1244, 992-1246, 992-1290, 995-1307, 998-1198, 998-1213, 998-1260, 1000-1159, 1006-1637, 1008-1255, 1016-1250, 1016-1588, 1017-1604, 1018-1195, 1018-1304, 1055-1362, 1057-1323, 1060-1249, 1062-1285, 1070-1323, 1070-1365, 1072-1693, 1074-1723, 1094-1795, 1105-1536, 1109-1613, 1109-1618, 1112-1237, 1114-1752, 1115-1702, 1120-1687, 1120-1764, 1124-1660, 1133-1367, 1134-1387, 1158-1402, 1167-1659, 1178-1427, 1180-1463, 1185-1876, 1194-1844, 1197-1461, 1197-1659, 1198-1890, 1204-1450, 1211-1479, 1214-1730, 1215-1784, 1225-1467, 1225-1910, 1237-1449, 1238-1485, 1243-1530, 1256-2125, 1265-2070, 1275-1507, 1275-1674, 1279-1873, 1292-1553, 1292-1787, 1293-1526, 1312-1588, 1314-1560, 1315-1621, 1317-1846, 1317-1941, 1327-1588, 1327-1590, 1332-1554, 1352-1901, 1359-2073, 1360-1417, 1364-2084, 1364-2098, 1369-1610, 1374-1602, 1375-2127, 1377-1708, 1380-1633, 1382-1935, 1384-1641, 1389-1642, 1389-1994, 1390-1976, 1394-1675, 1394-1688, 1394-2057, 1395-1656, 1396-1690, 1396-2015, 1409-1648, 1410-1643, 1410-1648, 1410-1677, 1410-1967, 1418-2092, 1419-1876, 1425-1993, 1431-2075, 1435-2066, 1436-2063, 1441-2083, 1444-2078, 1450-1988, 1452-2109, 1454-2049, 1454-2081, 1459-1941, 1461-1719, 1467-2033, 1470-2107, 1472-2137, 1473-2083, 1473-2129, 1480-1718, 1487-1737, 1487-1775, 1487-1891, 1489-1744, 1490-1742, 1492-2078, 1494-2124, 1496-1764, 1499-2120, 1500-1877, 1503-1598, 1507-1962, 1511-2112, 1511-2121, 1513-1785, 1515-2093, 1516-2110, 1520-2135, 1525-2119, 1527-2132, 1534-2050, 1537-2135, 1538-1832, 1541-2121, 1548-1849, 1548-2137, 1549-1785, 1551-1819, 1558-1775, 1558-1795, 1558-1858, 1558-1955, 1560-1704, 1560-1757, 1560-1815, 1560-2121, 1560-2134, 1567-2137, 1572-1823, 1572-2006, 1576-2107, 1582-2121, 1585-2137, 1589-1835, 1590-2137, 1595-1999, 1596-1742, 1606-1855, 1608-2121, 1610-1853, 1613-1894, 1613-2137, 1615-2081, 1619-1938, 1619-2103, 1620-1837, 1621-1844, 1621-2106, 1621-2136, 1625-1929, 1626-1880, 1629-1880, 1631-2136, 1635-2092, 1647-1917, 1653-1935, 1654-2093, 1654-2121, 1657-2119, 1658-1923, 1660-2137, 1664-2121, 1666-1908, 1668-2121, 1671-2137, 1674-2137, 1676-2120, 1681-2125, 1683-1778, 1683-2087, 1685-1948, 1686-2120, 1693-2088, 1694-2084, 1696-2130, 1698-2125, 1704-2121, 1705-2079, 1705-2123, 1705-2125, 1708-2122, 1708-2127, 1709-1968, 1710-1994, 1711-2121, 1713-2137,

Table 4

Polynucleotide SEQ ID NO./ Incyte ID/ Sequence Length	Sequence Fragments
61	1714-2121, 1714-2137, 1715-2052, 1715-2121, 1715-2135, 1715-2139, 1718-2123, 1720-2130, 1722-2128, 1724-2119, 1724-2127, 1725-2121, 1725-2122, 1726-2025, 1727-2094, 1728-2006, 1728-2121, 1730-2099, 1731-2121, 1737-2121, 1740-1910, 1743-2137, 1744-2028, 1753-1968, 1753-2121, 1757-2121, 1758-1989, 1758-2137, 1760-2121, 1764-2137, 1767-2121, 1774-1796, 1774-2093, 1778-2121, 1782-1983, 1782-2125, 1787-2007, 1787-2121, 1796-2121, 1798-2077, 1798-2121, 1800-2084, 1800-2121, 1803-2121, 1804-2121, 1805-2121, 1809-2009, 1809-2068, 1816-2105, 1817-2032, 1817-2042, 1817-2062, 1817-2125, 1818-2055, 1819-2103, 1819-2116, 1819-2133, 1821-2102, 1822-2102, 1830-2124, 1832-2127, 1866-2065, 1868-2132, 1874-2127, 1876-2125, 1877-2121, 1920-2137, 1921-2137, 1926-2137, 1940-2137, 1954-2121, 1961-2137, 1971-2137, 1983-2137, 1985-2121, 2007-2135, 2008-2121, 2010-2137, 2025-2084, 2025-2120, 2025-2121, 2032-2136, 2037-2124
62/7500929CB1/ 648	1-164, 1-250, 1-259, 1-269, 1-273, 1-354, 1-356, 1-469, 15-597, 43-329, 341-616, 343-598, 343-617, 343-622, 343-648, 345-594, 347-596, 384-606, 448-613, 451-610, 475-596, 511-601

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID:	Representative Library
32	8268274CB1	TESTTUT02
33	7500515CB1	LIVRTMR01
34	2256826CB1	UTRSNON03
35	7686186CB1	BRABDIK02
37	7501945CB1	UTRSTUE01
38	7500264CB1	LIVRTUT01
39	7499935CB1	BRABDIK02
40	7982285CB1	MIXDTME01
41	7758505CB1	HEL AUNT01
42	6885756CB1	BRAWTDR02
43	7500748CB1	SCORNON02
44	7500749CB1	BRSTTUT03
45	7503401CB1	PLACFEF05
46	7503485CB1	BRSTNOT01
47	7504076CB1	SPLNTUE01
48	7500926CB1	HELATXT01
49	7503216CB1	FIBRTXS07
50	7503233CB1	PROSTUT09
51	7726576CB1	HIPONON02
52	7503507CB1	KIDEUNE02
53	7503506CB1	KIDEUNE02
54	7503509CB1	LIVRTUT01
55	7503800CB1	MUSCNOT02
56	7503141CB1	LIVRTMR01
57	7500362CB1	COLNPOT01
58	7503328CB1	UTRSNOT11
59	7510464CB1	BONMTUE02
60	7510394CB1	BRSTTUT03
61	7500745CB1	BRSTTUT03
62	7500929CB1	LUNGNOT09

PAGE MISSING AT THE TIME OF THE  
PUBLICATION

Table 6

Library	Vector	Library Description
BRSTTUT03	PSPORT1	Library was constructed using RNA isolated from breast tumor tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated multicentric invasive grade 4 lobular carcinoma. The mass was identified in the upper outer quadrant, and three separate nodules were found in the lower outer quadrant of the left breast. Patient history included skin cancer, rheumatic heart disease, osteoarthritis, and tuberculosis. Family history included cerebrovascular disease, coronary artery aneurysm, breast cancer, prostate cancer, atherosclerotic coronary artery disease, and type I diabetes.
COLNPOT01	pINCY	Library was constructed using RNA isolated from colon polyp tissue removed from a 40-year-old Caucasian female during a total colectomy. Pathology indicated an inflammatory pseudopolyp; this tissue was associated with a focally invasive grade 2 adenocarcinoma and multiple tubovillous adenomas. Patient history included a benign neoplasm of the bowel.
FIBRTXS07	pINCY	This subtracted library was constructed using 1.3 million clones from a dermal fibroblast library and was subjected to two rounds of subtraction hybridization with 2.8 million clones from an untreated dermal fibroblast tissue library. The starting library for subtraction was constructed using RNA isolated from treated dermal fibroblast tissue removed from the breast of a 31-year-old Caucasian female. The cells were treated with 9CIS retinoic acid. The hybridization probe for subtraction was derived from a similarly constructed library from RNA isolated from untreated dermal fibroblast tissue from the same donor. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR (1991) 19:1954 and Bonaldo, et al., Genome Research (1996) 6:791.
HELATXT01	pINCY	Library was constructed using RNA isolated from HeLa cells treated with TNF- $\alpha$ and IL-1 $\beta$ , 10ng/ml each for 20 hours. The HeLa cell line is derived from cervical adenocarcinoma removed from a 31-year-old Black female.
HELAUNT01	pINCY	Library was constructed using RNA isolated from HeLa cells. The HeLa cell line is derived from cervical adenocarcinoma removed from a 31-year-old Black female.
HIPONON02	PSPORT1	This normalized hippocampus library was constructed from 1.13M independent clones from a hippocampus tissue library. RNA was isolated from the hippocampus tissue of a 72-year-old Caucasian female who died from an intracranial bleed. Patient history included nose cancer, hypertension, and arthritis. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228).
KIDEUNE02	pINCY	This 5' biased random primed library was constructed using RNA isolated from an untreated transformed embryonal cell line (293-EBNA) derived from kidney epithelial tissue (Invitrogen). The cells were transformed with adenovirus 5 DNA.



PAGE MISSING AT THE TIME OF THE  
PUBLICATION

Table 6

Library	Vector	Library Description
PLACFEF05	PCMV-ICIS	Library was constructed using RNA isolated from placental tissue removed from a Caucasian fetus, who died after 16 weeks' gestation from fetal demise and hydrocephalus. Patient history included umbilical cord wrapped around the head (3 times) and the shoulders (1 time). Serology was positive for anti-CMV and remaining serologies were negative. Family history included multiple pregnancies and live births, and an abortion in the mother.
PROSTUT09	pINCY	Library was constructed using RNA isolated from prostate tumor tissue removed from a 66-year-old Caucasian male during a radical prostatectomy, radical cystectomy, and urinary diversion. Pathology indicated grade 3 transitional cell carcinoma. The patient presented with prostatic inflammatory disease. Patient history included lung neoplasm, and benign hypertension. Family history included a malignant breast neoplasm, tuberculosis, cerebrovascular disease, atherosclerotic coronary artery disease and lung cancer.
SCORNON02	PSPORT1	This normalized spinal cord library was constructed from 3.24M independent clones from the a spinal cord tissue library. RNA was isolated from the spinal cord tissue removed from a 71-year-old Caucasian male who died from respiratory arrest. Patient history included myocardial infarction, gangrene, and end stage renal disease. The normalization and hybridization conditions were adapted from Soares et al.(PNAS (1994) 91:9228).
SPLNTUE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from spleen tumor tissue removed from a 28-year-old male during total splenectomy. Pathology indicated malignant lymphoma, diffuse large cell type, B-cell phenotype with abundant reactive T-cells and marked granulomatous response involving the spleen, where it formed approximately 45 nodules, liver, and multiple lymph nodes.
TESTTUT02	pINCY	Library was constructed using RNA isolated from testicular tumor removed from a 31-year-old Caucasian male during unilateral orchiectomy. Pathology indicated embryonal carcinoma.
UTRSNON03	pINCY	This normalized library was constructed from 6.4M independent clones from the UTRSNOT12 library. RNA was isolated from uterine myometrial tissue removed from a 41-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Pathology for the associated tumor tissue indicated uterine leiomyoma. Patient history included ventral hernia and a benign ovarian neoplasm. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228).

Table 6

Library	Vector	Library Description
UTRSNOT11	pINCY	Library was constructed using RNA isolated from uterine myometrial tissue removed from a 43-year-old female during a vaginal hysterectomy and removal of the fallopian tubes and ovaries. Pathology for the associated tumor tissue indicated that the myometrium contained an intramural and a submucosal leiomyoma. Family history included benign hypertension, hyperlipidemia, colon cancer, type II diabetes, and atherosclerotic coronary artery disease.
UTRSTUE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from uterus tumor tissue removed a 37-year-old Black female during myomectomy, dilation and curettage, right fimbrial region biopsy, and incidental appendectomy. Pathology indicated multiple (12) uterine leiomyomata. A fimbrial cyst was identified. The patient presented with deficiency anemia, an umbilical hernia, and premenopausal menorrhagia. Patient history included premenopausal menorrhagia and sarcoidosis of the lung. Previous surgeries included hysterectomy, dilation and curettage, and an endoscopic lung biopsy. Patient medications included Chromagen and Claritin. Family history included acute myocardial infarction and atherosclerotic coronary artery disease in the father.

PAGE MISSING AT THE TIME OF THE  
PUBLICATION

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score=120 or greater; Match length=56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
57	7500362	1239004H1	SNP00100193	124	603	T	T	C	Y198	n/a	n/a	n/a	n/a
57	7500362	1320137H1	SNP00061186	182	1013	G	G	C	G335	n/d	n/d	n/a	n/d
57	7500362	1402571H1	SNP00061186	251	1012	G	G	C	A335	n/d	n/d	n/a	n/d
57	7500362	1406223H1	SNP00051751	56	1340	C	C	A	T444	n/d	n/d	n/d	n/d
57	7500362	1493274H1	SNP00051750	17	1074	C	C	T	H355	n/a	n/a	n/a	n/a
57	7500362	1532731H1	SNP00061186	5	1011	G	G	C	Q334	n/d	n/d	n/a	n/d
57	7500362	1819683H1	SNP00051750	212	1076	C	C	T	A356	n/a	n/a	n/a	n/a
57	7500362	2563605H2	SNP00006074	54	1246	T	T	C	F413	n/a	n/a	n/a	n/a
57	7500362	3095949H1	SNP00051750	268	1073	C	C	T	P355	n/a	n/a	n/a	n/a
57	7500362	3095949H1	SNP00061186	205	1010	G	G	C	R334	n/d	n/d	n/a	n/d
57	7500362	3097881H1	SNP00051750	266	1072	C	C	T	H355	n/a	n/a	n/a	n/a
57	7500362	3097881H1	SNP00061186	203	1009	G	G	C	E334	n/d	n/d	n/a	n/d
57	7500362	3356734H1	SNP00100193	104	601	T	T	C	Y198	n/a	n/a	n/a	n/a
57	7500362	3481734H1	SNP00100193	17	597	T	T	C	N196	n/a	n/a	n/a	n/a
57	7500362	3773311H1	SNP00100193	92	602	T	T	C	F198	n/a	n/a	n/a	n/a
57	7500362	4071254H1	SNP00051751	39	1339	C	C	A	P444	n/d	n/d	n/d	n/d
57	7500362	4436184H1	SNP00100193	110	599	T	T	C	V197	n/a	n/a	n/a	n/a
57	7500362	4601138H1	SNP00006074	155	1245	C	C	A	F412	n/a	n/a	n/a	n/a
57	7500362	4833716H1	SNP00051750	228	1066	C	C	T	H353	n/a	n/a	n/a	n/a
57	7500362	4833716H1	SNP00061186	165	1001	G	G	C	G331	n/d	n/d	n/a	n/d
57	7500362	4839758H2	SNP00006074	170	1242	T	T	C	S411	n/a	n/a	n/a	n/a
57	7500362	4839758H2	SNP00051751	264	1336	C	C	A	L443	n/d	n/d	n/d	n/d
57	7500362	6809701J1	SNP00051750	123	1068	C	C	T	D353	n/a	n/a	n/a	n/a
57	7500362	6809701J1	SNP00061186	60	1005	G	G	C	M332	n/d	n/d	n/a	n/d
57	7500362	7191933H2	SNP00051750	225	1064	T	T	C	I352	n/a	n/a	n/a	n/a
57	7500362	7191933H2	SNP00061186	161	992	G	G	C	S328	n/d	n/d	n/a	n/d
58	7503328	055117H1	SNP00131006	104	2134	C	C	T	noncoding	n/a	n/a	n/a	n/a
58	7503328	1492725H1	SNP00025167	61	1794	G	G	A	noncoding	n/a	n/a	n/a	n/a

**PAGE MISSING AT THE TIME OF THE  
PUBLICATION**

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
59	7510464	3411052H1	SNP00131006	165	2222	C	C	T	noncoding	n/a	n/a	n/a	n/a
59	7510464	3512172H1	SNP00131006	181	2227	C	C	T	noncoding	n/a	n/a	n/a	n/a
59	7510464	3596460H1	SNP00025166	43	148	C	C	G	H28	n/d	n/d	n/d	n/d
59	7510464	3775141H1	SNP00025166	76	127	C	C	G	H21	n/d	n/d	n/d	n/d
59	7510464	4175359H1	SNP00131006	40	2224	C	C	T	noncoding	n/a	n/a	n/a	n/a
59	7510464	4465166H1	SNP00131006	41	2230	C	C	T	noncoding	n/a	n/a	n/a	n/a
59	7510464	4708867H1	SNP00068560	136	1378	T	T	C	Y438	n/a	n/a	n/a	n/a
59	7510464	4785219H1	SNP00141118	159	687	C	C	G	N207	n/a	n/a	n/a	n/a
59	7510464	4875944H1	SNP00068560	102	1380	T	T	C	H438	n/a	n/a	n/a	n/a
59	7510464	572892H1	SNP00131006	100	2239	C	C	T	noncoding	n/a	n/a	n/a	n/a
59	7510464	6550203H1	SNP00131006	188	2243	C	C	T	noncoding	n/a	n/a	n/a	n/a
60	7510394	1282428H1	SNP00075286	180	1054	C	C	T	noncoding	n/d	n/d	n/d	n/d
60	7510394	1282428H1	SNP00106459	90	964	C	C	G	noncoding	n/a	n/a	n/a	n/a
60	7510394	1299002H1	SNP00106460	204	1175	G	G	C	noncoding	n/a	n/a	n/a	n/a
60	7510394	1332279H1	SNP00009699	27	1457	G	A	G	noncoding	n/a	n/a	n/a	n/a
60	7510394	1332279H1	SNP00097916	45	1475	C	C	T	noncoding	n/a	n/a	n/a	n/a
60	7510394	1373856H1	SNP00009700	181	1706	T	C	T	noncoding	n/a	n/a	n/a	n/a
60	7510394	1401460H1	SNP00053363	87	95	G	G	A	R30	n/d	n/a	n/a	n/a
60	7510394	1483017H1	SNP00075287	63	1337	G	G	A	noncoding	n/a	n/a	n/a	n/a
60	7510394	1830710H1	SNP00009700	110	1707	T	C	T	noncoding	n/a	n/a	n/a	n/a
60	7510394	2113488H1	SNP00053363	86	94	G	G	A	V30	n/d	n/a	n/a	n/a
60	7510394	2113488H1	SNP00065694	65	73	C	C	T	P23	n/d	n/d	n/d	n/d
60	7510394	2481214H1	SNP00053363	81	93	G	G	A	R29	n/a	n/a	n/a	n/a
60	7510394	2481214H1	SNP00065694	59	71	C	C	T	P22	n/d	n/d	n/d	n/d
60	7510394	2492955H1	SNP00065694	74	74	C	C	T	A23	n/d	n/d	n/d	n/d
60	7510394	2530385H1	SNP00009700	5	1671	T	C	T	noncoding	n/a	n/a	n/a	n/a
60	7510394	2542562H1	SNP00009700	227	1712	C	C	T	noncoding	n/a	n/a	n/a	n/a
60	7510394	2560813H1	SNP00075287	163	1336	G	G	A	noncoding	n/a	n/a	n/a	n/a



**PAGE MISSING AT THE TIME OF THE  
PUBLICATION**

Table 8

SEQ ID NO:	PID	EST ID	SNP ID	EST SNP	CB1 SNP	EST Allele	Allele 1	Allele 2	Amino Acid	Caucasian Allele 1 frequency	African Allele 1 frequency	Asian Allele 1 frequency	Hispanic Allele 1 frequency
60	7510394	5074436H1	SNP00106459	136	963	C	C	G	noncoding	n/a	n/a	n/a	n/a
60	7510394	5594636H1	SNP00134605	86	856	A	A	C	noncoding	n/a	n/a	n/a	n/a
60	7510394	5879206H1	SNP00075286	37	971	C	C	T	noncoding	n/d	n/d	n/d	n/d
60	7510394	5879206H1	SNP00106460	158	1089	G	G	C	noncoding	n/d	n/a	n/a	n/a
60	7510394	5909982H1	SNP00053363	83	86	G	G	A	R27	n/d	n/a	n/a	n/a
60	7510394	5909982H1	SNP00065694	61	64	C	C	T	R20	n/d	n/d	n/d	n/d
60	7510394	5955251H1	SNP00075286	50	1049	T	C	T	noncoding	n/d	n/d	n/d	n/d
60	7510394	5955251H1	SNP00106460	171	1049	G	G	C	noncoding	n/d	n/a	n/a	n/a
60	7510394	5978154H1	SNP00075287	236	1334	G	G	A	noncoding	n/a	n/a	n/a	n/a
60	7510394	6171175H1	SNP00075287	212	1324	G	G	A	noncoding	n/d	n/a	n/a	n/a
60	7510394	6171175H1	SNP00106460	50	1162	G	G	C	noncoding	n/a	n/a	n/a	n/a
60	7510394	6849471H1	SNP00009699	177	1436	A	A	C	noncoding	n/d	n/a	n/a	n/a
60	7510394	6849471H1	SNP00097916	195	1454	C	C	G	noncoding	n/a	n/a	n/a	n/a
60	7510394	953943H1	SNP00106460	94	1176	G	G	T	noncoding	n/a	n/a	n/a	n/a
61	7500745	1282428H1	SNP00075286	180	937	C	C	C	noncoding	n/d	n/a	n/a	n/a
61	7500745	1282428H1	SNP00106459	90	847	C	C	T	noncoding	n/d	n/d	n/d	n/d
61	7500745	1299002H1	SNP00106460	204	1058	G	G	G	noncoding	n/a	n/a	n/a	n/a
61	7500745	1332279H1	SNP00009699	27	1340	G	G	C	noncoding	n/d	n/a	n/a	n/a
61	7500745	1332279H1	SNP00097916	45	1358	C	A	G	noncoding	n/a	n/a	n/a	n/a
61	7500745	1373856H1	SNP00009700	181	1589	T	C	T	noncoding	n/a	n/a	n/a	n/a
61	7500745	1401460H1	SNP00053363	87	99	G	C	T	L30	n/d	n/a	n/a	n/a
61	7500745	1483017H1	SNP00075287	63	1220	G	G	A	noncoding	n/a	n/a	n/a	n/a
61	7500745	2113488H1	SNP00065694	65	77	C	C	T	A23	n/d	n/d	n/d	n/d
61	7500745	5594636H1	SNP00134605	86	739	A	A	C	noncoding	n/a	n/a	n/a	n/a

**PAGE MISSING AT THE TIME OF THE  
PUBLICATION**

5. An isolated polynucleotide of claim 4 comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-62.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a  
5 polynucleotide of claim 3.

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

10

9. A method of producing a polypeptide of claim 1, the method comprising:

- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide  
15 encoding the polypeptide of claim 1, and
- b) recovering the polypeptide so expressed.

10. A method of claim 9, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-31.

20

11. An isolated antibody which specifically binds to a polypeptide of claim 1.

12. An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising a polynucleotide sequence selected from the group  
25 consisting of SEQ ID NO:32-62,
- b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:32-41, SEQ ID NO:43-56, and SEQ ID NO:61-62,
- c) a polynucleotide comprising a naturally occurring polynucleotide sequence at least  
30 92% identical to the polynucleotide sequence of SEQ ID NO:42,
- d) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 97% identical to the polynucleotide sequence of SEQ ID NO:59,

- e) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 98% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:58 and SEQ ID NO:60,
- f) a polynucleotide complementary to a polynucleotide of a),
- 5 g) a polynucleotide complementary to a polynucleotide of b),
- h) a polynucleotide complementary to a polynucleotide of c),
- i) a polynucleotide complementary to a polynucleotide of d),
- j) a polynucleotide complementary to a polynucleotide of e), and
- k) an RNA equivalent of a)-j).

10 13. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 12.

- 15 14. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:
- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target
  - 20 polynucleotide or fragments thereof, and
  - b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

25 15. A method of claim 14, wherein the probe comprises at least 60 contiguous nucleotides.

16. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:
- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
  - 30 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

17. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

18. A composition of claim 17, wherein the polypeptide comprises an amino acid sequence  
5 selected from the group consisting of SEQ ID NO:1-31.

19. A method for treating a disease or condition associated with decreased expression of functional PMMM, comprising administering to a patient in need of such treatment the composition of claim 17.

10

20. A method of screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

15

21. A composition comprising an agonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with decreased expression of  
20 functional PMMM, comprising administering to a patient in need of such treatment a composition of claim 21.

23. A method of screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- 25 a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

24. A composition comprising an antagonist compound identified by a method of claim 23 and a pharmaceutically acceptable excipient.

30

25. A method for treating a disease or condition associated with overexpression of functional PMMM, comprising administering to a patient in need of such treatment a composition of claim 24.

26. A method of screening for a compound that specifically binds to the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

27. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

28. A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

29. A method of assessing toxicity of a test compound, the method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 12 under conditions whereby a specific hybridization complex is formed between said probe and a target

polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 12 or fragment thereof,

- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

30. A method for a diagnostic test for a condition or disease associated with the expression of PMMM in a biological sample, the method comprising:

- a) combining the biological sample with an antibody of claim 11, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex, and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

31. The antibody of claim 11, wherein the antibody is:

- a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- d) a F(ab')<sub>2</sub> fragment, or
- e) a humanized antibody.

32. A composition comprising an antibody of claim 11 and an acceptable excipient.

33. A method of diagnosing a condition or disease associated with the expression of PMMM in a subject, comprising administering to said subject an effective amount of the composition of claim 32.

34. A composition of claim 32, wherein the antibody is labeled.



**PAGE MISSING AT THE TIME OF THE  
PUBLICATION**

41. A composition comprising the monoclonal antibody of claim 40 and a suitable carrier.

42. The antibody of claim 11, wherein the antibody is produced by screening a Fab expression library.

5

43. The antibody of claim 11, wherein the antibody is produced by screening a recombinant immunoglobulin library.

44. A method of detecting a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31 in a sample, the method comprising:

10   a)     incubating the antibody of claim 11 with the sample under conditions to allow specific binding of the antibody and the polypeptide, and

      b)     detecting specific binding, wherein specific binding indicates the presence of a polypeptide comprising an amino acid sequence selected from the group consisting of

15         SEQ ID NO:1-31 in the sample.

45. A method of purifying a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31 from a sample, the method comprising:

20   a)     incubating the antibody of claim 11 with the sample under conditions to allow specific binding of the antibody and the polypeptide, and

      b)     separating the antibody from the sample and obtaining the purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-31.

25   46. A microarray wherein at least one element of the microarray is a polynucleotide of claim 13.

47. A method of generating an expression profile of a sample which contains polynucleotides, the method comprising:

30   a)     labeling the polynucleotides of the sample,

      b)     contacting the elements of the microarray of claim 46 with the labeled polynucleotides of the sample under conditions suitable for the formation of a hybridization complex, and

- c) quantifying the expression of the polynucleotides in the sample..

48. An array comprising different nucleotide molecules affixed in distinct physical locations on a solid substrate, wherein at least one of said nucleotide molecules comprises a first oligonucleotide or polynucleotide sequence specifically hybridizable with at least 30 contiguous nucleotides of a target polynucleotide, and wherein said target polynucleotide is a polynucleotide of claim 12.

49. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 30 contiguous nucleotides of said target polynucleotide.

50. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 60 contiguous nucleotides of said target polynucleotide.

51. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to said target polynucleotide.

52. An array of claim 48, which is a microarray.

53. An array of claim 48, further comprising said target polynucleotide hybridized to a nucleotide molecule comprising said first oligonucleotide or polynucleotide sequence.

54. An array of claim 48, wherein a linker joins at least one of said nucleotide molecules to said solid substrate.

55. An array of claim 48, wherein each distinct physical location on the substrate contains multiple nucleotide molecules, and the multiple nucleotide molecules at any single distinct physical location have the same sequence, and each distinct physical location on the substrate contains nucleotide molecules having a sequence which differs from the sequence of nucleotide molecules at another distinct physical location on the substrate.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

PAGE MISSING AT THE TIME OF THE  
PUBLICATION

75. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.

76. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.

5 77. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:22.

78. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:23.

79. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.

10

80. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:25.

81. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:26.

15

82. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:27.

83. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:28.

84. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:29.

20

85. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:30.

86. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:31.

25

87. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:32.

88. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:33.

89. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:34.

30

90. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:35.

91. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:36.

92. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:37.

93. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:38.

5 94. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:39.

95. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:40.

10 96. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:41.

97. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:42.

98. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:43.

15 99. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:44.

100. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:45.

20 101. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:46.

102. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:47.

25 103. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:48.

30 104. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:49.

105. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:50.

106. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:51.

107. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
5 NO:52.

108. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:53.

109. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
10 NO:54.

110. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:55.

111. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
15 NO:56.

112. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
20 NO:57.

113. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:58.

114. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
25 NO:59.

115. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
NO:60.

116. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID  
30 NO:61.

117. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:62.



<110> INCYTE GENOMICS, INC.  
SPRAGUE, William W.  
CHAWLA, Narinder K.  
WARREN, Bridget A.  
TANG, Y. Tom  
ELLIOTT, Vicki S.  
MARQUIS, Joseph P.  
LI, Joana X.  
GRIFFIN, Jennifer A.  
GIETZEN, Kimberly J.  
YANG, Junming  
LU, Dyung Aina M.  
EMERLING, Brooke M.  
DUGGAN, Brendan M.  
RICHARDSON, Thomas W.  
LEE, Soo Yeun  
RAMKUMAR, Jayalaxmi  
BECHA, Shanya D.  
LEHR-MASON, Patricia M.  
SWARNAKAR, Anita  
TRAN, Uyen K.  
KABLE, Amy E.  
HAFALIA, April J.A.  
KHARE, Reena

<120> PROTEIN MODIFICATION AND MAINTENANCE MOLECULES

<130> PF-1186 PCT

<140> To Be Assigned

<141> Herewith

<150> US 60/322,196

<151> 2001-09-14

<150> US 60/324,134

<151> 2001-09-21

<150> US 60/327,233

<151> 2001-10-05

<150> US 60/346,198

<151> 2001-10-26

<150> US 60/343,980

<151> 2001-11-02

<150> US 60/348,887

<151> 2001-11-09

<150> US 60/332,423

<151> 2001-11-16

<150> US 60/334,145

<151> 2001-11-28

&lt;150&gt; US 60/334,229

&lt;151&gt; 2001-11-28

&lt;150&gt; US 60/337,451

&lt;151&gt; 2001-12-06

&lt;150&gt; US 60/351,928

&lt;151&gt; 2002-01-25

&lt;150&gt; US 60/366,837

&lt;151&gt; 2002-03-21

&lt;160&gt; 62

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 404

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 8268274CD1

&lt;400&gt; 1

```

Met Glu Thr His Ile Ser Cys Leu Phe Pro Glu Leu Leu Ala Met
  1           5           10           15
Ile Phe Gly Tyr Leu Asp Val Arg Asp Lys Gly Arg Ala Ala Gln
          20           25           30
Val Cys Thr Ala Trp Arg Asp Ala Ala Tyr His Lys Ser Val Trp
          35           40           45
Arg Gly Val Glu Ala Lys Leu His Leu Arg Arg Ala Asn Pro Ser
          50           55           60
Leu Phe Pro Ser Leu Gln Ala Arg Gly Ile Arg Arg Val Gln Ile
          65           70           75
Leu Ser Leu Arg Arg Ser Leu Ser Tyr Val Ile Gln Gly Met Ala
          80           85           90
Asn Ile Glu Ser Leu Asn Leu Ser Gly Cys Tyr Asn Leu Thr Asp
          95          100          105
Asn Gly Leu Gly His Ala Phe Val Gln Glu Ile Gly Ser Leu Arg
          110          115          120
Ala Leu Asn Leu Ser Leu Cys Lys Gln Ile Thr Asp Ser Ser Leu
          125          130          135
Gly Arg Ile Ala Gln Tyr Leu Lys Gly Leu Glu Val Leu Glu Leu
          140          145          150
Gly Gly Cys Ser Asn Ile Thr Asn Thr Gly Leu Leu Leu Ile Ala
          155          160          165
Trp Gly Leu Gln Arg Leu Lys Ser Leu Asn Leu Arg Ser Cys Arg
          170          175          180
His Leu Ser Asp Val Gly Ile Gly His Leu Ala Gly Met Thr Arg
          185          190          195
Ser Ala Ala Glu Gly Cys Leu Gly Leu Glu Gln Leu Thr Leu Gln
          200          205          210
Asp Cys Gln Lys Leu Thr Asp Leu Ser Leu Lys His Ile Ser Arg
          215          220          225

```

Gly	Leu	Thr	Gly	Leu	Arg	Leu	Leu	Asn	Leu	Ser	Phe	Cys	Gly	Gly
				230					235					240
Ile	Ser	Asp	Ala	Gly	Leu	Leu	His	Leu	Ser	His	Met	Gly	Ser	Leu
				245					250					255
Arg	Ser	Leu	Asn	Leu	Arg	Ser	Cys	Asp	Asn	Ile	Ser	Asp	Thr	Gly
				260					265					270
Ile	Met	His	Leu	Ala	Met	Gly	Ser	Leu	Arg	Leu	Ser	Gly	Leu	Asp
				275					280					285
Val	Ser	Phe	Cys	Asp	Lys	Val	Gly	Asp	Gln	Ser	Leu	Ala	Tyr	Ile
				290					295					300
Ala	Gln	Gly	Leu	Asp	Gly	Leu	Lys	Ser	Leu	Ser	Leu	Cys	Ser	Cys
				305					310					315
His	Ile	Ser	Asp	Asp	Gly	Ile	Asn	Arg	Met	Val	Arg	Gln	Met	His
				320					325					330
Gly	Leu	Arg	Thr	Leu	Asn	Ile	Gly	Gln	Cys	Val	Arg	Ile	Thr	Asp
				335					340					345
Lys	Gly	Leu	Glu	Leu	Ile	Ala	Glu	His	Leu	Ser	Gln	Leu	Thr	Gly
				350					355					360
Ile	Asp	Leu	Tyr	Gly	Cys	Thr	Arg	Ile	Thr	Lys	Arg	Gly	Leu	Glu
				365					370					375
Arg	Ile	Thr	Gln	Leu	Pro	Cys	Leu	Lys	Glu	Ala	Arg	Gly	Asp	Phe
				380					385					390
Ser	Pro	Leu	Phe	Thr	Val	Arg	Thr	Arg	Gly	Ser	Ser	Arg	Arg	
				395					400					

&lt;210&gt; 2

&lt;211&gt; 900

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500515CD1

&lt;400&gt; 2

Met	Lys	Pro	Pro	Arg	Pro	Val	Arg	Thr	Cys	Ser	Lys	Val	Leu	Val
1				5					10					15
Leu	Leu	Ser	Leu	Leu	Ala	Ile	His	Gln	Thr	Thr	Thr	Ala	Glu	Lys
				20					25					30
Asn	Gly	Ile	Asp	Ile	Tyr	Ser	Leu	Thr	Val	Asp	Ser	Arg	Val	Ser
				35					40					45
Ser	Arg	Phe	Ala	His	Thr	Val	Val	Thr	Ser	Arg	Val	Val	Asn	Arg
				50					55					60
Ala	Asn	Thr	Val	Gln	Glu	Ala	Thr	Phe	Gln	Met	Glu	Leu	Pro	Lys
				65					70					75
Lys	Ala	Phe	Ile	Thr	Asn	Phe	Ser	Met	Ile	Ile	Asp	Gly	Met	Thr
				80					85					90
Tyr	Pro	Gly	Ile	Ile	Lys	Glu	Lys	Ala	Glu	Ala	Gln	Ala	Gln	Tyr
				95					100					105
Ser	Ala	Ala	Val	Ala	Lys	Gly	Lys	Ser	Ala	Gly	Leu	Val	Lys	Ala
				110					115					120
Thr	Gly	Arg	Asn	Met	Glu	Gln	Phe	Gln	Val	Ser	Val	Ser	Val	Ala
				125					130					135
Pro	Asn	Ala	Lys	Ile	Thr	Phe	Glu	Leu	Val	Tyr	Glu	Glu	Leu	Leu
				140					145					150
Lys	Arg	Arg	Leu	Gly	Val	Tyr	Glu	Leu	Leu	Leu	Lys	Val	Arg	Pro

	155		160		165
Gln Gln Leu Val	Lys His Leu Gln Met	Asp Ile His Ile Phe	Glu		
	170		175		180
Pro Gln Gly Ile	Ser Phe Leu Glu Thr	Glu Ser Thr Phe Met	Thr		
	185		190		195
Asn Gln Leu Val	Asp Ala Leu Thr Thr	Trp Gln Asn Lys Thr	Lys		
	200		205		210
Ala His Ile Arg	Phe Lys Pro Thr Leu	Ser Gln Gln Gln Lys	Ser		
	215		220		225
Pro Glu Gln Gln	Glu Thr Val Leu Asp	Gly Asn Leu Ile Ile	Arg		
	230		235		240
Tyr Asp Val Asp	Arg Ala Ile Ser Gly	Gly Ser Ile Gln Ile	Glu		
	245		250		255
Asn Gly Tyr Phe	Val His Tyr Phe Ala	Pro Glu Gly Leu Thr	Thr		
	260		265		270
Met Pro Lys Asn	Val Val Phe Val Ile	Asp Lys Ser Gly Ser	Met		
	275		280		285
Ser Gly Arg Lys	Ile Gln Gln Thr Arg	Glu Ala Leu Ile Lys	Ile		
	290		295		300
Leu Asp Asp Leu	Ser Pro Arg Asp Gln	Phe Asn Leu Ile Val	Phe		
	305		310		315
Ser Thr Glu Ala	Thr Gln Trp Arg Pro	Ser Leu Val Pro Ala	Ser		
	320		325		330
Ala Glu Asn Val	Asn Lys Ala Arg Ser	Phe Ala Ala Gly Ile	Gln		
	335		340		345
Ala Leu Gly Gly	Thr Asn Ile Asn Asp	Ala Met Leu Met Ala	Val		
	350		355		360
Gln Leu Leu Asp	Ser Ser Asn Gln Glu	Glu Arg Leu Pro Glu	Gly		
	365		370		375
Ser Val Ser Leu	Ile Ile Leu Leu Thr	Asp Gly Asp Pro Thr	Val		
	380		385		390
Gly Glu Thr Asn	Pro Arg Ser Ile Gln	Asn Asn Val Arg Glu	Ala		
	395		400		405
Val Ser Gly Arg	Tyr Ser Leu Phe Cys	Leu Gly Phe Gly Phe	Asp		
	410		415		420
Val Ser Tyr Ala	Phe Leu Glu Lys Leu	Ala Leu Asp Asn Gly	Gly		
	425		430		435
Leu Ala Arg Arg	Ile His Glu Asp Ser	Asp Ser Ala Leu Gln	Leu		
	440		445		450
Gln Asp Phe Tyr	Gln Glu Val Ala Asn	Pro Leu Leu Thr Ala	Val		
	455		460		465
Thr Phe Glu Tyr	Pro Ser Asn Ala Val	Glu Glu Val Thr Gln	Asn		
	470		475		480
Asn Phe Arg Leu	Leu Phe Lys Gly Ser	Glu Met Val Val Ala	Gly		
	485		490		495
Lys Leu Gln Asp	Arg Gly Pro Asp Val	Leu Thr Ala Thr Val	Ser		
	500		505		510
Gly Lys Leu Pro	Thr Gln Asn Ile Thr	Phe Gln Thr Glu Ser	Ser		
	515		520		525
Val Ala Glu Gln	Glu Ala Glu Phe Gln	Ser Pro Lys Tyr Ile	Phe		
	530		535		540
His Asn Phe Met	Glu Arg Leu Trp Ala	Tyr Leu Thr Ile Gln	Gln		
	545		550		555
Leu Leu Glu Gln	Thr Val Ser Ala Ser	Asp Ala Asp Gln Gln	Ala		
	560		565		570
Leu Arg Asn Gln	Ala Leu Asn Leu Ser	Leu Ala Tyr Ser Phe	Val		

Thr	Pro	Leu	Thr	Ser	Met	Val	Val	Thr	Lys	Pro	Asp	Asp	Gln	Glu	575	580	585
															590	595	600
Gln	Ser	Gln	Val	Ala	Glu	Lys	Pro	Met	Glu	Gly	Glu	Ser	Arg	Asn	605	610	615
Arg	Asn	Val	His	Ser	Ala	Gly	Ala	Ala	Gly	Ser	Arg	Met	Asn	Phe	620	625	630
Arg	Pro	Gly	Val	Leu	Ser	Ser	Arg	Gln	Leu	Gly	Leu	Pro	Gly	Pro	635	640	645
Pro	Asp	Val	Pro	Asp	His	Ala	Ala	Tyr	His	Pro	Phe	Arg	Arg	Leu	650	655	660
Ala	Ile	Leu	Pro	Ala	Ser	Ala	Pro	Pro	Ala	Thr	Ser	Asn	Pro	Asp	665	670	675
Pro	Ala	Val	Ser	Arg	Val	Met	Asn	Met	Lys	Ile	Glu	Glu	Thr	Thr	680	685	690
Met	Thr	Thr	Gln	Thr	Pro	Ala	Pro	Ile	Gln	Ala	Pro	Ser	Ala	Ile	695	700	705
Leu	Pro	Leu	Pro	Gly	Gln	Ser	Val	Glu	Arg	Leu	Cys	Val	Asp	Pro	710	715	720
Arg	His	Arg	Gln	Gly	Pro	Val	Asn	Leu	Leu	Ser	Asp	Pro	Glu	Gln	725	730	735
Gly	Val	Glu	Val	Thr	Gly	Gln	Tyr	Glu	Arg	Glu	Lys	Ala	Gly	Phe	740	745	750
Ser	Trp	Ile	Glu	Val	Thr	Phe	Lys	Asn	Pro	Leu	Val	Trp	Val	His	755	760	765
Ala	Ser	Pro	Glu	His	Val	Val	Val	Thr	Arg	Asn	Arg	Arg	Ser	Ser	770	775	780
Ala	Tyr	Lys	Trp	Lys	Glu	Thr	Leu	Phe	Ser	Val	Met	Pro	Gly	Leu	785	790	795
Lys	Met	Thr	Met	Asp	Lys	Thr	Gly	Leu	Leu	Leu	Leu	Ser	Asp	Pro	800	805	810
Asp	Lys	Val	Thr	Ile	Gly	Leu	Leu	Phe	Trp	Asp	Gly	Arg	Gly	Glu	815	820	825
Gly	Leu	Arg	Leu	Leu	Leu	Arg	Asp	Thr	Asp	Arg	Phe	Ser	Ser	His	830	835	840
Val	Gly	Gly	Thr	Leu	Gly	Gln	Phe	Tyr	Gln	Glu	Val	Leu	Trp	Gly	845	850	855
Ser	Pro	Ala	Ala	Ser	Asp	Asp	Gly	Arg	Arg	Thr	Leu	Arg	Val	Gln	860	865	870
Gly	Asn	Asp	His	Ser	Ala	Thr	Arg	Glu	Arg	Arg	Leu	Asp	Tyr	Gln	875	880	885
Glu	Gly	Pro	Pro	Gly	Val	Glu	Ile	Ser	Cys	Trp	Ser	Val	Glu	Leu	890	895	900

&lt;210&gt; 3

&lt;211&gt; 436

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2256826CD1

&lt;400&gt; 3

Met Arg Arg Asp Val Asn Gly Val Thr Lys Ser Arg Phe Glu Met

1	5	10	15
Phe Ser Asn Ser Asp	Glu Ala Val Ile Asn Lys Lys Leu Pro Lys		
20	25	30	
Glu Leu Leu Leu Arg	Ile Phe Ser Phe Leu Asp Val Val Thr Leu		
35	40	45	
Cys Arg Cys Ala Gln	Val Ser Arg Ala Trp Asn Val Leu Ala Leu		
50	55	60	
Asp Gly Ser Asn Trp	Gln Arg Ile Asp Leu Phe Asp Phe Gln Arg		
65	70	75	
Asp Ile Glu Gly Arg	Val Val Glu Asn Ile Ser Lys Arg Cys Gly		
80	85	90	
Gly Phe Leu Arg Lys	Leu Ser Leu Arg Gly Cys Leu Gly Val Gly		
95	100	105	
Asp Asn Ala Leu Arg	Thr Phe Ala Gln Asn Cys Arg Asn Ile Glu		
110	115	120	
Val Leu Asn Leu Asn	Gly Cys Thr Lys Thr Thr Asp Ala Thr Cys		
125	130	135	
Thr Ser Leu Ser Lys	Phe Cys Ser Lys Leu Arg His Leu Asp Leu		
140	145	150	
Ala Ser Cys Thr Ser	Ile Thr Asn Met Ser Leu Lys Ala Leu Ser		
155	160	165	
Glu Gly Cys Pro Leu	Leu Glu Gln Leu Asn Ile Ser Trp Cys Asp		
170	175	180	
Gln Val Thr Lys Asp	Gly Ile Gln Ala Leu Val Arg Gly Cys Gly		
185	190	195	
Gly Leu Lys Ala Leu	Phe Leu Lys Gly Cys Thr Gln Leu Glu Asp		
200	205	210	
Glu Ala Leu Lys Tyr	Ile Gly Ala His Cys Pro Glu Leu Val Thr		
215	220	225	
Leu Asn Leu Gln Thr	Cys Leu Gln Ile Thr Asp Glu Gly Leu Ile		
230	235	240	
Thr Ile Cys Arg Gly	Cys His Lys Leu Gln Ser Leu Cys Ala Ser		
245	250	255	
Gly Cys Ser Asn Ile	Thr Asp Ala Ile Leu Asn Ala Leu Gly Gln		
260	265	270	
Asn Cys Pro Arg Leu	Arg Ile Leu Glu Val Ala Arg Cys Ser Gln		
275	280	285	
Leu Thr Asp Val Gly	Phe Thr Thr Leu Ala Arg Asn Cys His Glu		
290	295	300	
Leu Glu Lys Met Asp	Leu Glu Glu Cys Val Gln Ile Thr Asp Ser		
305	310	315	
Thr Leu Ile Gln Leu	Ser Ile His Cys Pro Arg Leu Gln Val Leu		
320	325	330	
Ser Leu Ser His Cys	Glu Leu Ile Thr Asp Asp Gly Ile Arg His		
335	340	345	
Leu Gly Asn Gly Ala	Cys Ala His Asp Gln Leu Glu Val Ile Glu		
350	355	360	
Leu Asp Asn Cys Pro	Leu Ile Thr Asp Ala Ser Leu Glu His Leu		
365	370	375	
Lys Ser Cys His Ser	Leu Glu Arg Ile Glu Leu Tyr Asp Cys Gln		
380	385	390	
Gln Ile Thr Arg Ala	Gly Ile Lys Arg Leu Arg Thr His Leu Pro		
395	400	405	
Asn Ile Lys Val His	Ala Tyr Phe Ala Pro Val Thr Pro Pro Pro		
410	415	420	
Ser Val Gly Gly Ser	Arg Gln Arg Phe Cys Arg Cys Cys Ile Ile		

	425		430		435
Leu					
<210>	4				
<211>	356				
<212>	PRT				
<213>	Homo sapiens				
<220>					
<221>	misc_feature				
<223>	Incyte ID No: 7686186CD1				
<400>	4				
Met Glu Asn Asp Val Thr Tyr Pro Asp Pro Tyr Ser Arg Pro Ala					
1 5 10 15					
Pro Asp Arg Phe Ile Arg Arg Trp Leu Val Ile Thr Gly Cys Ile					
20 25 30					
Ala Ala Leu Met Leu Leu Trp Gln Phe Leu Pro Ala Ile Glu Ala					
35 40 45					
Trp Phe Ser Pro His Glu Thr Gln Glu Arg Thr Val Thr Pro Arg					
50 55 60					
Gly Asp Leu Ala Ala Asp Glu Lys Thr Thr Ile Glu Leu Phe Glu					
65 70 75					
Lys Ser Arg Gly Ser Val Val Tyr Ile Thr Thr Ala Gln Leu Val					
80 85 90					
Arg Asp Val Trp Ser Arg Asn Val Phe Ser Val Pro Arg Gly Thr					
95 100 105					
Gly Ser Gly Phe Ile Trp Asp Asp Ala Gly His Val Val Thr Asn					
110 115 120					
Phe His Val Ile Gln Gly Ala Ser Ser Ala Thr Val Lys Leu Ala					
125 130 135					
Asp Gly Arg Asp Tyr Gln Ala Ala Leu Val Gly Ala Ser Pro Ala					
140 145 150					
His Asp Ile Ala Val Leu Lys Ile Gly Val Gly Phe Lys Arg Pro					
155 160 165					
Pro Ala Val Pro Val Gly Thr Ser Ala Asp Leu Lys Val Gly Gln					
170 175 180					
Lys Val Phe Ala Ile Gly Asn Pro Phe Gly Leu Asp Trp Thr Leu					
185 190 195					
Thr Thr Gly Ile Val Ser Ala Leu Asp Arg Thr Leu Ser Gly Asp					
200 205 210					
Ala Ser Gly Pro Ala Ile Asp His Leu Ile Gln Thr Asp Ala Ala					
215 220 225					
Ile Asn Pro Gly Asn Ser Gly Gly Pro Leu Leu Asp Ser Ala Gly					
230 235 240					
Arg Leu Ile Gly Ile Asn Thr Ala Ile Tyr Ser Pro Ser Gly Ala					
245 250 255					
Ser Ala Gly Ile Gly Phe Ala Val Pro Val Asp Thr Val Met Arg					
260 265 270					
Val Val Pro Gln Leu Ile Lys Thr Gly Lys Tyr Ile Arg Pro Ala					
275 280 285					
Leu Gly Ile Glu Val Asp Glu Gln Leu Asn Ala Arg Leu Gln Ala					
290 295 300					
Leu Thr Gly Ser Lys Gly Val Phe Val Leu Arg Val Thr Pro Gly					
305 310 315					

Ser	Ala	Ala	His	Arg	Ala	Gly	Leu	Val	Gly	Val	Glu	Val	Thr	Ala
				320					325					330
Gly	Gly	Ile	Val	Pro	Gly	Asp	Arg	Val	Ile	Ser	Ile	Asp	Gly	Ile
				335					340					345
Ala	Val	Asp	Pro	Gly	Ile	Pro	Asp	Arg	Thr	Cys				
				350					355					

&lt;210&gt; 5

&lt;211&gt; 432

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 72617436CD1

&lt;400&gt; 5

Met	Gly	Pro	Ser	Ser	Leu	Arg	Lys	Thr	Ser	Ser	Gly	Leu	Pro	Leu
1				5					10					15
Ile	Leu	His	Tyr	Gly	Val	Ile	Leu	Gly	Ala	Pro	Leu	Ala	Ser	Ser
				20					25					30
Cys	Ala	Gly	Ala	Cys	Gly	Thr	Ser	Phe	Pro	Asp	Gly	Leu	Thr	Pro
				35					40					45
Glu	Gly	Thr	Gln	Ala	Ser	Gly	Asp	Lys	Asp	Ile	Pro	Ala	Ile	Asn
				50					55					60
Gln	Gly	Leu	Ile	Leu	Glu	Glu	Thr	Pro	Glu	Ser	Ser	Phe	Leu	Ile
				65					70					75
Glu	Gly	Asp	Ile	Ile	Arg	Pro	Ser	Pro	Phe	Arg	Leu	Leu	Ser	Ala
				80					85					90
Thr	Ser	Asn	Lys	Trp	Pro	Met	Gly	Gly	Ser	Gly	Val	Val	Glu	Val
				95					100					105
Pro	Phe	Leu	Leu	Ser	Ser	Lys	Tyr	Asp	Glu	Pro	Ser	Arg	Gln	Val
				110					115					120
Ile	Leu	Glu	Ala	Leu	Ala	Glu	Phe	Glu	Arg	Ser	Thr	Cys	Ile	Arg
				125					130					135
Phe	Val	Thr	Tyr	Gln	Asp	Gln	Arg	Asp	Phe	Ile	Ser	Ile	Ile	Pro
				140					145					150
Met	Tyr	Gly	Cys	Phe	Ser	Ser	Val	Gly	Arg	Ser	Gly	Gly	Met	Gln
				155					160					165
Val	Val	Ser	Leu	Ala	Pro	Thr	Cys	Leu	Gln	Lys	Gly	Arg	Gly	Ile
				170					175					180
Val	Leu	His	Glu	Leu	Met	His	Val	Leu	Gly	Phe	Trp	His	Glu	His
				185					190					195
Thr	Arg	Ala	Asp	Arg	Asp	Arg	Tyr	Ile	Arg	Val	Asn	Trp	Asn	Glu
				200					205					210
Ile	Leu	Pro	Gly	Phe	Glu	Ile	Asn	Phe	Ile	Lys	Ser	Arg	Ser	Ser
				215					220					225
Asn	Met	Leu	Thr	Pro	Tyr	Asp	Tyr	Ser	Ser	Val	Met	His	Tyr	Gly
				230					235					240
Arg	Leu	Ala	Phe	Ser	Arg	Arg	Gly	Leu	Pro	Thr	Ile	Thr	Pro	Leu
				245					250					255
Trp	Ala	Pro	Ser	Val	His	Ile	Gly	Gln	Arg	Trp	Asn	Leu	Ser	Ala
				260					265					270
Ser	Asp	Ile	Thr	Arg	Val	Leu	Gln	Leu	Tyr	Gly	Cys	Ser	Pro	Ser
				275					280					285
Gly	Pro	Arg	Pro	Arg	Gly	Arg	Gly	Ser	His	Ala	His	Ser	Thr	Gly



290	295	300
Arg Ser Pro Ala Pro Ala Ser Leu Ser	Leu Gln Arg Leu Leu Glu	
305	310	315
Ala Leu Ser Ala Glu Ser Arg Ser Pro	Asp Pro Ser Gly Ser Ser	
320	325	330
Ala Gly Gly Gln Pro Val Pro Ala Gly	Pro Gly Glu Ser Pro His	
335	340	345
Gly Trp Glu Ser Pro Ala Leu Lys Lys	Leu Ser Ala Glu Ala Ser	
350	355	360
Ala Arg Gln Pro Gln Thr Leu Ala Ser	Ser Pro Arg Ser Arg Pro	
365	370	375
Gly Ala Gly Ala Pro Gly Val Ala Gln	Glu Gln Ser Trp Leu Ala	
380	385	390
Gly Val Ser Thr Lys Pro Thr Val Pro	Ser Ser Glu Ala Gly Ile	
395	400	405
Gln Pro Val Pro Val Gln Gly Ser Pro	Ala Leu Pro Gly Gly Cys	
410	415	420
Val Pro Arg Asn His Phe Lys Gly Met	Ser Glu Asp	
425	430	

&lt;210&gt; 6

&lt;211&gt; 248

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7501945CD1

&lt;400&gt; 6

Met Gly Pro Ser Ser Leu Arg Lys Thr	Ser Ser Gly Leu Pro Leu
1	5 10 15
Ile Leu His Tyr Gly Val Ile Leu Gly	Ala Pro Leu Ala Ser Ser
20	25 30
Cys Ala Gly Ala Cys Gly Thr Ser Phe	Pro Asp Gly Leu Thr Pro
35	40 45
Glu Gly Thr Gln Ala Ser Gly Asp Lys	Asp Ile Pro Ala Ile Asn
50	55 60
Gln Gly Leu Ile Leu Glu Glu Thr Pro	Glu Ser Ser Phe Leu Ile
65	70 75
Glu Gly Asp Ile Ile Arg Pro Ser Pro	Phe Arg Leu Leu Ser Ala
80	85 90
Thr Ser Asn Lys Trp Pro Met Gly Gly	Ser Gly Val Val Glu Val
95	100 105
Pro Phe Leu Leu Ser Ser Lys Tyr Asp	Glu Pro Ser Arg Gln Val
110	115 120
Ile Leu Glu Ala Leu Ala Glu Phe Glu	Arg Ser Thr Cys Ile Arg
125	130 135
Phe Val Thr Tyr Gln Asp Gln Arg Asp	Phe Ile Ser Ile Ile Pro
140	145 150
Met Tyr Gly Cys Phe Ser Ser Val Gly	Arg Ser Gly Gly Met Gln
155	160 165
Val Val Ser Leu Ala Pro Thr Cys Leu	Gln Lys Gly Arg Gly Ile
170	175 180
Val Leu His Glu Leu Met His Val Leu	Gly Phe Trp His Glu His
185	190 195

```

Thr Arg Ala Asp Arg Asp Arg Tyr Ile His Val Asn Trp Asn Glu
      200                      205                      210
Ile Leu Pro Gly Phe Glu Ile Asn Phe Ile Lys Ser Arg Ser Ser
      215                      220                      225
Asn Met Leu Thr Pro Tyr Asp Tyr Ser Ser Val Met His Tyr Gly
      230                      235                      240
Arg Val Pro Cys Pro Gln His Trp
      245

```

&lt;210&gt; 7

&lt;211&gt; 388

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500264CD1

&lt;400&gt; 7

```

Met Ala Ser Val Ala Gln Glu Ser Ala Gly Ser Gln Arg Arg Leu
  1          5          10          15
Pro Pro Arg His Gly Ala Leu Arg Gly Leu Leu Leu Leu Cys Leu
      20          25          30
Trp Leu Pro Ser Gly Arg Ala Ala Leu Pro Pro Ala Ala Pro Leu
      35          40          45
Ser Glu Leu His Ala Gln Leu Ser Gly Val Glu Gln Leu Leu Glu
      50          55          60
Glu Phe Arg Arg Gln Leu Gln Gln Glu Arg Pro Gln Glu Glu Leu
      65          70          75
Glu Leu Glu Leu Arg Ala Gly Gly Gly Pro Gln Glu Asp Cys Pro
      80          85          90
Gly Pro Gly Ser Gly Gly Tyr Ser Ala Met Pro Asp Ala Ile Ile
      95          100         105
Arg Thr Lys Asp Ser Leu Ala Ala Gly Ala Ser Phe Leu Arg Ala
     110         115         120
Pro Ala Ala Val Arg Gly Trp Arg Gln Cys Val Ala Ala Cys Cys
     125         130         135
Ser Glu Pro Arg Cys Ser Val Ala Val Val Glu Leu Pro Arg Arg
     140         145         150
Pro Ala Pro Pro Ala Ala Val Leu Gly Cys Tyr Leu Phe Asn Cys
     155         160         165
Thr Ala Arg Gly Arg Asn Val Cys Lys Phe Ala Leu His Ser Gly
     170         175         180
Tyr Ser Ser Tyr Ser Leu Ser Arg Ala Pro Asp Gly Ala Ala Leu
     185         190         195
Ala Thr Ala Arg Ala Ser Pro Arg Gln Glu Lys Asp Ala Pro Pro
     200         205         210
Leu Ser Lys Ala Gly Gln Asp Val Val Leu His Leu Pro Thr Asp
     215         220         225
Gly Val Val Leu Asp Gly Arg Glu Ser Thr Asp Asp His Ala Ile
     230         235         240
Val Gln Tyr Glu Trp Ala Leu Leu Gln Gly Asp Pro Ser Val Asp
     245         250         255
Met Lys Val Pro Gln Ser Gly Gly Asp Ser Leu Val Glu Lys Ser
     260         265         270
Gln Lys Ala Thr Ala Pro Asn Lys Pro Pro Ala Leu Ser Asn Thr

```

275	280	285
Glu Lys Arg Asn His Ser Ala Phe Trp	Gly Pro Glu Ser Gln Ile	
290	295	300
Ile Pro Val Met Pro Asp Ser Ser Ser	Ser Gly Lys Asn Arg Lys	
305	310	315
Glu Glu Ser Tyr Ile Phe Glu Ser Lys	Gly Asp Gly Gly Gly Gly	
320	325	330
Glu His Pro Ala Pro Glu Thr Gly Ala Val	Leu Pro Leu Ala Leu	
335	340	345
Gly Leu Ala Ile Thr Ala Leu Leu Leu	Leu Met Val Ala Cys Arg	
350	355	360
Leu Arg Leu Val Lys Gln Lys Leu Lys	Lys Ala Arg Pro Ile Thr	
365	370	375
Ser Glu Glu Ser Asp Tyr Leu Ile Asn	Gly Met Tyr Leu	
380	385	

&lt;210&gt; 8

&lt;211&gt; 467

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7499935CD1

&lt;400&gt; 8

Met Ala Ala Ala Thr Gly Pro Ser Phe Trp	Leu Gly Asn Glu Thr
1	5 10 15
Leu Lys Val Pro Leu Ala Leu Phe Ala Leu	Asn Arg Gln Arg Leu
20	25 30
Cys Glu Arg Leu Arg Lys Asn Pro Ala Val	Gln Ala Gly Ser Ile
35	40 45
Val Val Leu Gln Gly Gly Glu Glu Thr	Gln Arg Tyr Cys Thr Asp
50	55 60
Thr Gly Val Leu Phe Arg Gln Glu Ser Phe	Phe His Trp Ala Phe
65	70 75
Gly Val Thr Glu Pro Gly Cys Tyr Gly Val	Ile Asp Val Asp Thr
80	85 90
Gly Lys Ser Thr Leu Phe Val Pro Arg Leu	Pro Ala Ser His Ala
95	100 105
Thr Trp Met Gly Lys Ile His Ser Lys Glu	His Phe Lys Glu Lys
110	115 120
Tyr Ala Val Asp Asp Val Gln Tyr Val Asp	Glu Ile Ala Ser Val
125	130 135
Leu Thr Ser Gln Lys Pro Ser Val Leu Leu	Thr Leu Arg Gly Val
140	145 150
Asn Thr Asp Ser Gly Ser Val Cys Arg Glu	Ala Ser Phe Asp Gly
155	160 165
Ile Ser Lys Phe Glu Val Asn Asn Thr Ile	Leu His Pro Glu Ile
170	175 180
Val Glu Cys Arg Val Phe Lys Thr Asp Met	Glu Leu Glu Val Leu
185	190 195
Arg Tyr Thr Asn Lys Ile Ser Ser Glu Ala	His Arg Glu Val Met
200	205 210
Lys Ala Val Lys Val Gly Met Lys Glu Tyr	Glu Leu Glu Ser Leu
215	220 225

Phe Glu His Tyr Cys Tyr Ser Arg Gly Gly Met Arg His Ser Ser  
 230 235 240  
 Tyr Thr Cys Ile Cys Gly Ser Gly Glu Asn Ser Ala Val Leu His  
 245 250 255  
 Tyr Gly His Ala Gly Ala Pro Asn Asp Arg Thr Ile Gln Asn Gly  
 260 265 270  
 Asp Met Cys Leu Phe Asp Met Gly Gly Glu Tyr Tyr Cys Phe Ala  
 275 280 285  
 Ser Asp Ile Thr Cys Ser Phe Pro Ala Asn Gly Lys Phe Thr Ala  
 290 295 300  
 Asp Gln Lys Ala Val Tyr Glu Ala Val Leu Arg Ser Ser Arg Ala  
 305 310 315  
 Val Met Gly Ala Met Lys Pro Gly Val Trp Trp Pro Asp Met His  
 320 325 330  
 Arg Leu Ala Asp Arg Ile His Leu Glu Glu Leu Ala His Met Gly  
 335 340 345  
 Ile Leu Ser Gly Ser Val Asp Ala Met Val Gln Ala His Leu Gly  
 350 355 360  
 Ala Val Phe Met Pro His Gly Leu Gly His Phe Leu Gly Ile Asp  
 365 370 375  
 Val His Asp Val Gly Gly Tyr Pro Glu Gly Val Glu Arg Ile Tyr  
 380 385 390  
 Phe Ile Asp His Leu Leu Asp Glu Ala Leu Ala Asp Pro Ala Arg  
 395 400 405  
 Ala Ser Phe Leu Asn Arg Glu Val Leu Gln Arg Phe Arg Gly Phe  
 410 415 420  
 Gly Gly Val Arg Ile Glu Glu Asp Val Val Val Thr Asp Ser Gly  
 425 430 435  
 Ile Glu Leu Leu Thr Cys Val Pro Arg Thr Val Glu Glu Ile Glu  
 440 445 450  
 Ala Cys Met Ala Gly Cys Asp Lys Ala Phe Thr Pro Phe Ser Gly  
 455 460 465  
 Pro Lys

&lt;210&gt; 9

&lt;211&gt; 379

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7982285CD1

&lt;400&gt; 9

Met Glu Gly Asn Arg Asp Glu Ala Glu Lys Cys Val Glu Ile Ala  
 1 5 10 15  
 Arg Glu Ala Leu Asn Ala Gly Asn Arg Glu Lys Ala Gln Arg Phe  
 20 25 30  
 Leu Gln Lys Ala Glu Lys Leu Tyr Pro Leu Pro Ser Ala Arg Ala  
 35 40 45  
 Leu Leu Glu Ile Ile Met Lys Asn Gly Ser Thr Ala Gly Asn Ser  
 50 55 60  
 Pro His Cys Arg Lys Pro Ser Gly Ser Gly Asp Gln Ser Lys Pro  
 65 70 75  
 Asn Cys Thr Lys Asp Ser Thr Ser Gly Ser Gly Glu Gly Gly Lys

80	85	90
Gly Tyr Thr Lys Asp Gln Val Asp Gly Val Leu Ser Ile Asn Lys		
95	100	105
Cys Lys Asn Tyr Tyr Glu Val Leu Gly Val Thr Lys Asp Ala Gly		
110	115	120
Asp Glu Asp Leu Lys Lys Ala Tyr Arg Lys Leu Ala Leu Lys Phe		
125	130	135
His Pro Asp Lys Asn His Ala Pro Gly Ala Thr Asp Ala Phe Lys		
140	145	150
Lys Ile Gly Asn Ala Tyr Ala Val Leu Ser Asn Pro Glu Lys Arg		
155	160	165
Lys Gln Tyr Asp Leu Thr Gly Asn Glu Glu Gln Ala Cys Asn His		
170	175	180
Gln Asn Asn Gly Arg Phe Asn Phe His Arg Gly Cys Glu Ala Asp		
185	190	195
Ile Thr Pro Glu Asp Leu Phe Asn Ile Phe Phe Gly Gly Gly Phe		
200	205	210
Pro Ser Gly Ser Val His Ser Phe Ser Asn Gly Arg Ala Gly Tyr		
215	220	225
Ser Gln Gln His Gln His Arg His Ser Gly His Glu Arg Glu Glu		
230	235	240
Glu Arg Gly Asp Gly Gly Phe Ser Val Phe Ile Gln Leu Met Pro		
245	250	255
Ile Ile Val Leu Ile Leu Val Ser Leu Leu Ser Gln Leu Met Val		
260	265	270
Ser Asn Pro Pro Tyr Ser Leu Tyr Pro Arg Ser Gly Thr Gly Gln		
275	280	285
Thr Ile Lys Met Gln Thr Glu Asn Leu Gly Val Val Tyr Tyr Val		
290	295	300
Asn Lys Asp Phe Lys Asn Glu Tyr Lys Gly Met Leu Leu Gln Lys		
305	310	315
Val Glu Lys Ser Val Glu Glu Asp Tyr Val Thr Asn Ile Arg Asn		
320	325	330
Asn Cys Trp Lys Glu Arg Gln Gln Lys Thr Asp Met Gln Tyr Ala		
335	340	345
Ala Lys Val Tyr Arg Asp Asp Arg Leu Arg Arg Lys Ala Asp Ala		
350	355	360
Leu Ser Met Asp Asn Cys Lys Glu Leu Glu Arg Leu Thr Ser Leu		
365	370	375
Tyr Lys Gly Gly		

<210> 10  
 <211> 737  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7758505CD1

<400> 10  
 Met Gly Val Leu Lys Val Trp Leu Gly Leu Ala Leu Ala Leu Ala  
 1 5 10 15  
 Glu Phe Ala Val Leu Pro His His Ser Glu Gly Ala Cys Val Tyr  
 20 25 30

Gln	Asp	Ser	Leu	Leu	Ala	Asp	Ala	Thr	Ile	Trp	Lys	Pro	Asp	Ser
				35					40					45
Cys	Gln	Ser	Cys	Arg	Cys	His	Gly	Asp	Ile	Val	Ile	Cys	Lys	Pro
				50					55					60
Ala	Val	Cys	Arg	Asn	Pro	Gln	Cys	Ala	Phe	Glu	Lys	Gly	Glu	Val
				65					70					75
Leu	Gln	Ile	Ala	Ala	Asn	Gln	Cys	Cys	Pro	Glu	Cys	Val	Leu	Arg
				80					85					90
Thr	Pro	Gly	Ser	Cys	His	His	Glu	Lys	Lys	Ile	His	Glu	His	Gly
				95					100					105
Thr	Glu	Trp	Ala	Ser	Ser	Pro	Cys	Ser	Val	Cys	Ser	Cys	Asn	His
				110					115					120
Gly	Glu	Val	Arg	Cys	Thr	Pro	Gln	Pro	Cys	Pro	Pro	Leu	Ser	Cys
				125					130					135
Gly	His	Gln	Glu	Leu	Ala	Phe	Ile	Pro	Glu	Gly	Ser	Cys	Cys	Pro
				140					145					150
Val	Cys	Val	Gly	Leu	Gly	Lys	Pro	Cys	Ser	Tyr	Glu	Gly	His	Val
				155					160					165
Phe	Gln	Asp	Gly	Glu	Asp	Trp	Arg	Leu	Ser	Arg	Cys	Ala	Lys	Cys
				170					175					180
Leu	Cys	Arg	Asn	Gly	Val	Ala	Gln	Cys	Phe	Thr	Ala	Gln	Cys	Gln
				185					190					195
Pro	Leu	Phe	Cys	Asn	Gln	Asp	Glu	Thr	Val	Val	Arg	Val	Pro	Gly
				200					205					210
Lys	Cys	Cys	Pro	Gln	Cys	Ser	Ala	Arg	Ser	Cys	Ser	Ala	Ala	Gly
				215					220					225
Gln	Val	Tyr	Glu	His	Gly	Glu	Gln	Trp	Ser	Glu	Asn	Ala	Cys	Thr
				230					235					240
Thr	Cys	Ile	Cys	Asp	Arg	Gly	Glu	Val	Arg	Cys	His	Lys	Gln	Ala
				245					250					255
Cys	Leu	Pro	Leu	Arg	Cys	Gly	Lys	Gly	Gln	Ser	Arg	Ala	Arg	Arg
				260					265					270
His	Gly	Gln	Cys	Cys	Glu	Glu	Cys	Val	Ser	Pro	Ala	Gly	Ser	Cys
				275					280					285
Ser	Tyr	Asp	Gly	Val	Val	Arg	Tyr	Gln	Asp	Glu	Met	Trp	Lys	Gly
				290					295					300
Ser	Ala	Cys	Glu	Phe	Cys	Met	Cys	Asp	His	Gly	Gln	Val	Thr	Cys
				305					310					315
Gln	Thr	Gly	Glu	Cys	Ala	Lys	Val	Glu	Cys	Ala	Arg	Asp	Glu	Glu
				320					325					330
Leu	Ile	His	Leu	Asp	Gly	Lys	Cys	Cys	Pro	Glu	Cys	Ile	Ser	Arg
				335					340					345
Asn	Gly	Tyr	Cys	Val	Tyr	Glu	Glu	Thr	Gly	Glu	Phe	Met	Ser	Ser
				350					355					360
Asn	Ala	Ser	Glu	Val	Lys	Arg	Ile	Pro	Glu	Gly	Glu	Lys	Trp	Glu
				365					370					375
Asp	Gly	Pro	Cys	Lys	Val	Cys	Glu	Cys	Arg	Gly	Ala	Gln	Val	Thr
				380					385					390
Cys	Tyr	Glu	Pro	Ser	Cys	Pro	Pro	Cys	Pro	Val	Gly	Thr	Leu	Ala
				395					400					405
Leu	Glu	Val	Lys	Gly	Gln	Cys	Cys	Pro	Asp	Cys	Thr	Ser	Val	His
				410					415					420
Cys	His	Pro	Asp	Cys	Leu	Thr	Cys	Ser	Gln	Ser	Pro	Asp	His	Cys
				425					430					435
Asp	Leu	Cys	Gln	Asp	Pro	Thr	Lys	Leu	Leu	Gln	Asn	Gly	Trp	Cys
				440					445					450

Val His Ser Cys Gly Leu Gly Phe Tyr Gln Ala Gly Ser Leu Cys  
 455 460 465  
 Ile Ala Cys Gln Pro Gln Cys Ser Thr Cys Thr Ser Gly Leu Glu  
 470 475 480  
 Cys Ser Ser Cys Gln Pro Pro Leu Leu Met Arg His Gly Gln Cys  
 485 490 495  
 Val Pro Thr Cys Gly Asp Gly Phe Tyr Gln Asp Arg His Ser Cys  
 500 505 510  
 Ala Val Cys His Glu Ser Cys Ala Gly Cys Trp Gly Pro Thr Glu  
 515 520 525  
 Lys His Cys Leu Ala Cys Arg Asp Pro Leu His Val Leu Arg Asp  
 530 535 540  
 Gly Gly Cys Glu Ser Ser Cys Gly Lys Gly Phe Tyr Asn Arg Gln  
 545 550 555  
 Gly Thr Cys Ser Ala Cys Asp Gln Ser Cys Asp Ser Cys Gly Pro  
 560 565 570  
 Ser Ser Pro Arg Cys Leu Thr Cys Thr Glu Lys Thr Val Leu His  
 575 580 585  
 Asp Gly Lys Cys Met Ser Glu Cys Pro Gly Gly Tyr Tyr Ala Asp  
 590 595 600  
 Ala Thr Gly Arg Cys Lys Val Cys His Asn Ser Cys Ala Ser Cys  
 605 610 615  
 Ser Gly Pro Thr Pro Ser His Cys Thr Ala Cys Ser Pro Pro Lys  
 620 625 630  
 Ala Leu Arg Gln Gly His Cys Leu Pro Arg Cys Gly Glu Gly Phe  
 635 640 645  
 Tyr Ser Asp His Gly Val Cys Lys Ala Cys His Ser Ser Cys Leu  
 650 655 660  
 Ala Cys Met Gly Pro Ala Pro Ser His Cys Thr Gly Cys Lys Lys  
 665 670 675  
 Pro Glu Glu Gly Leu Gln Val Glu Gln Leu Ser Gly Val Gly Ile  
 680 685 690  
 Pro Ser Gly Glu Cys Leu Ala Gln Cys Arg Ala His Phe Tyr Leu  
 695 700 705  
 Glu Ser Thr Gly Leu Cys Glu Gly Gln Asn Leu Asp Phe Cys Gln  
 710 715 720  
 Asn Leu Glu Val Ile Ser Ala Val Cys Leu Gly Ile Ser Ser Thr  
 725 730 735  
 Glu Asn

<210> 11  
 <211> 530  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 6885756CD1

<400> 11  
 Met Glu Asp Asp Ser Leu Tyr Leu Gly Gly Asp Trp Gln Phe Asn  
 1 5 10 15  
 His Phe Ser Lys Leu Thr Ser Ser Arg Leu Asp Ala Ala Phe Ala  
 20 25 30  
 Glu Ile Gln Arg Thr Ser Leu Ser Glu Lys Ser Pro Leu Ser Ser

	35	40	45
Glu Thr Arg Phe Asp Leu Cys Asp Asp Leu Ala Pro Val Ala Arg			
	50	55	60
Gln Leu Ala Pro Arg Glu Lys Leu Pro Leu Ser Ser Arg Arg Pro			
	65	70	75
Ala Ala Val Gly Ala Gly Leu Gln Lys Ile Gly Asn Thr Phe Tyr			
	80	85	90
Val Asn Val Ser Leu Gln Cys Leu Thr Tyr Thr Leu Pro Leu Ser			
	95	100	105
Asn Tyr Met Leu Ser Arg Glu Asp Ser Gln Thr Cys His Leu His			
	110	115	120
Lys Cys Cys Met Phe Cys Thr Met Gln Ala His Ile Thr Trp Ala			
	125	130	135
Leu Tyr Arg Pro Gly His Val Ile Gln Pro Ser Gln Val Leu Ala			
	140	145	150
Ala Gly Phe His Arg Gly Glu Gln Glu Asp Ala His Glu Phe Leu			
	155	160	165
Met Phe Thr Val Asp Ala Met Lys Lys Ala Cys Leu Pro Gly His			
	170	175	180
Lys Gln Leu Asp His His Ser Lys Asp Thr Thr Leu Ile His Gln			
	185	190	195
Ile Phe Gly Ala Tyr Trp Arg Ser Gln Ile Lys Tyr Leu His Cys			
	200	205	210
His Gly Ile Ser Asp Thr Phe Asp Pro Tyr Leu Asp Ile Ala Leu			
	215	220	225
Asp Ile Gln Ala Ala Gln Ser Val Lys Gln Ala Leu Glu Gln Leu			
	230	235	240
Val Lys Pro Lys Glu Leu Asn Gly Glu Asn Ala Tyr His Cys Gly			
	245	250	255
Leu Cys Leu Gln Lys Ala Pro Ala Ser Lys Thr Leu Thr Leu Pro			
	260	265	270
Thr Ser Ala Lys Val Leu Ile Leu Val Leu Lys Arg Phe Ser Asp			
	275	280	285
Val Thr Gly Asn Lys Leu Ala Lys Asn Val Gln Tyr Pro Lys Cys			
	290	295	300
Arg Asp Met Gln Pro Tyr Met Ser Gln Gln Asn Thr Gly Pro Leu			
	305	310	315
Val Tyr Val Leu Tyr Ala Val Leu Val His Ala Gly Trp Ser Cys			
	320	325	330
His Asn Gly His Tyr Phe Ser Tyr Val Lys Ala Gln Glu Gly Gln			
	335	340	345
Trp Tyr Lys Met Asp Asp Ala Glu Val Thr Ala Ser Gly Ile Thr			
	350	355	360
Ser Val Leu Ser Gln Gln Ala Tyr Val Leu Phe Tyr Ile Gln Lys			
	365	370	375
Ser Glu Trp Glu Arg His Ser Glu Ser Val Ser Arg Gly Arg Glu			
	380	385	390
Pro Arg Ala Leu Gly Ala Glu Asp Thr Asp Arg Pro Ala Thr Gln			
	395	400	405
Gly Glu Leu Lys Arg Asp His Pro Cys Leu Gln Val Pro Glu Leu			
	410	415	420
Asp Glu His Leu Val Glu Arg Ala Thr Gln Glu Ser Thr Leu Asp			
	425	430	435
His Trp Lys Phe Pro Gln Lys Gln Asn Lys Thr Lys Pro Glu Phe			
	440	445	450
Asn Val Arg Lys Val Glu Gly Thr Leu Pro Pro Asn Val Leu Val			



	455		460		465
Ile His Gln Ser Lys Tyr Lys Cys Gly		Met Lys Asn His His Pro			
	470		475		480
Glu Gln Gln Ser Ser Leu Leu Asn Leu		Ser Ser Thr Lys Pro Thr			
	485		490		495
Asp Gln Glu Ser Met Asn Thr Gly Thr		Leu Ala Ser Leu Gln Gly			
	500		505		510
Ser Thr Arg Arg Ser Lys Gly Asn Asn		Lys His Ser Lys Arg Ser			
	515		520		525
Leu Leu Val Cys Gln					
	530				

<210> 12  
 <211> 511  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7500748CD1

<400> 12

Met Ala Ala Ala Met Pro Leu Ala Leu Leu Val Leu Leu Leu Leu			
1	5	10	15
Gly Pro Gly Gly Trp Cys Leu Ala Glu Pro Pro Arg Asp Ser Leu			
	20	25	30
Arg Glu Glu Leu Val Ile Thr Pro Leu Pro Ser Gly Asp Val Ala			
	35	40	45
Ala Thr Phe Gln Phe Arg Thr Arg Trp Asp Ser Glu Leu Gln Arg			
	50	55	60
Glu Gly Val Ser His Tyr Arg Leu Phe Pro Lys Ala Leu Gly Gln			
	65	70	75
Leu Ile Ser Lys Tyr Ser Leu Arg Glu Leu His Leu Ser Phe Thr			
	80	85	90
Gln Gly Phe Trp Arg Thr Arg Tyr Trp Gly Pro Pro Phe Leu Gln			
	95	100	105
Ala Pro Ser Gly Ala Glu Leu Trp Val Trp Phe Gln Asp Thr Val			
	110	115	120
Thr Asp Val Asp Lys Ser Trp Lys Glu Leu Ser Asn Val Leu Ser			
	125	130	135
Gly Ile Phe Cys Ala Ser Leu Asn Phe Ile Asp Ser Thr Asn Thr			
	140	145	150
Val Thr Pro Thr Ala Ser Phe Lys Pro Leu Gly Leu Ala Asn Asp			
	155	160	165
Thr Asp His Tyr Phe Leu Arg Tyr Ala Val Leu Pro Arg Glu Val			
	170	175	180
Val Cys Thr Glu Asn Leu Thr Pro Trp Lys Lys Leu Leu Pro Cys			
	185	190	195
Ser Ser Lys Ala Gly Leu Ser Val Leu Leu Lys Ala Asp Arg Leu			
	200	205	210
Phe His Thr Ser Tyr His Ser Gln Ala Val His Ile Arg Pro Val			
	215	220	225
Cys Arg Asn Ala Arg Cys Thr Ser Ile Ser Trp Glu Leu Arg Gln			
	230	235	240
Thr Leu Ser Val Val Phe Asp Ala Phe Ile Thr Gly Gln Gly Lys			
	245	250	255

Lys Asp Trp Ser Leu Phe Arg Met Phe Ser Arg Thr Leu Thr Glu  
 260 265 270  
 Pro Cys Pro Leu Ala Ser Glu Ser Arg Val Tyr Val Asp Ile Thr  
 275 280 285  
 Thr Tyr Asn Gln Asp Asn Glu Thr Leu Glu Val His Pro Pro Pro  
 290 295 300  
 Thr Thr Thr Tyr Gln Asp Val Ile Leu Gly Thr Arg Lys Thr Tyr  
 305 310 315  
 Ala Ile Tyr Asp Leu Leu Asp Thr Ala Met Ile Asn Asn Ser Arg  
 320 325 330  
 Asn Leu Asn Ile Gln Leu Lys Trp Lys Arg Pro Pro Glu Asn Gly  
 335 340 345  
 Tyr Ile His Tyr Gln Pro Ala Gln Asp Arg Leu Gln Pro His Leu  
 350 355 360  
 Leu Glu Met Leu Ile Gln Leu Pro Ala Asn Ser Val Thr Lys Val  
 365 370 375  
 Ser Ile Gln Phe Glu Arg Ala Leu Leu Lys Trp Thr Glu Tyr Thr  
 380 385 390  
 Pro Asp Pro Asn His Gly Phe Tyr Val Ser Pro Ser Val Leu Ser  
 395 400 405  
 Ala Leu Val Pro Ser Met Val Ala Ala Lys Pro Val Asp Trp Glu  
 410 415 420  
 Glu Ser Pro Leu Phe Asn Ser Leu Phe Pro Val Ser Asp Gly Ser  
 425 430 435  
 Asn Tyr Phe Val Arg Leu Tyr Thr Glu Pro Leu Leu Val Asn Leu  
 440 445 450  
 Pro Thr Pro Asp Phe Ser Met Pro Tyr Asn Val Ile Cys Leu Thr  
 455 460 465  
 Cys Thr Val Val Ala Val Cys Tyr Gly Ser Phe Tyr Asn Leu Leu  
 470 475 480  
 Thr Arg Thr Phe His Ile Glu Glu Pro Arg Thr Gly Gly Leu Ala  
 485 490 495  
 Lys Arg Leu Ala Asn Leu Ile Arg Arg Ala Arg Gly Val Pro Pro  
 500 505 510  
 Leu

&lt;210&gt; 13

&lt;211&gt; 476

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500749CD1

&lt;400&gt; 13

Met Ala Ala Ala Met Pro Leu Ala Leu Leu Val Leu Leu Leu Leu  
 1 5 10 15  
 Gly Pro Gly Gly Trp Cys Leu Ala Glu Pro Pro Arg Asp Ser Leu  
 20 25 30  
 Arg Glu Glu Leu Val Ile Thr Pro Leu Pro Ser Gly Asp Val Ala  
 35 40 45  
 Ala Thr Phe Gln Phe Arg Thr Arg Trp Asp Ser Glu Leu Gln Arg  
 50 55 60  
 Glu Gly Asp Thr Asp His Tyr Phe Leu Arg Tyr Ala Val Leu Pro

	65		70		75
Arg Glu Val Val Cys Thr Glu Asn Leu Thr Pro Trp Lys Lys Leu					
	80		85		90
Leu Pro Cys Ser Ser Lys Ala Gly Leu Ser Val Leu Leu Lys Ala					
	95		100		105
Asp Arg Leu Phe His Thr Ser Tyr His Ser Gln Ala Val His Ile					
	110		115		120
Arg Pro Val Cys Arg Asn Ala Arg Cys Thr Ser Ile Ser Trp Glu					
	125		130		135
Leu Arg Gln Thr Leu Ser Val Val Phe Asp Ala Phe Ile Thr Gly					
	140		145		150
Gln Gly Lys Lys Asp Trp Ser Leu Phe Arg Met Phe Ser Arg Thr					
	155		160		165
Leu Thr Glu Pro Cys Pro Leu Ala Ser Glu Ser Arg Val Tyr Val					
	170		175		180
Asp Ile Thr Thr Tyr Asn Gln Asp Asn Glu Thr Leu Glu Val His					
	185		190		195
Pro Pro Pro Thr Thr Thr Tyr Gln Asp Val Ile Leu Gly Thr Arg					
	200		205		210
Lys Thr Tyr Ala Ile Tyr Asp Leu Leu Asp Thr Ala Met Ile Asn					
	215		220		225
Asn Ser Arg Asn Leu Asn Ile Gln Leu Lys Trp Lys Arg Pro Pro					
	230		235		240
Glu Asn Glu Ala Pro Pro Val Pro Phe Leu His Ala Gln Arg Tyr					
	245		250		255
Val Ser Gly Tyr Gly Leu Gln Lys Gly Glu Leu Ser Thr Leu Leu					
	260		265		270
Tyr Asn Thr His Pro Tyr Arg Ala Phe Pro Val Leu Leu Leu Asp					
	275		280		285
Thr Val Pro Trp Tyr Leu Arg Leu Tyr Val His Thr Leu Thr Ile					
	290		295		300
Thr Ser Lys Gly Lys Glu Asn Lys Pro Ser Tyr Ile His Tyr Gln					
	305		310		315
Pro Ala Gln Asp Arg Leu Gln Pro His Leu Leu Glu Met Leu Ile					
	320		325		330
Gln Leu Pro Ala Asn Ser Val Thr Lys Val Ser Ile Gln Phe Glu					
	335		340		345
Arg Ala Leu Leu Lys Trp Thr Glu Tyr Thr Pro Asp Pro Asn His					
	350		355		360
Gly Phe Tyr Val Ser Pro Ser Val Leu Ser Ala Leu Val Pro Ser					
	365		370		375
Met Val Ala Ala Lys Pro Val Asp Trp Glu Glu Ser Pro Leu Phe					
	380		385		390
Asn Ser Leu Phe Pro Val Ser Asp Gly Ser Asn Tyr Phe Val Arg					
	395		400		405
Leu Tyr Thr Glu Pro Leu Leu Val Asn Leu Pro Thr Pro Asp Phe					
	410		415		420
Ser Met Pro Tyr Asn Val Ile Cys Leu Thr Cys Thr Val Val Ala					
	425		430		435
Val Cys Tyr Gly Ser Phe Tyr Asn Leu Leu Thr Arg Thr Phe His					
	440		445		450
Ile Glu Glu Pro Arg Thr Gly Gly Leu Ala Lys Arg Leu Ala Asn					
	455		460		465
Leu Ile Arg Arg Ala Arg Gly Val Pro Pro Leu					
	470		475		

<210> 14  
 <211> 344  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7503401CD1

<400> 14  
 Met Ala Ala Thr Glu Gly Val Gly Glu Ala Ala Gln Gly Gly Glu  
 1 5 10 15  
 Pro Gly Gln Pro Ala Gln Pro Pro Pro Gln Pro His Pro Pro Pro  
 20 25 30  
 Pro Gln Gln Gln His Lys Glu Glu Met Ala Ala Glu Ala Gly Glu  
 35 40 45  
 Ala Val Ala Ser Pro Met Asp Asp Gly Phe Val Ser Leu Asp Ser  
 50 55 60  
 Pro Ser Tyr Val Leu Tyr Arg Asp Arg Ala Glu Trp Ala Asp Ile  
 65 70 75  
 Asp Pro Val Pro Gln Asn Asp Gly Pro Asn Pro Val Val Gln Ile  
 80 85 90  
 Ile Tyr Ser Asp Lys Phe Arg Asp Val Tyr Asp Tyr Phe Arg Ala  
 95 100 105  
 Val Leu Gln Arg Asp Glu Arg Ser Glu Arg Ala Phe Lys Leu Thr  
 110 115 120  
 Arg Asp Ala Ile Glu Leu Asn Ala Ala Asn Tyr Thr Val Trp His  
 125 130 135  
 His Arg Arg Val Leu Val Glu Trp Leu Arg Asp Pro Ser Gln Glu  
 140 145 150  
 Leu Glu Phe Ile Ala Asp Ile Leu Asn Gln Asp Ala Lys Asn Tyr  
 155 160 165  
 His Ala Trp Gln His Arg Gln Trp Val Ile Gln Glu Phe Lys Leu  
 170 175 180  
 Trp Asp Asn Glu Leu Gln Tyr Val Asp Gln Leu Leu Lys Glu Asp  
 185 190 195  
 Val Arg Asn Asn Ser Val Trp Asn Gln Arg Tyr Phe Val Ile Ser  
 200 205 210  
 Asn Thr Thr Gly Tyr Asn Asp Arg Ala Val Leu Glu Arg Glu Val  
 215 220 225  
 Gln Tyr Thr Leu Glu Met Ile Lys Leu Val Pro His Asn Glu Ser  
 230 235 240  
 Ala Trp Asn Tyr Leu Lys Gly Ile Leu Gln Asp Arg Gly Leu Ser  
 245 250 255  
 Lys Tyr Pro Asn Leu Leu Asn Gln Leu Leu Asp Leu Gln Pro Ser  
 260 265 270  
 His Ser Ser Pro Tyr Leu Ile Ala Phe Leu Val Asp Ile Tyr Glu  
 275 280 285  
 Asp Met Leu Glu Asn Gln Cys Asp Asn Lys Glu Asp Ile Leu Asn  
 290 295 300  
 Lys Ala Leu Glu Leu Cys Glu Ile Leu Ala Lys Glu Lys Asp Thr  
 305 310 315  
 Ile Arg Lys Glu Tyr Trp Arg Tyr Ile Gly Arg Ser Leu Gln Ser  
 320 325 330  
 Lys His Ser Thr Glu Asn Asp Ser Pro Thr Asn Val Gln Gln  
 335 340

<210> 15  
 <211> 122  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7503485CD1

<400> 15  
 Met Ser Thr Pro Ala Arg Arg Arg Leu Met Arg Asp Phe Lys Arg  
     1                    5                    10                    15  
 Leu Gln Glu Asp Pro Pro Ala Gly Val Ser Gly Ala Pro Ser Glu  
                     20                    25                    30  
 Asn Asn Ile Met Val Trp Asn Ala Val Ile Phe Gly Pro Glu Gly  
                     35                    40                    45  
 Thr Pro Phe Glu Asp Val Tyr Ala Asp Gly Ser Ile Cys Leu Asp  
                     50                    55                    60  
 Ile Leu Gln Asn Arg Trp Ser Pro Thr Tyr Asp Val Ser Ser Ile  
                     65                    70                    75  
 Leu Thr Ser Ile Gln Ser Leu Leu Asp Glu Pro Asn Pro Asn Ser  
                     80                    85                    90  
 Pro Ala Asn Ser Gln Ala Ala Gln Leu Tyr Gln Glu Asn Lys Arg  
                     95                    100                    105  
 Glu Tyr Glu Lys Arg Val Ser Ala Ile Val Glu Gln Ser Trp Arg  
                     110                    115                    120  
 Asp Cys

<210> 16  
 <211> 255  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7504076CD1

<400> 16  
 Met Ala Val Gly Asn Ile Asn Glu Leu Pro Glu Asn Ile Leu Leu  
     1                    5                    10                    15  
 Glu Leu Phe Thr His Val Pro Ala Arg Gln Leu Leu Leu Asn Cys  
                     20                    25                    30  
 Arg Leu Val Cys Ser Leu Trp Arg Asp Leu Ile Asp Leu Val Thr  
                     35                    40                    45  
 Leu Trp Lys Arg Lys Cys Leu Arg Glu Gly Phe Ile Thr Glu Asp  
                     50                    55                    60  
 Trp Asp Gln Pro Val Ala Asp Trp Lys Ile Phe Tyr Phe Leu Arg  
                     65                    70                    75  
 Ser Leu His Arg Asn Leu Leu His Asn Pro Cys Ala Glu Glu Gly  
                     80                    85                    90  
 Phe Glu Phe Trp Ser Leu Asp Val Asn Gly Gly Asp Glu Trp Lys  
                     95                    100                    105  
 Val Glu Asp Leu Ser Arg Asp Gln Arg Lys Glu Phe Pro Asn Asp  
                     110                    115                    120  
 Gln Val Lys Lys Tyr Phe Val Thr Ser Tyr Tyr Thr Cys Leu Lys

	125	130	135
Ser Gln Val Val	Asp Leu Lys Ala Glu Gly Tyr Trp Glu Glu Leu		
	140	145	150
Met Asp Thr Thr	Arg Pro Asp Ile Glu Val Lys Asp Trp Phe Ala		
	155	160	165
Ala Arg Pro Asp	Cys Gly Ser Lys Tyr Gln Leu Cys Val Gln Leu		
	170	175	180
Leu Ser Ser Ala	His Ala Pro Leu Gly Thr Phe Gln Pro Asp Pro		
	185	190	195
Ala Thr Ile Gln	Gln Lys Ser Asp Ala Lys Trp Arg Glu Val Ser		
	200	205	210
His Thr Phe Ser	Asn Tyr Pro Pro Gly Val Arg Tyr Ile Trp Phe		
	215	220	225
Gln His Gly Gly	Val Asp Thr His Tyr Trp Ala Gly Trp Tyr Gly		
	230	235	240
Pro Arg Val Thr	Asn Ser Ser Ile Thr Ile Gly Pro Pro Leu Pro		
	245	250	255

&lt;210&gt; 17

&lt;211&gt; 166

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500926CD1

&lt;400&gt; 17

Met Ser Ala Trp	Ala Ala Ala Ser Leu Ser Arg Ala Ala Ala Arg	
1	5	10
Cys Leu Leu Ala	Arg Gly Pro Gly Val Arg Ala Ala Pro Pro Arg	15
	20	25
Asp Pro Arg Pro	Ser His Pro Glu Pro Arg Gly Cys Gly Ala Ala	30
	35	40
Pro Gly Arg Thr	Leu His Phe Thr Ala Ala Val Pro Ala Gly His	45
	50	55
Asn Lys Trp Ser	Lys Val Arg His Ile Lys Gly Pro Lys Asp Val	60
	65	70
Glu Arg Ser Arg	Ile Phe Ser Lys Leu Cys Leu Asn Ile Arg Leu	75
	80	85
Ala Val Lys Glu	Gly Gly Pro Asn Pro Glu His Asn Ser Asn Leu	90
	95	100
Ala Asn Ile Leu	Glu Val Cys Arg Ser Lys His Met Pro Lys Ser	105
	110	115
Thr Ile Glu Thr	Ala Leu Lys Met Glu Lys Ser Lys Asp Thr Tyr	120
	125	130
Leu Leu Tyr Glu	Gly Arg Gly Pro Gly Gly Ser Ser Leu Leu Ile	135
	140	145
Glu Ala Leu Ser	Asn Ser Ser His Lys Cys Gln Ala Asp Leu Arg	150
	155	160
Pro		165

&lt;210&gt; 18

&lt;211&gt; 591

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503216CD1

&lt;400&gt; 18

Met	Pro	Pro	Lys	Val	Thr	Ser	Glu	Leu	Leu	Arg	Gln	Leu	Arg	Gln			
1				5					10					15			
Ala	Met	Arg	Asn	Ser	Glu	Tyr	Val	Thr	Glu	Pro	Ile	Gln	Ala	Tyr			
			20						25					30			
Ile	Ile	Pro	Ser	Gly	Asp	Ala	His	Gln	Ser	Glu	Tyr	Ile	Ala	Pro			
			35						40					45			
Cys	Asp	Cys	Arg	Arg	Ala	Phe	Val	Ser	Gly	Phe	Asp	Gly	Ser	Ala			
			50						55					60			
Gly	Thr	Ala	Ile	Ile	Thr	Glu	Glu	His	Ala	Ala	Met	Trp	Thr	Asp			
			65						70					75			
Gly	Arg	Tyr	Phe	Leu	Gln	Ala	Ala	Lys	Gln	Met	Asp	Ser	Asn	Trp			
			80						85					90			
Thr	Leu	Met	Lys	Met	Gly	Leu	Lys	Asp	Thr	Pro	Thr	Gln	Glu	Asp			
			95						100					105			
Trp	Leu	Val	Ser	Val	Leu	Pro	Glu	Gly	Ser	Arg	Val	Gly	Val	Asp			
			110						115					120			
Pro	Leu	Ile	Ile	Pro	Thr	Asp	Tyr	Trp	Lys	Lys	Met	Ala	Lys	Val			
			125						130					135			
Leu	Arg	Ser	Ala	Gly	His	His	Leu	Ile	Pro	Val	Lys	Glu	Asn	Leu			
			140						145					150			
Val	Asp	Lys	Ile	Trp	Thr	Asp	Arg	Pro	Glu	Arg	Pro	Cys	Lys	Pro			
			155						160					165			
Leu	Leu	Thr	Leu	Gly	Leu	Asp	Tyr	Thr	Gly	Leu	Phe	Asn	Leu	Arg			
			170						175					180			
Gly	Ser	Asp	Val	Glu	His	Asn	Pro	Val	Phe	Phe	Ser	Tyr	Ala	Ile			
			185						190					195			
Ile	Gly	Leu	Glu	Thr	Ile	Met	Leu	Phe	Ile	Asp	Gly	Asp	Arg	Ile			
			200						205					210			
Asp	Ala	Pro	Ser	Val	Lys	Glu	His	Leu	Leu	Leu	Asp	Leu	Gly	Leu			
			215						220					225			
Glu	Ala	Glu	Tyr	Arg	Ile	Gln	Val	His	Pro	Tyr	Lys	Ser	Ile	Leu			
			230						235					240			
Ser	Glu	Leu	Lys	Ala	Leu	Cys	Ala	Asp	Leu	Ser	Pro	Arg	Glu	Lys			
			245						250					255			
Val	Trp	Val	Ser	Asp	Lys	Ala	Ser	Tyr	Ala	Val	Ser	Glu	Thr	Ile			
			260						265					270			
Pro	Lys	Asp	His	Arg	Cys	Cys	Met	Pro	Tyr	Thr	Pro	Ile	Cys	Ile			
			275						280					285			
Ala	Lys	Ala	Val	Lys	Asn	Ser	Ala	Glu	Ser	Glu	Gly	Met	Arg	Arg			
			290						295					300			
Ala	His	Ile	Lys	Asp	Ala	Val	Ala	Leu	Cys	Glu	Leu	Phe	Asn	Trp			
			305						310					315			
Leu	Glu	Lys	Glu	Val	Pro	Lys	Gly	Gly	Val	Thr	Glu	Ile	Ser	Ala			
			320						325					330			
Ala	Asp	Lys	Ala	Glu	Glu	Phe	Arg	Arg	Gln	Gln	Ala	Asp	Phe	Val			
			335						340					345			
Asp	Leu	Ser	Phe	Pro	Thr	Ile	Ser	Ser	Thr	Gly	Pro	Asn	Gly	Ala			
			350						355					360			

```

Ile Ile His Tyr Ala Pro Val Pro Glu Thr Asn Arg Thr Leu Ser
365 370 375
Leu Asp Glu Val Tyr Leu Ile Asp Ser Gly Ala Gln Tyr Lys Asp
380 385 390
Gly Thr Thr Asp Val Thr Arg Thr Met His Phe Gly Thr Pro Thr
395 400 405
Ala Tyr Glu Lys Glu Cys Phe Thr Tyr Val Leu Lys Gly His Ile
410 415 420
Ala Val Ser Ala Ala Val Phe Pro Thr Gly Thr Lys Gly His Leu
425 430 435
Leu Asp Ser Phe Ala Arg Ser Ala Leu Trp Asp Ser Gly Leu Asp
440 445 450
Tyr Leu His Gly Thr Gly His Gly Val Gly Ser Phe Leu Asn Val
455 460 465
His Glu Gly Pro Cys Gly Ile Ser Tyr Lys Thr Phe Ser Asp Glu
470 475 480
Pro Leu Glu Ala Gly Met Ile Val Thr Asp Glu Pro Gly Tyr Tyr
485 490 495
Glu Asp Gly Ala Phe Gly Ile Arg Ile Glu Asn Val Val Leu Val
500 505 510
Val Pro Val Lys Thr Lys Tyr Asn Phe Asn Asn Arg Gly Ser Leu
515 520 525
Thr Phe Glu Pro Leu Thr Leu Val Pro Ile Gln Thr Lys Met Ile
530 535 540
Asp Val Asp Ser Leu Thr Asp Lys Glu Cys Asp Trp Leu Asn Asn
545 550 555
Tyr His Leu Thr Cys Arg Asp Val Ile Gly Lys Glu Leu Gln Lys
560 565 570
Gln Gly Arg Gln Glu Ala Leu Glu Trp Leu Ile Arg Glu Thr Gln
575 580 585
Pro Ile Ser Lys Gln His
590

```

&lt;210&gt; 19

&lt;211&gt; 652

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503233CD1

&lt;400&gt; 19

```

Met Ser Glu Glu Ile Ile Thr Pro Val Tyr Cys Thr Gly Val Ser
1 5 10 15
Ala Gln Val Gln Lys Gln Arg Ala Arg Glu Leu Gly Leu Gly Arg
20 25 30
His Glu Asn Ala Ile Lys Tyr Leu Gly Gln Asp Tyr Glu Gln Leu
35 40 45
Arg Val Arg Cys Leu Gln Ser Gly Thr Leu Phe Arg Asp Glu Ala
50 55 60
Phe Pro Pro Val Pro Gln Ser Leu Gly Tyr Lys Asp Leu Gly Pro
65 70 75
Asn Ser Ser Lys Thr Tyr Gly Tyr Ala Gly Ile Phe His Phe Gln
80 85 90
Leu Trp Gln Phe Gly Glu Trp Val Asp Val Val Val Asp Asp Leu

```



	95		100		105
Leu Pro Ile Lys Asp Gly Lys Leu Val Phe Val His Ser Ala Glu					
	110		115		120
Gly Asn Glu Phe Trp Ser Ala Leu Leu Glu Lys Ala Tyr Ala Lys					
	125		130		135
Val Asn Gly Ser Tyr Glu Ala Leu Ser Gly Gly Ser Thr Ser Glu					
	140		145		150
Gly Phe Glu Asp Phe Thr Gly Gly Val Thr Glu Trp Tyr Glu Leu					
	155		160		165
Arg Lys Ala Pro Ser Asp Leu Tyr Gln Ile Ile Leu Lys Ala Leu					
	170		175		180
Glu Arg Gly Ser Leu Leu Gly Cys Ser Ile Asp Ile Ser Ser Val					
	185		190		195
Leu Asp Met Glu Ala Ile Thr Phe Lys Lys Leu Val Lys Gly His					
	200		205		210
Ala Tyr Ser Val Thr Gly Ala Lys Gln Val Asn Tyr Arg Gly Gln					
	215		220		225
Val Val Ser Leu Ile Arg Met Arg Asn Pro Trp Gly Glu Val Glu					
	230		235		240
Trp Thr Gly Ala Trp Ser Asp Ser Ser Ser Glu Trp Asn Asn Val					
	245		250		255
Asp Pro Tyr Glu Arg Asp Gln Leu Arg Val Lys Met Glu Asp Gly					
	260		265		270
Glu Phe Trp Met Ser Phe Arg Asp Phe Met Arg Glu Phe Thr Arg					
	275		280		285
Leu Glu Ile Cys Asn Leu Thr Pro Asp Ala Leu Lys Ser Arg Thr					
	290		295		300
Ile Arg Lys Trp Asn Thr Thr Leu Tyr Glu Gly Thr Trp Arg Arg					
	305		310		315
Gly Ser Thr Ala Gly Gly Cys Arg Asn Tyr Pro Ala Thr Phe Trp					
	320		325		330
Val Asn Pro Gln Phe Lys Ile Arg Leu Asp Glu Thr Asp Asp Pro					
	335		340		345
Asp Asp Tyr Gly Asp Arg Glu Ser Gly Cys Ser Phe Val Leu Ala					
	350		355		360
Leu Met Gln Lys His Arg Arg Arg Glu Arg Arg Phe Gly Arg Asp					
	365		370		375
Met Glu Thr Ile Gly Phe Ala Val Tyr Glu Val Pro Pro Glu Leu					
	380		385		390
Val Gly Gln Pro Ala Val His Leu Lys Arg Asp Phe Phe Leu Ala					
	395		400		405
Asn Ala Ser Arg Ala Arg Ser Glu Gln Phe Ile Asn Leu Arg Glu					
	410		415		420
Val Ser Thr Arg Phe Arg Leu Pro Pro Gly Glu Tyr Val Val Val					
	425		430		435
Pro Ser Thr Phe Glu Pro Asn Lys Glu Gly Asp Phe Val Leu Arg					
	440		445		450
Phe Phe Ser Glu Lys Ser Ala Gly Thr Val Glu Leu Asp Asp Gln					
	455		460		465
Ile Gln Ala Asn Leu Pro Asp Glu Gln Val Leu Ser Glu Glu Glu					
	470		475		480
Ile Asp Glu Asn Phe Lys Ala Leu Phe Arg Gln Leu Ala Gly Glu					
	485		490		495
Asp Met Glu Ile Ser Val Lys Glu Leu Arg Thr Ile Leu Asn Arg					
	500		505		510
Ile Ile Ser Lys His Lys Asp Leu Arg Thr Lys Gly Phe Ser Leu					

515	520	525
Glu Ser Cys Arg Ser Met Val Asn Leu Met Asp Arg Asp Gly Asn		
530	535	540
Gly Lys Leu Gly Leu Val Glu Phe Asn Ile Leu Trp Asn Arg Ile		
545	550	555
Arg Asn Tyr Leu Ser Ile Phe Arg Lys Phe Asp Leu Asp Lys Ser		
560	565	570
Gly Ser Met Ser Ala Tyr Glu Met Arg Met Ala Ile Glu Ser Ala		
575	580	585
Gly Phe Lys Leu Asn Lys Lys Leu Tyr Glu Leu Ile Ile Thr Arg		
590	595	600
Tyr Ser Glu Pro Asp Leu Ala Val Asp Phe Asp Asn Phe Val Cys		
605	610	615
Cys Leu Val Arg Leu Glu Thr Met Phe Arg Phe Phe Lys Thr Leu		
620	625	630
Asp Thr Asp Leu Asp Gly Val Val Thr Phe Asp Leu Phe Lys Trp		
635	640	645
Leu Gln Leu Thr Met Phe Ala		
650		

&lt;210&gt; 20

&lt;211&gt; 861

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7726576CD1

&lt;400&gt; 20

Met Ala Gly Pro Gly Pro Gly Ala Val Leu Glu Ser Pro Arg Gln		
1	5	10
Leu Leu Gly Arg Val Arg Phe Leu Ala Glu Ala Ala Arg Ser Leu		
20	25	30
Arg Ala Gly Arg Pro Leu Pro Ala Ala Leu Ala Phe Val Pro Arg		
35	40	45
Glu Val Leu Tyr Lys Leu Tyr Lys Asp Pro Ala Gly Pro Ser Arg		
50	55	60
Val Leu Leu Pro Val Trp Glu Ala Glu Gly Leu Gly Leu Arg Val		
65	70	75
Gly Ala Ala Gly Pro Ala Pro Gly Thr Gly Ser Gly Pro Leu Arg		
80	85	90
Ala Ala Arg Asp Ser Ile Glu Leu Arg Arg Gly Ala Cys Val Arg		
95	100	105
Thr Thr Gly Glu Glu Leu Cys Asn Gly His Gly Leu Trp Val Lys		
110	115	120
Leu Thr Lys Glu Gln Leu Ala Glu His Leu Gly Asp Cys Gly Leu		
125	130	135
Gln Glu Gly Trp Leu Leu Val Cys Arg Pro Ala Glu Gly Gly Ala		
140	145	150
Arg Leu Val Pro Ile Asp Thr Pro Asn His Leu Gln Arg Gln Gln		
155	160	165
Gln Leu Phe Gly Val Asp Tyr Arg Pro Val Leu Arg Trp Glu Gln		
170	175	180
Val Val Asp Leu Thr Tyr Ser His Arg Leu Gly Ser Arg Pro Gln		
185	190	195

Pro Ala Glu Ala Tyr Ala Glu Ala Val	Gln Arg Leu Leu Tyr Val
200	205 210
Pro Pro Thr Trp Thr Tyr Glu Cys Asp	Glu Asp Leu Ile His Phe
215	220 225
Leu Tyr Asp His Leu Gly Lys Glu Asp	Glu Asn Leu Gly Ser Val
230	235 240
Lys Gln Tyr Val Glu Ser Ile Asp Val	Ser Ser Tyr Thr Glu Glu
245	250 255
Phe Asn Val Ser Cys Leu Thr Asp Ser	Asn Ala Asp Thr Tyr Trp
260	265 270
Glu Ser Asp Gly Ser Gln Cys Gln His	Trp Val Arg Leu Thr Met
275	280 285
Lys Lys Gly Thr Ile Val Lys Lys Leu	Leu Leu Thr Val Asp Thr
290	295 300
Thr Asp Asp Asn Phe Met Pro Lys Arg	Val Val Val Tyr Gly Gly
305	310 315
Glu Gly Asp Asn Leu Lys Lys Leu Ser	Asp Val Ser Ile Asp Glu
320	325 330
Thr Leu Ile Gly Asp Val Cys Val Leu	Glu Asp Met Thr Val His
335	340 345
Leu Pro Ile Ile Glu Ile Arg Ile Val	Glu Cys Arg Asp Asp Gly
350	355 360
Ile Asp Val Arg Leu Arg Gly Val Lys	Ile Lys Ser Ser Arg Gln
365	370 375
Arg Glu Leu Gly Leu Asn Ala Asp Leu	Phe Gln Pro Thr Ser Leu
380	385 390
Val Arg Tyr Pro Arg Leu Glu Gly Thr	Asp Pro Glu Val Leu Tyr
395	400 405
Arg Arg Ala Val Leu Leu Gln Arg Leu	Ile Lys Ile Leu Asp Ser
410	415 420
Val Leu His His Leu Val Pro Ala Trp	Asp His Thr Leu Gly Thr
425	430 435
Phe Ser Glu Ile Lys Gln Val Lys Gln	Phe Leu Leu Leu Ser Arg
440	445 450
Gln Arg Pro Gly Leu Val Ala Gln Cys	Leu Arg Asp Ser Glu Ser
455	460 465
Ser Lys Pro Ser Phe Met Pro Arg Leu	Tyr Ile Asn Arg Arg Leu
470	475 480
Ala Met Glu His Arg Ala Cys Pro Ser	Arg Asp Pro Ala Cys Lys
485	490 495
Asn Ala Val Phe Thr Gln Val Tyr Glu	Gly Leu Lys Pro Ser Asp
500	505 510
Lys Tyr Glu Lys Pro Leu Asp Tyr Arg	Trp Pro Met Arg Tyr Asp
515	520 525
Gln Trp Trp Glu Cys Lys Phe Ile Ala	Glu Gly Ile Ile Asp Gln
530	535 540
Gly Gly Gly Phe Arg Asp Ser Leu Ala	Asp Met Ser Glu Glu Leu
545	550 555
Cys Pro Ser Ser Ala Asp Thr Pro Val	Pro Leu Pro Phe Phe Val
560	565 570
Arg Thr Ala Asn Gln Gly Asn Gly Thr	Gly Glu Ala Arg Asp Met
575	580 585
Tyr Val Pro Asn Pro Ser Cys Arg Asp	Phe Ala Lys Tyr Glu Trp
590	595 600
Ile Gly Gln Leu Met Gly Ala Ala Leu	Arg Gly Lys Glu Phe Leu
605	610 615

```

Val Leu Ala Leu Pro Gly Phe Val Trp Lys Gln Leu Ser Gly Glu
      620                      625                      630
Glu Val Ser Trp Ser Lys Asp Phe Pro Ala Val Asp Ser Val Leu
      635                      640                      645
Val Lys Leu Leu Glu Val Met Glu Gly Met Asp Lys Glu Thr Phe
      650                      655                      660
Glu Phe Lys Phe Gly Lys Glu Leu Thr Phe Thr Thr Val Leu Ser
      665                      670                      675
Asp Gln Gln Val Val Glu Leu Ile Pro Gly Gly Ala Gly Ile Val
      680                      685                      690
Val Gly Tyr Gly Asp Arg Ser Arg Phe Ile Gln Leu Val Gln Lys
      695                      700                      705
Ala Arg Leu Glu Glu Ser Lys Glu Gln Val Ala Ala Met Gln Ala
      710                      715                      720
Gly Leu Leu Lys Val Val Pro Gln Ala Val Leu Asp Leu Leu Thr
      725                      730                      735
Trp Gln Glu Leu Glu Lys Lys Val Cys Gly Asp Pro Glu Val Thr
      740                      745                      750
Val Asp Ala Leu Arg Lys Leu Thr Arg Phe Glu Asp Phe Glu Pro
      755                      760                      765
Ser Asp Ser Arg Val Gln Tyr Phe Trp Glu Ala Leu Asn Asn Phe
      770                      775                      780
Thr Asn Glu Asp Arg Ser Arg Val Leu Arg Phe Val Thr Gly Arg
      785                      790                      795
Ser Arg Leu Pro Ala Arg Ile Tyr Ile Tyr Pro Asp Lys Leu Gly
      800                      805                      810
Tyr Glu Thr Thr Asp Ala Leu Pro Glu Ser Ser Thr Cys Ser Ser
      815                      820                      825
Thr Leu Phe Leu Pro His Tyr Ala Ser Ala Lys Val Cys Glu Glu
      830                      835                      840
Lys Leu Arg Tyr Ala Ala Tyr Asn Cys Val Ala Ile Asp Thr Asp
      845                      850                      855
Met Ser Pro Trp Glu Glu
      860

```

&lt;210&gt; 21

&lt;211&gt; 447

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503507CD1

&lt;400&gt; 21

```

Met Ala Ser Val Ala Gln Glu Ser Ala Gly Ser Gln Arg Arg Leu
  1              5              10              15
Pro Pro Arg His Gly Ala Leu Arg Gly Leu Leu Leu Cys Leu
      20              25              30
Trp Leu Pro Ser Gly Arg Ala Ala Leu Pro Pro Ala Ala Pro Leu
      35              40              45
Ser Glu Leu His Ala Gln Leu Ser Gly Val Glu Gln Leu Leu Glu
      50              55              60
Glu Phe Arg Arg Gln Leu Gln Gln Glu Arg Pro Gln Glu Glu Leu
      65              70              75
Glu Leu Glu Leu Arg Ala Gly Gly Gly Pro Gln Glu Asp Cys Pro

```

	80		85		90
Gly Arg Gly Ser	Gly Gly Tyr Ser Ala Met Pro Asp Ala Ile Ile				
	95		100		105
Arg Thr Lys Asp	Ser Leu Ala Ala Gly Ala Ser Phe Leu Arg Ala				
	110		115		120
Pro Ala Ala Val	Arg Gly Trp Arg Gln Cys Val Ala Ala Cys Cys				
	125		130		135
Ser Glu Pro Arg	Cys Ser Val Ala Val Val Glu Leu Pro Arg Arg				
	140		145		150
Pro Ala Pro Pro	Ala Ala Val Leu Gly Cys Tyr Leu Phe Asn Cys				
	155		160		165
Thr Ala Arg Gly	Arg Asn Val Cys Lys Phe Ala Leu His Ser Gly				
	170		175		180
Tyr Ser Ser Tyr	Ser Leu Ser Arg Ala Pro Asp Gly Ala Ala Leu				
	185		190		195
Ala Thr Ala Arg	Ala Ser Pro Arg Gln Glu Lys Asp Ala Pro Pro				
	200		205		210
Leu Ser Lys Ala	Gly Gln Asp Val Val Leu His Leu Pro Thr Asp				
	215		220		225
Gly Val Val Leu	Asp Gly Arg Glu Ser Thr Asp Asp His Ala Ile				
	230		235		240
Val Gln Tyr Glu	Trp Ala Leu Leu Gln Gly Asp Pro Ser Val Asp				
	245		250		255
Met Lys Val Pro	Gln Ser Gly Thr Leu Lys Leu Ser His Leu Gln				
	260		265		270
Glu Gly Thr Tyr	Thr Phe Gln Leu Thr Val Thr Asp Thr Ala Gly				
	275		280		285
Gln Arg Ser Ser	Asp Asn Val Ser Val Thr Val Leu Arg Ala Ala				
	290		295		300
Tyr Ser Thr Gly	Gly Cys Leu His Thr Cys Ser Arg Tyr His Phe				
	305		310		315
Phe Cys Asp Asp	Gly Cys Cys Ile Asp Ile Thr Leu Ala Cys Asp				
	320		325		330
Gly Val Gln Gln	Cys Pro Asp Gly Ser Asp Glu Asp Phe Cys Gln				
	335		340		345
Asn Leu Gly Leu	Asp Arg Lys Met Val Thr His Thr Ala Ala Ser				
	350		355		360
Pro Ala Leu Pro	Arg Thr Thr Gly Pro Ser Glu Asp Ala Gly Gly				
	365		370		375
Asp Ser Leu Val	Glu Lys Ser Gln Lys Ala Thr Ala Pro Asn Lys				
	380		385		390
Pro Pro Ala Leu	Ser Asn Thr Glu Lys Arg Lys Val Ile Tyr Leu				
	395		400		405
Ser Gln Arg Val	Met Glu Glu Glu Gly Asn Thr Gln Pro Gln Lys				
	410		415		420
Gln Val Gln Cys	Tyr Pro Trp Arg Trp Val Trp Leu Ser Leu Leu				
	425		430		435
Cys Cys Phe Ser	Trp Leu His Ala Asp Tyr Asp Trp				
	440		445		

&lt;210&gt; 22

&lt;211&gt; 468

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503506CD1

&lt;400&gt; 22

```

Met Ala Ser Val Ala Gln Glu Ser Ala Gly Ser Gln Arg Arg Leu
 1           5           10           15
Pro Pro Arg His Gly Ala Leu Arg Gly Leu Leu Leu Leu Cys Leu
          20           25           30
Trp Leu Pro Ser Gly Arg Ala Ala Leu Pro Pro Ala Ala Pro Leu
          35           40           45
Ser Glu Leu His Ala Gln Leu Ser Gly Val Glu Gln Leu Leu Glu
          50           55           60
Glu Phe Arg Arg Gln Leu Gln Gln Glu Arg Pro Gln Glu Glu Leu
          65           70           75
Glu Leu Glu Leu Arg Ala Gly Gly Gly Pro Gln Glu Asp Cys Pro
          80           85           90
Gly Arg Gly Ser Gly Gly Tyr Ser Ala Met Pro Asp Ala Ile Ile
          95          100          105
Arg Thr Lys Asp Ser Leu Ala Ala Gly Ala Ser Phe Leu Arg Ala
          110          115          120
Pro Ala Ala Val Arg Gly Trp Arg Gln Cys Val Ala Ala Cys Cys
          125          130          135
Ser Glu Pro Arg Cys Ser Val Ala Val Val Glu Leu Pro Arg Arg
          140          145          150
Pro Ala Pro Pro Ala Ala Val Leu Gly Cys Tyr Leu Phe Asn Cys
          155          160          165
Thr Ala Arg Gly Arg Asn Val Cys Lys Phe Ala Leu His Ser Gly
          170          175          180
Tyr Ser Ser Tyr Ser Leu Ser Arg Ala Pro Asp Gly Ala Ala Leu
          185          190          195
Ala Thr Ala Arg Ala Ser Pro Arg Gln Glu Lys Asp Ala Pro Pro
          200          205          210
Leu Ser Lys Ala Gly Gln Asp Val Val Leu His Leu Pro Thr Asp
          215          220          225
Gly Val Val Leu Asp Gly Arg Glu Ser Thr Asp Asp His Ala Ile
          230          235          240
Val Gln Tyr Glu Trp Ala Leu Leu Gln Gly Asp Pro Ser Val Asp
          245          250          255
Met Lys Val Pro Gln Ser Gly Thr Leu Lys Leu Ser His Leu Gln
          260          265          270
Glu Gly Thr Tyr Thr Phe Gln Leu Thr Val Thr Asp Thr Ala Gly
          275          280          285
Gln Arg Ser Ser Asp Asn Val Ser Val Thr Val Leu Arg Ala Ala
          290          295          300
Tyr Ser Thr Gly Gly Cys Leu His Thr Cys Ser Arg Tyr His Phe
          305          310          315
Phe Cys Asp Asp Gly Cys Cys Ile Asp Ile Thr Leu Ala Cys Asp
          320          325          330
Gly Val Gln Gln Cys Pro Asp Gly Ser Asp Glu Asp Phe Cys Gln
          335          340          345
Asn Leu Gly Leu Asp Arg Lys Met Val Thr His Thr Ala Ala Ser
          350          355          360
Pro Ala Leu Pro Arg Thr Thr Gly Pro Ser Glu Asp Ala Gly Gly
          365          370          375
Asp Ser Leu Val Glu Lys Ser Gln Lys Ala Thr Ala Pro Asn Lys
          380          385          390

```

```

Pro Pro Ala Leu Ser Asn Thr Glu Lys Arg Asn His Ser Ala Phe
      395                      400                      405
Trp Gly Pro Glu Ser Gln Ile Ile Pro Val Met Pro Gly Ala Val
      410                      415                      420
Leu Pro Leu Ala Leu Gly Leu Ala Ile Thr Ala Leu Leu Leu Leu
      425                      430                      435
Met Val Ala Cys Arg Leu Arg Leu Val Lys Gln Lys Leu Lys Lys
      440                      445                      450
Ala Arg Pro Ile Thr Ser Glu Glu Ser Asp Tyr Leu Ile Asn Gly
      455                      460                      465
Met Tyr Leu

```

```

<210> 23
<211> 236
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 7503509CD1

```

```

<400> 23
Met Ala Ser Val Ala Gln Glu Ser Ala Gly Ser Gln Arg Arg Leu
  1          5          10          15
Pro Pro Arg His Gly Ala Leu Arg Gly Leu Leu Leu Leu Cys Leu
      20          25          30
Trp Leu Pro Ser Gly Arg Ala Ala Leu Pro Pro Ala Ala Pro Leu
      35          40          45
Ser Glu Leu His Ala Gln Leu Ser Gly Val Glu Gln Leu Leu Glu
      50          55          60
Glu Phe Arg Arg Gln Leu Gln Gln Glu Arg Pro Gln Glu Glu Leu
      65          70          75
Glu Leu Glu Leu Arg Ala Gly Gly Gly Pro Gln Glu Asp Cys Pro
      80          85          90
Gly Pro Gly Ser Gly Gly Tyr Ser Ala Met Pro Asp Ala Ile Ile
      95          100         105
Arg Thr Lys Asp Ser Leu Ala Ala Gly Ala Ser Phe Leu Arg Ala
     110         115         120
Pro Ala Ala Val Arg Gly Trp Arg Gln Cys Val Ala Ala Cys Cys
     125         130         135
Ser Glu Pro Arg Cys Ser Val Ala Val Val Glu Leu Pro Arg Arg
     140         145         150
Pro Ala Pro Pro Ala Ala Val Leu Gly Cys Tyr Leu Phe Asn Cys
     155         160         165
Thr Ala Arg Gly Arg Asn Val Cys Lys Phe Ala Leu His Ser Gly
     170         175         180
Tyr Ser Ser Tyr Ser Leu Ser Arg Ala Pro Asp Gly Ala Ala Leu
     185         190         195
Ala Thr Ala Arg Ala Ser Pro Arg Gln Gly Ala Ser Ile Arg Asn
     200         205         210
Pro Glu Ala Val Pro Pro Thr Gly Gly Asn Leu His Leu Pro Ala
     215         220         225
Asp Arg Asp Gly His Cys Arg Ala Glu Lys Leu
     230         235

```

<210> 24  
 <211> 312  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7505800CD1

<400> 24  
 Met Ala Ala Thr Glu Gly Val Gly Glu Ala Ala Gln Gly Gly Glu  
 1 5 10 15  
 Pro Gly Gln Pro Ala Gln Pro Pro Pro Gln Pro His Pro Pro Pro  
 20 25 30  
 Pro Gln Gln Gln His Lys Glu Glu Met Ala Ala Glu Ala Gly Glu  
 35 40 45  
 Ala Val Ala Ser Pro Met Asp Asp Gly Phe Val Ser Leu Asp Ser  
 50 55 60  
 Pro Ser Tyr Val Leu Tyr Arg His Phe Arg Arg Val Leu Leu Lys  
 65 70 75  
 Ser Leu Gln Lys Asp Leu His Glu Glu Met Asn Tyr Ile Thr Ala  
 80 85 90  
 Ile Ile Glu Glu Gln Pro Lys Asn Tyr Gln Val Trp His His Arg  
 95 100 105  
 Arg Val Leu Val Glu Trp Leu Arg Asp Pro Ser Gln Glu Leu Glu  
 110 115 120  
 Phe Ile Ala Asp Ile Leu Asn Gln Asp Ala Lys Asn Tyr His Ala  
 125 130 135  
 Trp Gln His Arg Gln Trp Val Ile Gln Glu Phe Lys Leu Trp Asp  
 140 145 150  
 Asn Glu Leu Gln Tyr Val Asp Gln Leu Leu Lys Glu Asp Val Arg  
 155 160 165  
 Asn Asn Ser Val Trp Asn Gln Arg Tyr Phe Val Ile Ser Asn Thr  
 170 175 180  
 Thr Gly Tyr Asn Asp Arg Ala Val Leu Glu Arg Glu Val Gln Tyr  
 185 190 195  
 Thr Leu Glu Met Ile Lys Leu Val Pro His Asn Glu Ser Ala Trp  
 200 205 210  
 Asn Tyr Leu Lys Gly Ile Leu Gln Asp Arg Gly Leu Ser Lys Tyr  
 215 220 225  
 Pro Asn Leu Leu Asn Gln Leu Leu Asp Leu Gln Pro Ser His Ser  
 230 235 240  
 Ser Pro Tyr Leu Ile Ala Phe Leu Val Asp Ile Tyr Glu Asp Met  
 245 250 255  
 Leu Glu Asn Gln Cys Asp Asn Lys Glu Asp Ile Leu Asn Lys Ala  
 260 265 270  
 Leu Glu Leu Cys Glu Ile Leu Ala Lys Glu Lys Asp Thr Ile Arg  
 275 280 285  
 Lys Glu Tyr Trp Arg Tyr Ile Gly Arg Ser Leu Gln Ser Lys His  
 290 295 300  
 Ser Thr Glu Asn Asp Ser Pro Thr Asn Val Gln Gln  
 305 310

<210> 25  
 <211> 452  
 <212> PRT



&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503141CD1

&lt;400&gt; 25

```

Met Ala Ala Ala Thr Gly Pro Ser Phe Trp Leu Gly Asn Glu Thr
  1           5           10           15
Leu Lys Val Pro Leu Ala Leu Phe Ala Leu Asn Arg Gln Arg Leu
          20           25           30
Cys Glu Arg Leu Arg Lys Asn Pro Ala Val Gln Ala Gly Ser Ile
          35           40           45
Val Val Leu Gln Gly Gly Glu Glu Thr Gln Arg Tyr Cys Thr Asp
          50           55           60
Thr Gly Val Leu Phe Arg Gln Glu Ser Phe Phe His Trp Ala Phe
          65           70           75
Gly Val Thr Glu Pro Gly Cys Tyr Gly Val Ile Asp Val Asp Thr
          80           85           90
Gly Lys Ser Thr Leu Phe Val Pro Arg Leu Pro Ala Ser His Ala
          95          100          105
Thr Trp Met Gly Lys Ile His Ser Lys Glu His Phe Lys Glu Lys
          110          115          120
Tyr Ala Val Asp Asp Val Gln Tyr Val Asp Glu Ile Ala Ser Val
          125          130          135
Leu Thr Ser Gln Lys Pro Ser Val Leu Leu Thr Leu Arg Gly Val
          140          145          150
Asn Thr Asp Ser Gly Ser Val Cys Arg Glu Ala Ser Phe Asp Gly
          155          160          165
Ile Ser Lys Phe Glu Val Asn Asn Thr Ile Leu His Pro Glu Ile
          170          175          180
Val Glu Cys Leu Phe Glu His Tyr Cys Tyr Ser Arg Gly Gly Met
          185          190          195
Arg His Ser Ser Tyr Thr Cys Ile Cys Gly Ser Gly Glu Asn Ser
          200          205          210
Ala Val Leu His Tyr Gly His Ala Gly Ala Pro Asn Asp Arg Thr
          215          220          225
Ile Gln Asn Gly Asp Met Cys Leu Phe Asp Met Gly Gly Glu Tyr
          230          235          240
Tyr Cys Phe Ala Ser Asp Ile Thr Cys Ser Phe Pro Ala Asn Gly
          245          250          255
Lys Phe Thr Ala Asp Gln Lys Ala Val Tyr Glu Ala Val Leu Arg
          260          265          270
Ser Ser Arg Ala Val Met Gly Ala Met Lys Pro Gly Val Trp Trp
          275          280          285
Pro Asp Met His Arg Leu Ala Asp Arg Ile His Leu Glu Glu Leu
          290          295          300
Ala His Met Gly Ile Leu Ser Gly Ser Val Asp Ala Met Val Gln
          305          310          315
Ala His Leu Gly Ala Val Phe Met Pro His Gly Leu Gly His Phe
          320          325          330
Leu Gly Ile Asp Val His Asp Val Gly Gly Tyr Pro Glu Gly Val
          335          340          345
Glu Arg Ile Asp Glu Pro Gly Leu Arg Ser Leu Arg Thr Ala Arg
          350          355          360
His Leu Gln Pro Gly Met Val Leu Thr Val Glu Pro Gly Ile Tyr

```

	365		370		375
Phe Ile Asp His	Leu Leu Asp Glu Ala	Leu Ala Asp Pro Ala	Arg		
	380		385		390
Ala Ser Phe Leu	Asn Arg Glu Val Leu	Gln Arg Phe Arg Gly	Phe		
	395		400		405
Gly Gly Val Arg	Ile Glu Glu Asp Val	Val Val Thr Asp Ser	Gly		
	410		415		420
Ile Glu Leu Leu	Thr Cys Val Pro Arg	Thr Val Glu Glu Ile	Glu		
	425		430		435
Ala Cys Met Ala	Gly Cys Asp Lys Ala	Phe Thr Pro Phe Ser	Gly		
	440		445		450
Pro Lys					

&lt;210&gt; 26

&lt;211&gt; 471

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500362CD1

&lt;400&gt; 26

Met Ala Ala Ala Thr Gly Pro Ser Phe Trp Leu Gly Asn Glu Thr		
1	5	10 15
Leu Lys Val Pro Leu Ala Leu Phe Ala Leu Asn Arg Gln Arg Leu		
	20	25 30
Cys Glu Arg Leu Arg Lys Asn Pro Ala Val Gln Ala Gly Ser Ile		
	35	40 45
Val Ser Phe Phe His Trp Ala Phe Gly Val Thr Glu Pro Gly Cys		
	50	55 60
Tyr Gly Val Ile Asp Val Asp Thr Gly Lys Ser Thr Leu Phe Val		
	65	70 75
Pro Arg Leu Pro Ala Ser His Ala Thr Trp Met Gly Lys Ile His		
	80	85 90
Ser Lys Glu His Phe Lys Glu Lys Tyr Ala Val Asp Asp Val Gln		
	95	100 105
Tyr Val Asp Glu Ile Ala Ser Val Leu Thr Ser Gln Lys Pro Ser		
	110	115 120
Val Leu Leu Thr Leu Arg Gly Val Asn Thr Asp Ser Gly Ser Val		
	125	130 135
Cys Arg Glu Ala Ser Phe Asp Gly Ile Ser Lys Phe Glu Val Asn		
	140	145 150
Asn Thr Ile Leu His Pro Glu Ile Val Glu Cys Arg Val Phe Lys		
	155	160 165
Thr Asp Met Glu Leu Glu Val Leu Arg Tyr Thr Asn Lys Ile Ser		
	170	175 180
Ser Glu Ala His Arg Glu Val Met Lys Ala Val Lys Val Gly Met		
	185	190 195
Lys Glu Tyr Glu Leu Glu Ser Leu Phe Glu His Tyr Cys Tyr Ser		
	200	205 210
Arg Gly Gly Met Arg His Ser Ser Tyr Thr Cys Ile Cys Gly Ser		
	215	220 225
Gly Glu Asn Ser Ala Val Leu His Tyr Gly His Ala Gly Ala Pro		
	230	235 240

Asn Asp Arg Thr	Ile Gln Asn Gly Asp Met Cys Leu Phe Asp Met	
	245	250 255
Gly Gly Glu Tyr	Tyr Cys Phe Ala Ser Asp Ile Thr Cys Ser Phe	
	260	265 270
Pro Ala Asn Gly	Lys Phe Thr Ala Asp Gln Lys Ala Val Tyr Glu	
	275	280 285
Ala Val Leu Arg	Ser Ser Arg Ala Val Met Gly Ala Met Lys Pro	
	290	295 300
Gly Val Trp Trp	Pro Asp Met His Arg Leu Ala Asp Arg Ile His	
	305	310 315
Leu Glu Glu Leu	Ala His Met Gly Ile Leu Ser Gly Ser Val Asp	
	320	325 330
Ala Met Val Gln	Ala His Leu Gly Ala Val Phe Met Pro His Gly	
	335	340 345
Leu Gly His Phe	Leu Gly Ile Asp Val His Asp Val Gly Gly Tyr	
	350	355 360
Pro Glu Gly Val	Glu Arg Ile Asp Glu Pro Gly Leu Arg Ser Leu	
	365	370 375
Arg Thr Ala Arg	His Leu Gln Pro Gly Met Val Leu Thr Val Glu	
	380	385 390
Pro Gly Ile Tyr	Phe Ile Asp His Leu Leu Asp Glu Ala Leu Ala	
	395	400 405
Asp Pro Ala Arg	Ala Ser Phe Leu Asn Arg Glu Val Leu Gln Arg	
	410	415 420
Phe Arg Gly Phe	Gly Gly Val Arg Ile Glu Glu Asp Val Val Val	
	425	430 435
Thr Asp Ser Gly	Ile Glu Leu Leu Thr Cys Val Pro Arg Thr Val	
	440	445 450
Glu Glu Ile Glu	Ala Cys Met Ala Gly Cys Asp Lys Ala Phe Thr	
	455	460 465
Pro Phe Ser Gly	Pro Lys	
	470	

<210> 27  
 <211> 458  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7503328CD1

<400> 27

Met Ala Ala Ser Arg Lys Pro Pro Arg Val Arg Val Asn His Gln	
1	5 10 15
Asp Phe Gln Leu Arg Asn Leu Arg Ile Ile Glu Pro Asn Glu Val	
	20 25 30
Thr His Ser Gly Asp Thr Gly Val Glu Thr Asp Gly Arg Met Pro	
	35 40 45
Pro Lys Val Thr Ser Glu Leu Leu Arg Gln Leu Arg Gln Ala Met	
	50 55 60
Arg Asn Ser Glu Tyr Val Thr Glu Pro Ile Gln Ala Tyr Ile Ile	
	65 70 75
Pro Ser Gly Asp Ala His Gln Ser Glu Tyr Ile Ala Pro Cys Asp	
	80 85 90
Cys Arg Arg Ala Phe Val Ser Gly Phe Asp Gly Ser Ala Gly Thr	

	95		100		105
Ala Ile Ile Thr	Glu Glu His Ala Ala	Met Trp Thr Asp Gly Arg			
	110		115		120
Tyr Phe Leu Gln	Ala Ala Lys Gln Met	Asp Ser Asn Trp Thr Leu			
	125		130		135
Met Lys Met Gly	Leu Lys Asp Thr Pro	Thr Gln Glu Asp Trp Leu			
	140		145		150
Val Ser Val Leu	Pro Glu Gly Ser Arg	Val Gly Val Asp Pro Leu			
	155		160		165
Ile Ile Pro Thr	Asp Tyr Trp Lys Lys	Met Ala Lys Val Leu Arg			
	170		175		180
Ser Ala Gly His	His Leu Ile Pro Val	Lys Glu Asn Leu Val Asp			
	185		190		195
Lys Ile Trp Thr	Asp Arg Pro Glu Arg	Pro Cys Lys Pro Leu Leu			
	200		205		210
Thr Leu Gly Leu	Asp Tyr Thr Gly Ile	Ser Trp Lys Asp Lys Val			
	215		220		225
Ala Asp Leu Arg	Leu Lys Met Ala Glu	Arg Asn Val Met Trp Phe			
	230		235		240
Val Val Thr Ala	Leu Asp Glu Ile Ala	Trp Leu Phe Asn Leu Arg			
	245		250		255
Gly Ser Asp Val	Glu His Asn Pro Val	Phe Phe Ser Tyr Ala Ile			
	260		265		270
Ile Gly Leu Glu	Thr Ile Met Leu Phe	Ile Asp Gly Asp Arg Ile			
	275		280		285
Asp Ala Pro Ser	Val Lys Glu His Leu	Leu Leu Asp Leu Gly Leu			
	290		295		300
Glu Ala Glu Tyr	Arg Ile Gln Val His	Pro Tyr Lys Ser Ile Leu			
	305		310		315
Ser Glu Leu Lys	Ala Leu Cys Ala Asp	Leu Ser Pro Arg Glu Lys			
	320		325		330
Val Trp Val Ser	Asp Lys Ala Ser Tyr	Ala Val Ser Glu Thr Ile			
	335		340		345
Pro Lys Asp His	Arg Cys Cys Met Pro	Tyr Thr Pro Ile Cys Ile			
	350		355		360
Ala Lys Ala Val	Lys Asn Ser Ala Glu	Ser Glu Gly Met Arg Arg			
	365		370		375
Ala His Ile Lys	Asp Ala Val Ala Leu	Cys Glu Leu Phe Asn Trp			
	380		385		390
Leu Glu Lys Glu	Val Pro Lys Gly Gly	Val Thr Glu Ile Ser Ala			
	395		400		405
Ala Asp Lys Ala	Glu Glu Phe Arg Arg	Gln Gln Ala Asp Phe Val			
	410		415		420
Asp Leu Ser Phe	Pro Thr Ile Ser Ser	Gln Ser Leu Arg Arg Ile			
	425		430		435
Gly Pro Cys Pro	Trp Met Arg Cys Thr	Leu Leu Thr Arg Val Leu			
	440		445		450
Asn Thr Arg Met	Ala Pro Gln Met				
	455				

&lt;210&gt; 28

&lt;211&gt; 695

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7510464CD1

&lt;400&gt; 28

Met	Ala	Ala	Ser	Arg	Lys	Pro	Pro	Arg	Val	Arg	Val	Asn	His	Gln	1	5	10	15
Asp	Phe	Gln	Leu	Arg	Asn	Leu	Arg	Ile	Ile	Glu	Pro	Asn	Glu	Val	20	25	30	
Thr	His	Ser	Gly	Asp	Thr	Gly	Val	Glu	Thr	Asp	Gly	Arg	Met	Pro	35	40	45	
Pro	Lys	Val	Thr	Ser	Glu	Leu	Leu	Arg	Gln	Leu	Arg	Gln	Ala	Met	50	55	60	
Arg	Asn	Ser	Glu	Tyr	Val	Thr	Glu	Pro	Ile	Gln	Ala	Tyr	Ile	Ile	65	70	75	
Pro	Ser	Gly	Asp	Ala	His	Gln	Ser	Glu	Tyr	Ile	Ala	Pro	Cys	Asp	80	85	90	
Cys	Arg	Arg	Ala	Phe	Val	Ser	Gly	Phe	Asp	Gly	Ser	Ala	Gly	Thr	95	100	105	
Ala	Ile	Ile	Thr	Glu	Glu	His	Ala	Ala	Met	Trp	Thr	Asp	Gly	Arg	110	115	120	
Tyr	Phe	Leu	Gln	Ala	Ala	Lys	Gln	Met	Asp	Ser	Asn	Trp	Thr	Leu	125	130	135	
Met	Lys	Met	Gly	Leu	Lys	Asp	Thr	Pro	Thr	Gln	Glu	Asp	Trp	Leu	140	145	150	
Val	Ser	Val	Leu	Pro	Glu	Gly	Ser	Arg	Val	Gly	Val	Asp	Pro	Leu	155	160	165	
Ile	Ile	Pro	Thr	Asp	Tyr	Trp	Lys	Lys	Met	Ala	Lys	Val	Leu	Arg	170	175	180	
Ser	Ala	Gly	His	His	Leu	Ile	Pro	Val	Lys	Glu	Asn	Leu	Val	Asp	185	190	195	
Lys	Ile	Trp	Thr	Asp	Arg	Pro	Glu	Arg	Pro	Cys	Lys	Pro	Leu	Leu	200	205	210	
Thr	Leu	Gly	Leu	Asp	Tyr	Thr	Gly	Ile	Ser	Trp	Lys	Asp	Lys	Val	215	220	225	
Ala	Asp	Leu	Arg	Leu	Lys	Met	Ala	Glu	Arg	Asn	Val	Met	Trp	Phe	230	235	240	
Val	Val	Thr	Ala	Leu	Asp	Glu	Ile	Ala	Trp	Leu	Phe	Asn	Leu	Arg	245	250	255	
Gly	Ser	Asp	Val	Glu	His	Asn	Pro	Val	Phe	Phe	Ser	Tyr	Ala	Ile	260	265	270	
Ile	Gly	Leu	Glu	Thr	Ile	Met	Leu	Phe	Ile	Asp	Gly	Asp	Arg	Ile	275	280	285	
Asp	Ala	Pro	Ser	Val	Lys	Glu	His	Leu	Leu	Leu	Asp	Leu	Gly	Leu	290	295	300	
Glu	Ala	Glu	Tyr	Arg	Ile	Gln	Val	His	Pro	Tyr	Lys	Ser	Ile	Leu	305	310	315	
Ser	Glu	Leu	Lys	Ala	Leu	Cys	Ala	Asp	Leu	Ser	Pro	Arg	Glu	Lys	320	325	330	
Val	Trp	Val	Ser	Asp	Lys	Ala	Ser	Tyr	Ala	Val	Ser	Glu	Thr	Ile	335	340	345	
Pro	Lys	Asp	His	Arg	Cys	Cys	Met	Pro	Tyr	Thr	Pro	Ile	Cys	Ile	350	355	360	
Ala	Lys	Ala	Val	Lys	Asn	Ser	Ala	Glu	Ser	Glu	Gly	Met	Arg	Arg	365	370	375	
Ala	His	Ile	Lys	Asp	Ala	Val	Ala	Leu	Cys	Glu	Leu	Phe	Asn	Trp	380	385	390	

Leu	Glu	Lys	Glu	Val	Pro	Lys	Gly	Gly	Val	Thr	Glu	Ile	Ser	Ala	
				395					400					405	
Ala	Asp	Lys	Ala	Glu	Glu	Phe	Arg	Arg	Gln	Gln	Ala	Asp	Phe	Val	
				410					415					420	
Asp	Leu	Ser	Phe	Pro	Thr	Ile	Ser	Ser	Thr	Gly	Pro	Asn	Gly	Ala	
				425					430					435	
Ile	Ile	His	Tyr	Ala	Pro	Val	Pro	Glu	Thr	Asn	Arg	Thr	Leu	Ser	
				440					445					450	
Leu	Asp	Glu	Val	Tyr	Leu	Ile	Asp	Ser	Gly	Ala	Gln	Tyr	Lys	Asp	
				455					460					465	
Gly	Thr	Thr	Asp	Val	Thr	Arg	Thr	Met	His	Phe	Gly	Thr	Pro	Thr	
				470					475					480	
Ala	Tyr	Glu	Lys	Glu	Cys	Phe	Thr	Tyr	Val	Leu	Lys	Gly	His	Ile	
				485					490					495	
Ala	Val	Ser	Ala	Ala	Val	Phe	Pro	Thr	Gly	Thr	Lys	Gly	His	Leu	
				500					505					510	
Leu	Asp	Ser	Phe	Ala	Arg	Ser	Ala	Leu	Trp	Asp	Ser	Gly	Leu	Asp	
				515					520					525	
Tyr	Leu	His	Gly	Thr	Gly	His	Gly	Val	Gly	Ser	Phe	Leu	Asn	Val	
				530					535					540	
His	Glu	Gly	Pro	Cys	Gly	Ile	Ser	Tyr	Lys	Thr	Phe	Ser	Asp	Glu	
				545					550					555	
Pro	Leu	Glu	Ala	Gly	Met	Ile	Val	Thr	Asp	Glu	Pro	Gly	Tyr	Tyr	
				560					565					570	
Glu	Asp	Gly	Ala	Phe	Gly	Ile	Arg	Ile	Glu	Asn	Val	Val	Leu	Val	
				575					580					585	
Val	Pro	Val	Lys	Thr	Lys	Tyr	Asn	Phe	Asn	Asn	Arg	Gly	Ser	Leu	
				590					595					600	
Thr	Phe	Glu	Pro	Leu	Thr	Leu	Val	Pro	Ile	Gln	Thr	Lys	Met	Ile	
				605					610					615	
Asp	Val	Asp	Ser	Leu	Thr	Asp	Lys	Glu	Glu	Leu	Trp	Asn	Gly	Ile	
				620					625					630	
Leu	Pro	Ala	Arg	Ser	Leu	Phe	Cys	Leu	Phe	Gln	Phe	Thr	Val	Arg	
				635					640					645	
Leu	Ala	Gln	Gln	Leu	Pro	Pro	Asp	Leu	Gln	Gly	Cys	Asp	Trp	Glu	
				650					655					660	
Gly	Ile	Ala	Glu	Thr	Gly	Pro	Pro	Gly	Ser	Ser	Arg	Val	Ala	His	
				665					670					675	
Gln	Arg	Asp	Ala	Thr	His	Leu	Gln	Thr	Ala	Leu	Ile	Asn	Thr	Ser	
				680					685					690	
Pro	Val	Leu	Phe	Leu											
				695											

&lt;210&gt; 29

&lt;211&gt; 140

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7510394CD1

&lt;400&gt; 29

Met	Ala	Ala	Ala	Met	Pro	Leu	Ala	Leu	Leu	Val	Leu	Leu	Leu	Leu	
1				5				10					15		
Gly	Pro	Gly	Gly	Trp	Cys	Leu	Ala	Glu	Pro	Pro	Arg	Asp	Ser	Leu	

	20		25		30
Arg	Glu	Glu	Leu	Val	Ile
	35		40		45
Ala	Thr	Phe	Gln	Phe	Arg
	50		55		60
Glu	Gly	Val	Ser	His	Tyr
	65		70		75
Leu	Ile	Ser	Lys	Tyr	Ser
	80		85		90
Gln	Gly	Phe	Trp	Arg	Thr
	95		100		105
Ala	Pro	Ser	Gly	Ala	Glu
	110		115		120
Thr	Glu	Phe	Ser	Ser	Gln
	125		130		135
Val	Ala	Pro	Gly	Gln	
	140				

&lt;210&gt; 30

&lt;211&gt; 191

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500745CD1

&lt;400&gt; 30

Met	Ala	Ala	Ala	Met	Pro	Leu	Ala	Leu	Leu	Val	Leu	Leu	Leu	Leu
1				5						10				15
Gly	Pro	Gly	Gly	Trp	Cys	Leu	Ala	Glu	Pro	Pro	Arg	Asp	Ser	Leu
				20						25				30
Arg	Glu	Glu	Leu	Val	Ile	Thr	Pro	Leu	Pro	Ser	Gly	Asp	Val	Ala
				35						40				45
Ala	Thr	Phe	Gln	Phe	Arg	Thr	Arg	Trp	Asp	Ser	Glu	Leu	Gln	Arg
				50						55				60
Glu	Gly	Val	Ser	His	Tyr	Arg	Leu	Phe	Pro	Lys	Ala	Leu	Gly	Gln
				65						70				75
Leu	Ile	Ser	Lys	Tyr	Ser	Leu	Arg	Glu	Leu	His	Leu	Ser	Phe	Thr
				80						85				90
Gln	Gly	Phe	Trp	Arg	Thr	Arg	Tyr	Trp	Gly	Pro	Pro	Phe	Leu	Gln
				95						100				105
Ala	Pro	Ser	Val	Trp	Ile	Asn	Leu	Gly	Arg	Ser	Ser	Val	Met	Ser
				110						115				120
Ser	Gln	Gly	Ser	Ser	Ala	Pro	Leu	Ser	Thr	Ser	Ser	Thr	Pro	Pro
				125						130				135
Thr	Gln	Ser	Leu	Pro	Leu	Pro	Pro	Ser	Asn	Pro	Trp	Val	Trp	Pro
				140						145				150
Met	Thr	Leu	Thr	Thr	Thr	Phe	Cys	Ala	Met	Leu	Cys	Cys	Arg	Gly
				155						160				165
Arg	Trp	Ser	Ala	Pro	Lys	Thr	Ser	Pro	Pro	Gly	Arg	Ser	Ser	Cys
				170						175				180
Pro	Val	Val	Pro	Arg	Gln	Ala	Ser	Leu	Cys	Cys				
				185						190				

&lt;210&gt; 31

<211> 145  
 <212> PRT  
 <213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 7500929CD1

<400> 31

```

Met Ser Ala Trp Ala Ala Ala Ser Leu Ser Arg Ala Ala Ala Arg
 1          5          10          15
Cys Leu Leu Ala Arg Gly Pro Gly Val Arg Ala Ala Pro Pro Arg
          20          25          30
Asp Pro Arg Pro Ser His Pro Glu Pro Arg Gly Cys Gly Ala Ala
          35          40          45
Pro Gly Arg Thr Leu His Phe Thr Ala Ala Val Pro Ala Gly His
          50          55          60
Asn Lys Trp Ser Lys Val Arg His Ile Lys Gly Pro Lys Asp Val
          65          70          75
Glu Arg Ser Arg Ile Phe Ser Lys Leu Cys Leu Asn Ile Arg Leu
          80          85          90
Ala Val Lys Ala Arg Arg Pro Lys Asp Arg Thr Cys Asp Leu Glu
          95          100         105
Ala Lys Gly Ile Ser Leu Val Gly Pro Pro Cys Gln Leu Cys Cys
          110         115         120
Cys Leu Arg Ala Ile Trp Met Ser Val Pro Thr Pro Ser Arg Met
          125         130         135
Gln Gly Arg Thr Thr Gln Leu Val Arg Leu
          140         145

```

<210> 32  
 <211> 2129  
 <212> DNA  
 <213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 8268274CB1

<400> 32

```

ggcggcgccg gccggagccg gaaggcgggg aggggcccgc cgttggggcc gaggcggcgg 60
cggcgggcggc ggcggctggg gagaagcgct ctctgcgcct gcccgaggcc ggagcggcgg 120
ggcccgccgc tcctccccc agcgccgcgg aggggggagg aggaagatgg agaccacat 180
ctcatgcctg ttcccgagc tgctggccat gatcttcggc tacctggacg tccgggacaa 240
ggggcgccgc gcgcaggtgt gcaccgcctg gcgggacgcc gcctaccaca agtcggtgtg 300
gcgggggggtg gaggccaaagc tgcaacctgc cggggccaac ccgtcgctgt tcccagcct 360
gcaggcccg ggcacccgcc ggggtgcagat cctgagcctc cggcgagcc tcagctacgt 420
gatccagggc atggccaaca tcgagagcct caacctcagc ggctgctaca acctcaccga 480
caacgggctg ggccaacgct ttgtgcagga gatcggtctc ctgcgcgctc tcaacctgag 540
ctctgcaag cagatcactg acagcagcct gggccgcata gccagttacc tcaagggcct 600
ggaggtgctg gagctgggag gttgcagcaa catcaccac actggccttc tgctcatcgc 660
ctgggggtctg cagcgccctca agagccttaa cctccgcagc tgccgccacc ttctggatgt 720
gggcatcggg cacctggccg gcatgacgcg cagcgcgggc gagggctgcc tgggcctgga 780
gcagctcacg ctacaggact gccagaagct cacagatctt tctctaaagc acatctcccg 840
agggctgacg ggctgaggc tcctcaacct cagcttctgt gggggaatct cggacgctgg 900
cctcctgcac ctgtgcaca tgggcagcct gcgcagcctc aacctgcgct cctgtgacaa 960

```



```

catcagtgac acgggcatca tgcattctggc catgggcagc ctgcgcctct cggggctgga 1020
tgtttcgttc tgtgacaagg tgggagacca gagtctggct tacatagccc aggggctgga 1080
tggcctcaag tctctctccc tctgctcctg ccacatcagt gatgatggca tcaaccgcat 1140
ggtgcggcag atgcacgggc tgcgcacgct caacattgga cagtgtgtgc gcatcacgga 1200
caagggcctg gagctgatcg ctgagcacct gagccaactc accggcatag acctgtacgg 1260
ctgcaccoga atcaccaagc gcggcctgga gcgcatcacg cagctgccgt gcctcaagga 1320
ggcacgaggg gattttttctc cattattcac tgtgagaact cggggaagct ccagaagggtg 1380
agggagaggg gacaacgaca tggttcccgt ggatctttaa cttccagact tgcccgtctc 1440
gcgcctctgg cactctgggtg atgacagctc aggtttccct gcctgtcact gctcgggcag 1500
aggctgctgc ccagggtctc tgctccggtg ccttgtgaag ctgcattctc ctgccggttt 1560
ctccagttct ggggacagtg gtttgctctg agacctcgct tcctttatgg atccaaggag 1620
acttgctttt tcagtctgtt cagcttttta cttgctagga tggaattgca atttgcaagc 1680
ttcttcgaca ggaaactaca agttccacac tttaatttta tacatataaa tatatacatg 1740
tgtacatata tctatgtaca ggggtattat atatatacat ataagatgat gatatatata 1800
atgatgatat gtattactga gaacgtaaaa tatcattaca tagtgatagc tggacacaca 1860
aggaattcac aactcccaa agaaaataca tctggatgac ctgcctagca gtttcccat 1920
gagatagagg aatgtctacg tatttcattc cctgttcctg ccctgaaaca atttcaatca 1980
ctgacaaatc attatcattc attaataatg tttactgagt gcccatatgt gaaagaaatc 2040
cactctacat tccacagatg catttcctct cccacgggg tttccatttt aatgggaaca 2100
atgtagaata tatctgtctt cccttaaaa 2129

```

<210> 33

<211> 3489

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 7500515CB1

<400> 33

```

ggcagacact ggagccacga tgaagcccc aaggcctgtc cgtacctgca gcaaagttct 60
cgtctctgtt tcaactgttg ccattccacca gactactact gccgaaaaga atggcatcga 120
catctacagc ctaccgttg actccagggt ctcatccga tttgccaca cggctcgtcac 180
cagccgagtg gtcaataggg ccaatactgt gcaggaggcc accttcaga tggagctgcc 240
caagaaagcc ttcatcacca acttctccat gatcatcgat ggcatgacct acccagggat 300
catcaaggag aaggctgaag cccaggcaca gtacagcgca gcagtggcca agggaaagag 360
cgctggcctc gtcaaggcca cggggagaaa catggagcag tccagggtgt cggtcagtgt 420
ggctcccaat gccaatgaca cctttgagct ggtctatgag gagctgctca agcggcggtt 480
gggggtgtac gagctgctgc tgaaagtgcg gcccagcag ctggtcaagc acctgcagat 540
ggacattcac atcttcgagc cccagggtat cagctttctg gagacagaga gcaccttcat 600
gaccaaccag ctggttagacg ccctcaccac ctggcagaat aagaccaagg ctccatccg 660
gttcaagcca acactttccc agcagcaaaa gtcccagag cagcaagaaa cagtcttgga 720
cggcaacctc attatccgct atgatgtgga cggggccatc tccggggggt ccattcagat 780
cgagaacggc tactttgtac actactttgc cccgaggggc ctaaccacaa tgcccaagaa 840
tgtgtgtctt gtcatgtaca agagcggctc catgagtggc aggaaaatcc agcagacctg 900
ggaagcccta atcaagatcc tggatgacct cagccccaga gaccagttca acctcatcgt 960
cttcagtaca gaagcaactc agtggaggcc atcactggtg ccagcctcag ccgagaacgt 1020
gaacaaggcc aggagctttg ctgcgggcat ccaggccctg ggagggacca acatcaatga 1080
tgcaatgctg atggctgtgc agttgtctga cagcagcaac caggaggagc ggctgcccga 1140
agggagtgtc tcaactcatc tctgtctcac cgatggcgac cccactgtgg gggagactaa 1200
ccccaggagc atccagaata acgtgcggga agctgtaagt ggccggtaca gcctcttctg 1260
cctgggcttc ggtttcgacg tcagctatgc cttcctggag aagctggcac tggacaatgg 1320
cggcctggcc cggcgcatcc atgaggactc agactctgcc ctgcagctcc aggacttcta 1380
ccaggaagtg gccaaaccac tgctgacagc agtgacctt gagtaccacaa gcaatgccgt 1440
ggaggagggtc actcagaaca acttccggct cctcttcaag ggctcagaga tgggtgggtggc 1500

```

```

tggaagctc caggaccggg ggctgatgt gtcacagcc acagtcagt ggaagctgcc 1560
tacacagaac atcactttcc aaacggagtc cagtgtggca gagcaggagg cggagttcca 1620
gagccccaag tatatcttcc acaacttcat ggagaggctc tgggcatacc tgactatcca 1680
gcagctgctg gagcaaactg tctccgcctc cgatgctgat cagcaggccc tccggaacca 1740
agcgtgaat ttatcacttg cctacagctt tgtcacgcct ctcacatcta tggtagtcac 1800
caaaccgat gaccaagagc agtctcaagt tgctgagaag cccatggaag gcgaaagtag 1860
aaacaggaat gtccactcag ctggagctgc tggctcccg atgaatttca gacctggggt 1920
tctcagctcc aggcaacttg gactcccagg acctcctgat gtctctgacc atgctgctta 1980
ccaccccttc cgccgtcttg ccatcttgcc tgcttcagca ccaccagcca cctcaaacc 2040
tgatccagct gtgtctctg tcatgaatat gaaaatcgaa gaaacaacca tgacaacca 2100
aaccaccagc cccatacagg ctccctctgc catcctgcca ctgcctgggc agagtgtgga 2160
gcggtctctg gtggaccca gacaccgcca ggggccagt aacctgctct cagaccctga 2220
gcaaggggtt gaggtgactg gccagtatga gagggagaag gctgggttct catggatcga 2280
agtgaccttc aagaaccccc tggatgggt tcacgcctcc cctgaacacg tgggtgtgac 2340
tcggaaccga agaagctctg cgtacaagt gaaggagacg ctattctcag tgatgcccg 2400
cctgaagatg accatggaca agacgggtct cctgctgctc agtgaccag acaaagtac 2460
catcgccctg ttgttctggg atggccgtgg ggaggggctc cggtccttc tgcgtgacac 2520
tgaccgcttc tccagccacg ttggagggac cttggccag tttaccagg aggtgctctg 2580
gggatctcca gcagcatcag atgacggcag acgcacgctg agggttcagg gcaatgacca 2640
ctctgccacc agagagcgca ggctggatta ccaggagggg ccccgaggag tggagatttc 2700
ctgctggtct gtggagctgt agttctgat gaaggagctg tgccaccct gtacacttgg 2760
cttccccctg caactgcagg gccgttctg gggcctggac caccatgggg aggaagagtc 2820
ccactcatta caaataaaga aaggtggtgt gagcctggga aaaaaaaaaa aaaaaaaaaa 2880
aaaaaaaaaa aaaaaaaagg gggggccccc aaaataagga ccccaaccc cgggggatat 2940
aaatactcgg ggacaagcgc ttaccactgg cggaggcgtg ttaatacca cccccacat 3000
gggggggggca acgttatatt ccgtattgt cagagggggc atccccacta aatgaggggc 3060
ggcgtaatta aactatctcg gcaaaaggac ccagtggaat gaccccgta tttatatgta 3120
ctgacgcaga caacgacaca ctagtcaac aacacgacag ccacatcagt acctcgtcga 3180
catgctgacg aagagtcgga cccacatac acacaactaa aacaaccaa ctctacaca 3240
caactacac acatctaata tccgactcag ccccccaacc cacaccata acacacacac 3300
acagaacaac caacaatata atactatcat taactataaa acgacaaacc ctcataacac 3360
ttatataatg cagtacatcc taatcacacc acaacaacaa aaaaacaacc atcatacatc 3420
atccactaac actacattac aaaaccatca aaaaacgcca cacaccacc acactctcca 3480
ctattctct
3489

```

&lt;210&gt; 34

&lt;211&gt; 2996

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2256826CB1

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 360, 370

&lt;223&gt; a, t, c, g, or other

&lt;400&gt; 34

```

ctgtttgcgc gcggacggag gagcggtgga ctgggggcag cggagggggc cccgcgcacc 60
gtgcggtcct gctctcctct cctcgtctgg tccgcgagca cgcgcgcctc tgcacccgcc 120
ccccagaccc ccttttcccc ccccccgcc gcttccgtgt ctgcctcct cccggtgggc 180
tcgctgcct cgaggcgca gggggtcgtg gcgcgcgcgc gcaccggtg ctccggggag 240
cgccgtcagg gcgggccccg tgggggggag ggggtggttt ggaccttttc cgtaggggtc 300
ctccctttcg gccctcccc ctaccgggag ctccgaggcc ctggcggtc tgtccaatgn 360

```

```

agcagtggcn gttgctggca ggtggggagt gtttttgttt tgggttgaag ttgaggctga 420
ggagagagcc gagctagcga cgagcagtcg ttgcccgcgc cggcgccgcg ggaggtggtg 480
gaggcctagc cggagccgag aggtctcttg ttcccgtccc acggtcccgc cgtcaccctt 540
ccggcgccca gtcccgcgtc cggaactccc gggcctgtcc tgggccccgc gtctgtgcac 600
tccgctcgcc gcagcgcccg gcccgggccg caccgcgcgc ccccatgagg agggacgtga 660
acggagtgac caagagcagg tttgagatgt tctcaaatag tgatgaagct gtaatcaata 720
aaaaacttcc caaagaactc ctgttacgga tattttcttt tctagatgtt gttaccctgt 780
gccgctgtgc tcaggctctc agggcctgga atgttctggc tctggatggc agtaactggc 840
agcgaattga cctatttgat ttccagaggg atattgaggg ccgagtagtg gagaatattt 900
caaaacgatg tgggggcttt ttacgaaagt taagtcttcg tggatgtctt ggagtgggag 960
acaatgcatt aagaaccttt gcacaaaact gcaggaacat tgaagtactg aatctaaatg 1020
ggtgtacaaa gacaacagac gctacatgta ctagccttag caagttctgt tccaaactca 1080
ggcaccttga cttggcttcc tgtacatcaa taacaaacat gtctctaaaa gctctgagtg 1140
agggatgtcc actgttggag cagtgaaca tttcctgggtg tgaccaagta accaaggatg 1200
gcattcaagc actagtgagg ggctgtgggg gtctcaaggc cttattctta aaaggctgca 1260
cgcagctaga agatgaagct ctcaagtaca taggtgcaca ctgccctgaa ctgggtgactt 1320
tgaacttgca gacttgcttg caaatcacag atgaaggtct cattactata tgcagagggt 1380
gccataagtt acaatccctt tgtgcctctg gctgctccaa catcacagat gccatcctga 1440
atgctctagg tcagaactgc ccacggctta gaatttgga agtggcaaga tgttctcaat 1500
taacagatgt gggctttacc actctagcca ggaattgcca tgaacttgaa aagatggacc 1560
tggaagagtg tgttcagata acagatagca cattaatcca actttctata cactgtcctc 1620
gacttcaagt attgagtctg tctcactgtg agctgatcac agatgatgga attcgtcacc 1680
tggggaatgg ggctgcgc catgaccagc taggagtgat tgagctggac aactgccac 1740
taatcacaga tgcacccctg gagcacttga agagctgtca tagccttgag cggatagaac 1800
tctatgactg ccagcaaatc acacgggctg gaatcaagag actcaggacc catttacc 1860
atattaaagt ccacgcctac ttgcacctg tcaactccac cccatcagta gggggcagca 1920
gacagcgctt ctgcagatgc tgcacatcc tatgacaatg gaggtggtca accttggcga 1980
actgagtatt taatgacact tctagagcta ccgtggagtc tctccagtgg aagcaacccc 2040
agtgttctga gcaagggtta caaagtgagg gagggcagtg tccagatccc cagagccaca 2100
catacatata catacacacc cttaccccca tccactctag ctttgtgacc atgggactga 2160
agtttgtgat ggctttttta tcaagtagat tggtaaaatt taaccattcc tgttgagggtg 2220
cccataagaa aatcataggc caagataggg aggggcattc cagcaaacc cgtgttaatg 2280
ctactgtggt ttttaaattt ttgtctaggg gtttctttgg ggattttaga acagcatctg 2340
ctgtcctccg gggtaagaa aagcatggaa agacaatata tgatgtaccc agggaccaga 2400
aagaaaattt ctttgcattc tagaaatggt agacattcat tgtgactaaa gagcttctat 2460
gcttccttgt ttccatgcca acatgctgag catgctcaca aagaaggctc gtccattcct 2520
cctgtgtttt agtatttggc ccagaggttt cctaaatggt tgccttgaaa tcaactgtggt 2580
ccaaatgtaa ttcttacaca ctcaaattat cactgtctgt agcacacttg tgcacctgtc 2640
ttacattctc tgttgctccc cccacactc ttgctcagtc tgtcacctgt tcagtctgct 2700
tactcactca attgttacc ttttgcgtgt gtcgtgttta cagtgttgc tttgaatgat 2760
tagttgggat taccaaacat tttttaaaaa gatattatca ataaatattt ttttaattct 2820
aaattttaaa aaaaaaaaaa aagggggggg ccgcttaaaa ggtcccaagt ttgattacgc 2880
ttgcttccga cgtcatagcg ggccgcagaa ttccgatatc aagcttttgg atccggggac 2940
ttcggggggg cccgttccca atgcccctat gtgattgatt acgccccaca ggcgct 2996

```

<210> 35

<211> 1860

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 7686186CB1

<400> 35

gcctccggac tgaccttgcg gctgtatggc gcacgcccc tgtaccggga tgaagcgccc 60

```

gatctttggg ctgtgttcgc gcgaactggc ggcgcgggcc gggctgcctg ccgtgcccg 120
taccgcacta cgtgcccagc ggggtcgtca acgccttcgc caccggttcg aagcatcacg 180
cggccatcgc gctgaccgac ggcttgcctg gtagcctcac gcccgcgag ttgaccggcg 240
tgctcggcca cgaaatcgcg catattgcga acgaggattt gcgtgtcatg ggcttggcgg 300
attccatcag ccggctgacc catctgctgg ccctgctggg gcagcttgcg atcgtgtctca 360
gcttgccagc gctgctgctt ggagtcgagg aagtcaattg gcccgcggtg cttctactgg 420
cggtcgcgcc acagctggcc ttgctggctc agttgggctt gtccagggtg cggaattcg 480
acgccgaccg gctcgtgccc gaattgaccg gcgaccgcca cgggctggcc tcggcgctcg 540
ccaagatcga gcgggtgagc cgctcctggc gcgcctggct gctgccccgg atgaggcaat 600
ccggaacctt cctgggttcg cagcatccg gcgacggctg aacgcattga gcgcttgctg 660
gaacttgctc cgccgcccgc gatgccgcg tttccatcgg cccgtttcgt ccccagggtg 720
accgtatcac cagctccgcc acgctggggc accggcgggc tttgacgctg atttcaacat 780
ggagaatgac gtgacctacc ccgacccta cagccggccg gcgcgggacc gcttcatccg 840
gcgtgggtc gtcatcactg gctgcatcgc cgactcatg ctgctgtggc agttcctgcc 900
cgccatcgaa gcctggttca gtccccacga aacgcaggag cgacgggtga cgccgcggcg 960
cgacctggcc gccgacgaaa aaaccacat cgagctgttc gagaaatcgc gcgggtcggg 1020
ggtttacatc accacggcac aactagtgcg tgacgtctgg tcgcgcaatg tcttttcctg 1080
gcgcgcggc accgggtccg gcttcatctg ggacgatgcc ggccacgtgg tgaccaactt 1140
ccacgtgata cagggggcat cgtctgccac ggtcaaaact gccgacggtc gcgattatca 1200
ggctgcgctc gttggcgcca gtctgcgca cgacatcgcg gtactcaaga ttggcgctcg 1260
cttcaagcgc ccgcggggcg tgccggtggg caccagtgcg gatctcaagg tggggcaaaa 1320
ggtctttgcc attggcaatc cttcggggt cgactggagc ctcaccaccg gcacgtctc 1380
ggcgcttgac cgcaccttt ccggcgagc cagtggcccg gccattgacc acctgatcca 1440
gaccgacgcc gctatcaacc ccggcaattc cgggtggccg ctgctcgatt cggctggggc 1500
gctgatcgcc atcaataacc ccatctacag tccgtctggc gcctcgcccg gcacggctt 1560
tgcggtgcg gtcgataacc tcatgcgcgt ggtgccgcaa ctcataaaga ccggcaagta 1620
catccgtccg gcgctgggca tcgaggtgga tgagcagctc aacgcgcgctc tgcaggcgct 1680
gaccggcagt aaggggcgat tcgtattgcg cgtgacgccg ggctcggcg cgacaggggc 1740
cgggctcgtc ggctgcagg tcaccgcagg cggcatcgct cccggcgatc gcgttatcag 1800
catcgacggg atcgccgtcg acccggaat tcgggacgct acctgctgac ctcttcagac 1860

```

<210> 36

<211> 1334

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 72617436CB1

<400> 36

```

cctgatgata tgtcactgtc tcccatcact ccagatggg accgtctagt ctcaggaaaa 60
caagctcttg gctcccactg attctacatt atggtgtgat cctaggagcg cccctggcct 120
ccagctgcgc aggagcctgt ggtaccagct tcccagatgg cctcaccctt gagggaacct 180
aggcctccgg ggacaaggac attcctgcaa ttaaccaagg gctcatcctg gaagaaacct 240
cagagagcag cttcctcatc gagggggaca tcatccggcc gagtcccttc cgactgctgt 300
cagcaaccag caacaaatgg cccatgggtg gtagtgggtg cgtggagggt cccttctctg 360
tctccagcaa gtacgatgag cccagccgcc aggtcatcct ggaggctctt gcggagtctt 420
aacgttcac gtgcatcagg tttgtcacct atcaggacca gagagacttc atttccatca 480
tccccatcta tgggtgcttc tcgagtgtgg ggcgagtg agggatgcag gtggtctccc 540
tggcgccac gtgtctccag aaggggccgg gcattgtcct tcatgagctc atgcatgtgc 600
tgggcttctg gcacgagcac acggggcccg cttatatcgt gtcaactgga 660
acgagatcct gccaggcttt gaaatcaact tcatcaagtc tcggagcagc aacatgctga 720
cgccctatga ctactcctct gtgatgcact atgggaggct tgcttcagc cggcggtggc 780
tgccaccat cacaccactt tgggccccca gtgtccacat cggccagcga tggaaacctga 840
gtgcctcgga catcaccggg gtctccaaac tctacggctg cagcccaagt ggccccaggc 900

```

```

ccccgtgggag aggggtcccat gcccacagca ctggtaggag ccccgctccg gcctccctat 960
ctctgcagcg gcttttggag gcaactgtcgg cggaatccag gagccccgac cccagtgggtt 1020
ccagtgcggg agggccagccc gttcctgcag ggctgggga gagccacat ggggtgggagt 1080
cccctgccct gaaaaagctc agtgagagg cctcggcaag gcagcctcag accctagctt 1140
cctccccaag atcaaggcct ggagcaggtg ccccgggtgt tgctcaggag cagtccctggc 1200
tggcgggagt gtccaccaag cccacagtcc catcttcaga agcaggaatc cagccagtcc 1260
ctgtccaggg aagcccagct ctgccagggg gctgtgtacc tagaaatcat ttcaagggga 1320
tgtccgaaga ttaa                                     1334

```

&lt;210&gt; 37

&lt;211&gt; 2070

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7501945CB1

&lt;400&gt; 37

```

gctggaacag tgggtccctcg tcgtgactca gagccaggcc cagtgtctgcc cccatcagac 60
cccgttccct ttccggaatg tggaaggatg cgagtgtgtg cctgaggggtg tggtcctgtg 120
gggggtgggtg gtcttggccc gtgcacacct gcagtgtgtg catggtgttc ctgtgggagc 180
gtgtgtgcac gccacacaa gaggcacttc ccaggccgtg tgggtctcta ctccctcaca 240
gcaggcctgc atggcctccc tactcttctc ccacatccca gggctggggc tcacagcgtc 300
ccctgttcca gagtcccttc atgtccccc aaggcatgtg gccctgtgtc tgggtgcagag 360
gacctgtcac cagggatggg attcattgct ctgcacaccc acccaactag cacatcttgg 420
ccgagcttca attacggggc cagcatagga ctaaggttca gtggagacct gagaccagt 480
ccccgcccac gccagaagcc aaggggtacc tgccgacgt gccgggaggg ggtgctgtga 540
tctgcctgtg agcaggggtg ctgaggcctt tggaaatggt tgtgcgggag ccagtcctgc 600
ttcaggctca ctgggacact cggccatttg gagtgtcatc cagccaaggg ttgtctctgc 660
acagtgttac ctctaaggat ctgtccgtgg tctgttagga actgggctgc acagccggag 720
gtgagcagct tcacagtat tacagccga cccattgct agcattactg cctgagctcc 780
gcctccatc agatcagcgg tggcattaga ttctcatagg agcttgaacc ctgttgtgaa 840
ctgcattgga gggatatagg atgtatgtc cttatgaaac tctaactaat gcctgatgat 900
ctgtcactgt ctcccatcac tcccagatgg gaccgtctag tctcaggaaa acaagctctg 960
ggctcccact gattctacat tatggtgtga tcctaggagc gccctggcc tccagctgcg 1020
caggagcctg tggatccagc ttcccagatg gcctcaccac tgagggaacc caggcctccg 1080
gggacaagga cattcctgca attaaccaag ggctcatcct ggaagaaacc ccagagagca 1140
gcttccctcat cgagggggac atcatccggc cgagtccctt ccgactgctg tcagcaacca 1200
gcaacaaatg gcccatgggt ggtagtgggt tcgtggaggt ccccttccgt ctctccagca 1260
agtacgatga gccagccgc caggtcatcc tggaggctct tgccggagttt gaacgttcca 1320
cgtgcatcag gtttgtcacc tatcaggacc agagagactt catttccatc atccccatgt 1380
atgggtgctt ctcgagtgtg gggcgagtg gagggatgca ggtgggtctcc ctggcgccca 1440
cgtgtctcca gaagggccgg ggcattgtcc ttcagtgtg catgcatgtg ctgggcttct 1500
ggcacgagca cacgcggggc gaccgggacc gctatatcca tgtcaactgg aacgagatcc 1560
tgccaggctt tgaaatcaac ttcataaggt ctcgagcag caacatgctg acgccctatg 1620
actactctc tgtgatgcac tatgggaggg tcccatgccc acagcactgg taggagcccc 1680
gctccggcct ccctatctct gcagcggctt ttggaggcac tgtcggcgga atccaggagc 1740
cccgaaccca gtggttccag tgccggaggc cagcccgttc ctgcagggcc tggggagagc 1800
ccacatgggt gggagtcccc tgccctgaaa aagctcagtg cagaggcctc ggcaaggcag 1860
ctcagaccc tggcttccct cccaagatca aggcctggag cagggtcccc cgggtgttct 1920
caggagcagt cctggctggc cggagtgtcc accaagccca cagtcccatc ttcagaagca 1980
ggaatccagc cagtccctgt ccagggaagc ccagctctgc cagggggctg tgtacctaga 2040
aatcatttca aggggatgtc cgaagattaa                                     2070

```

&lt;210&gt; 38

&lt;211&gt; 2265

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500264CB1

&lt;400&gt; 38

```
gggttcgggg gcgcccgcgt gtgaggccgg ggcctagagc cagccgcggc cgcgcaggag 60
gggcccaggg cccgcgctcg cccgcgtccc cgccttcttc cccgcgtcag ccccgccctcg 120
gctcgctgcc cttggctctc gtccgcatgg cctccgtcgc ccaggagagc gcgggctcgc 180
agcggcggtc accgcccgcg cacggggcgc tgcgcggggt gctactgctc tgcctgtggc 240
tgccaagcgg ccgtgcggcc ttgcccgcgc cggcgccggt gtccgaactg cacgcgcagc 300
tgtcggggcg ggagcagctg ctggaggagt tccgcccggc actgcagcag gagcgccctc 360
aggaggagct ggagctggag ctgcgcgcgg gcggcgggcc ccaggaggac tgcccggggc 420
cgggcagcgg cggctacagc gcaatgcctg acgcatcat ccgcaccaag gactccctgg 480
cggcggtgct cagcttcttc cgggcgcggc cggccgtgcg gggctggcgg caatgcgtgg 540
cgccctgctg ctccgagccg cgctgctccg tggccgtggg ggagctgccc cggcgccccg 600
cgccccgggc agccgtgctc ggctgctacc tcttcaactg cacggcgcgc ggccgcaacg 660
tctgcaagtt cgcgtgcac agcggtaca gcagctacag cctcagcgc gcgcccggacg 720
gcgcccgcct ggccaccgcg cgcgcctcgc cccggcagga aaaggatgcg cctccactta 780
gcaaggctgg gcaggatgtg gttctgcac tgcccacaga cggggtggtt ctagacggcc 840
gcgagagcac agatgaccac gccatcgctc agtatgagtg ggactgctg cagggggacc 900
cgtcagtgga catgaagggt cctcaatcag ggggtgactc cttggtggaa aagtctcaga 960
aagccactgc cccaaacaag ccacctgcat tatcaaacac agagaagagg aatcattccg 1020
ccttttgggg accagagagt caaatcattc ctgtgatgcc agatagtagt tcctcaggga 1080
agaacagaaa agaggaaagt tatatatatt agtcaaaggg tgatggagga ggaggggaac 1140
acccagcccc agaaacaggt gcagtgtac ccctggcgct gggtttggct atcactgctc 1200
tgctgcttct catggttgca tgccgactac gactggtgaa acagaaactg aaaaagctc 1260
gtcccattac atctgaggaa tcggactacc tcataaatgg gatgtatcta tagtaatgta 1320
atctcaatac cttggggcag ggacatgttt tgtttataat ttatacatct attaatgtct 1380
ggatatttac agcttctttt gtttttaatt gggccagaag attctgcaa tcccaaatct 1440
ttctttatta ttattgttaa aaaaagtctt cttagaagtc ataaatatt ttgaaattta 1500
gagaggaatt catgattaaa gattcctaaa aatataattc tgatttatgt aagctgtccc 1560
tgaaaataga aatgtgtact tagctgagag aaaattcagc atctcaggag gtggtattag 1620
gatgactgtg ttaacccatt accttttaga agccaaactgt tggccctta ccatgctgga 1680
ctgctatagg ccagcttcc ccttgtctg tggccctttt cttcctcctt gaagctccca 1740
gtattctttt tcttttcccc tctaaacctg tttctgagag tggatctcaa gcaagtcat 1800
gccttcaatc agatgttact tagggtgggt atacctaaat tataaacctt atgtacaagt 1860
cagtaagcct tagggaagggt gagtgtgggt ccttccta at cctctgacg tcatgtcata 1920
taggtggctg cctccttaga ctgacctttg ggagaaaaaa accccagact ttgaattagt 1980
aacagctcta agatggtcat gcagtgagat aggaaatcaa gatggaagca gagaatctgg 2040
catgccaaaa actaacagaa acttagttga aggcaaagag agcaaggaga acgtttaata 2100
cttcattaca tcaaatcaac actgctccat ggtgagagca cagcaactca tttatatata 2160
tatatatagg ctttgttgat gaaaaacgac aattgaagag aggacgttga gtggattcct 2220
gggtacagct tttgtaaaaa tgtcaccatg gctttcatcc aatgg 2265
```

&lt;210&gt; 39

&lt;211&gt; 1834

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7499935CB1

&lt;400&gt; 39

```

gtgacgccgg tgcggggcga acatggcggc ggccaccgga ccctcgtttt ggctggggaa 60
tgaaaccttg aaggtgccgc tggcgctctt tgccttgaac cggcagcgcc tgtgtgagcg 120
gctgcggaag aacctgctg tgcaggccgg ctccatcggt gtcctgcagg gcggggagga 180
gactcagcgc tactgcaccg acaccggggt cctcttcgcg caggagtcct tctttcactg 240
ggcgttcggt gtcactgagc caggctgcta tgggtgcatc gatgttgaca ctgggaagtc 300
gaccctgttt gtgccaggc ttctgcccag ccacgccacc tggatgggaa agatccattc 360
caaggagcac ttcaaggaga agtatgccgt ggacgacgtc cagtacgtag atgagattgc 420
cagcgtcctg acgtcacaga agcctctgtt cctcctcact ttgcgtggcg tcaacacgga 480
cagcggcagt gtctgcaggg aggcctcctt tgacggcatc agcaagtctg aagtcaacaa 540
taccattctt caccagaga tcgttgagtg ccgagtgttt aagacggata tggagctgga 600
ggttctgcgc tataccaata aaatctocag cgaggccac cgtgaggtaa tgaaggctgt 660
aaaagtggga atgaaagaat atgagttgga aagcctcttc gagcactact gctactcccg 720
ggcgggcatg cgccacagct cctacacctg catctgcggc agtggtgaga actcagccgt 780
gctacactac ggacacgccg gagctcccaa cgaccgaacg atccagaatg gggatatgtg 840
cctgttcgac atggggcggt agtattactg cttcgcttcc gacatcacct gctcctttcc 900
cgccaacggc aagttcactg cagaccagaa ggccgtctat gaggcagtgc tgcggagctc 960
ccgtgcggtc atgggtgcca tgaagccagg tgtctgggtg cctgacatgc accgcctggc 1020
tgaccgcatc cacctggagg agctggccca catgggcatc ctgagcggca gcgtggacgc 1080
catggtccag gctcacctgg gggccgtggt tatgcctcac gggcttggcc acttcctggg 1140
cattgacgtg cagcagctgg gaggtacccc agaggcggtg gagcgcatct acttcacga 1200
ccacctcctg gatgaggccc tggcggaacc ggcccgcgcc tccttcctta accgcgaggt 1260
cctgcagcgc tttcgcggtt ttggcggggt ccgcacgcag gaggacgtcg tggtgactga 1320
cagcggcata gagctgctga cctgcgtgcc ccgcactgtg gaagagattg aagcatgcat 1380
ggcaggctgt gacaaggcct ttacccctt ctctggcccc aagtagagcc agccagaaat 1440
cccagcgcac ctgggggcct ggccctgcaa cctcttttgc tgatgggcag cctgctggtc 1500
agcactccag tagcgagaga cggcaccacg aatcagatcc cagcttcggc atttgatcag 1560
accaaacagt gctgtttccc ggggagggaa cactttttta attacccttt tgcaggctcc 1620
cacctttaat ctgttttata ccttgcttat taaatgagcg acttaaaatg attgaaaata 1680
atgctgttct ttagtagcaa ctaaaatgtg tcttgctgtc atttatattc cttttcccag 1740
gaaagaagca tttctgatac tttctgtcaa aaatcaatat gcagaatggc atttgcaata 1800
aaaggtttcc taaaaaaaaa aaaaaaaaaa aaaa 1834

```

&lt;210&gt; 40

&lt;211&gt; 1524

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7982285CB1

&lt;400&gt; 40

```

gggacatata aaaatgccat tgtaactact gtagagtaaa gtgttagctg cgctgccgga 60
ggaaacggaa gaaggagcaa gctatggagg ggaacaggga tgaggctgag aaatgtgtcg 120
agatcgcccc ggaggccctg aacgccggca accgcgagaa ggcccagcgc ttctgcaga 180
aggccgagaa gctctaccca ctgccctcgg ccgcgcact attggaaata attatgaaaa 240
atggaagcac ggctggaaat agccctcatt gccgaaaacc atcaggtagt ggcgatcaaa 300
gcaagcctaa ttgcacaaag gacagcacat ctggtagtgg tgaagggtga aaaggctata 360
ccaaagacca agtagatgga gttctcagca taaacaaatg taaaaattac tatgaagtac 420
ttggagttac gaaagatgct ggtgatgaag atttgaaaaa agcttataga aagcttgctt 480
tgaagtttca tccagacaaa aaccatgcac ctggagcaac agatgctttt aaaaagattg 540
gaaatgctta tgctgtttta agtaatccag aaaagcgaac acagtatgac ctcacgggca 600
atgaagaaca agcatgtaac caccaaaaca atggcagatt taatttccat agaggttggt 660
aagctgatat aactccagaa gacttgttta atatattttt tgggggtgga tttccttcag 720
gtagtgtaca ttctttttca aatggaagag ctggttatag ccaacaacat cagcatcgac 780

```

```

atagtggaca tgaagagaa gaggaagag gagatggagg tttttctgtg tttatccagc 840
tgatgcccac aattgtattg atcctcgtgt cattattaag ccagttgatg gtctcctaac 900
ctccttattc cttatatccc agatctggaa ctgggcaaac tattaaaatg caaacagaaa 960
acttgggtgt tgtttattat gtcaacaagg acttcaaaaa tgaatataaa ggaatgttat 1020
tacaaaaggt agaaaagagt gtggaggaag attatgtgac taatattcga aataactgct 1080
ggaaagaaag acaacaaaaa acagatatgc agtatgcagc aaaagtatac cgtgatgatc 1140
gactccgaag gaaggcagat gccttgagca tggacaactg taaagaatta gagcggctta 1200
ccagtcttta taaaggagga tgaactggaa tttttattta taccttttag cgtactcttt 1260
attttttctg taagtaagtt tggtttcac atgagggatg aaggaaaaga tttgatactg 1320
aaaactaaac tgaatagttg gttcctgaaa tcttggactg tttatgacct actggctcct 1380
ttaaatagta actgaaaact aaaaaggaa attttagtta acgcttctac aagtattttc 1440
attttaaaag cttacatgat tcctaaacta aagtgtcatg agaaaggatt atcacacctg 1500
tagcaatttc cagtttttagt gatt                                     1524

```

&lt;210&gt; 41

&lt;211&gt; 2973

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7758505CB1

&lt;400&gt; 41

```

cgagcggggt ggctcgacca aggagtgggt gtgggtgggt ctttcggggc tcccggcgac 60
ccgcggtgcc gacgcaacgc cgacgtatgg tgccaagcga actttaaaaa gctgcttcgg 120
acaaaccaga gccaggatct cactgtcgg ggaccggga tcggaagggt ctagcccgag 180
ggaaatgctg gaagatccca tcggccagtg accagcaact ttccggcgag attttgacgc 240
ggagaactgt gctctgcctc ctcttattct cccaaagctc acgttggcgt cctgccttgc 300
gggggaactc ggcgcgctct ctgectgagc agcagtgaa ttgaaccca gccgcctccg 360
gcgcctccgg gctgatgagt gtcgctctcc gcccgccat ctctttttcc cggaggtaaa 420
ggcccgcggt ccccaacctt cagtgcgcc gggtccaag gcccgagcc agcgttttg 480
cggagccgct tcttggatgc tgaaggctgg gctcctccat cgtgggtgcc gaggcggcga 540
tgggtgtcct caaagtgtgg ctcggtctgg ccctagcgtt ggcggaattt gcagtattgc 600
ctcatcatc cgaagggtgt tgtgtctatc aggatctcct gttggcggt gccacaattt 660
ggaagcccca ttcatgccag agctgccgtt gccatggtga tattgttata tgcaaacctg 720
ctgtttgcag aaacctcaa tgtgcctttg agaagggaga agtgcttcaa atagctgcca 780
accaatgctg tcctgagtggt gttttgagga ctccaggatc ttgccatcat gaaaagaaaa 840
tccatgagca tgggacagaa tgggcctctt ctccatgtag tgtgtgctct tgcaatcatg 900
gggaagtccg atgtaccccc caaccatgcc caccgtgtc atgtggacac caggagctgg 960
cattcatccc tgaagggaagc tgctgccag tttgtgtggg ccttgggaaa ccctgttctc 1020
atgaaggcca tgtgtttcag gatggggagg actggcggt gagccggtgt gccaaatgtc 1080
tgtgtagaaa tgggggttgc cagtgttca cagctcagtg tcagcctcta ttttgtaacc 1140
aggatgagac tgtagtccga gtccctggaa aatgttgccc gcagtgtctt gcaagatcct 1200
gctctgcagc tggccaagta tacgagcatg gtgagcagtg gagcgaatat gcctgcacca 1260
cgtgtatatg tgaccggggt gaggtcaggt gtcacaagca ggctgcctg ccctgagat 1320
gcggaagggt tcagagcagg gctcggcgtc atgggcaatg ctgtgaggaa tgtgtgtctc 1380
ctgcccggag ctgtcctat gatggagtgt tgcggtacca ggacgaaatg tggaagggt 1440
cggcctgtga gttctgctga tgtgatcat gccaaagtgc ctgccagact ggagagtgtg 1500
ccaaagtga gttgtgcccg gatgaagaat taattcact agatggaaag tgtgtctc 1560
aatgcatttc aaggaaatggt tattgtgttt atgaagaaac tggagaattt atgtcatcaa 1620
atgctagtga agttaaacgt attccagagg gagagaagt ggaagatggc ccttgcaagg 1680
tgtgtgagtg ccgagggtgt caggtaactt gctacgagcc ctcttgccca ccatgtccag 1740
tgggcacact ggccttagag gtgaagggtc agtgctgtcc agactgcaca tcagttcatt 1800
gccatccaga ttgtttgaca tgctctcagt ctccagacca ctgtgacctc tgccaagatc 1860
ctaccaagtt actgcagaat ggatggtgtg tgcacagctg tggactgggt ttttaccagg 1920

```



```

ctggcagtct ctgtatagcc tgccagcccc agtgetccac gtgtaccagt gggctggagt 1980
gctcatcctg ccagcctccc ctgctgatgc ggcacgggca gtgtgtgcct acctgtgggg 2040
acggcttcta ccaagatcgc cattcctgtg cagtctgcc tgaagtctgt gcaggttgct 2100
ggggcccaac ggagaagcac tgcttggcct gcagagatcc cctccacgtg ctgagagatg 2160
gcggtgtga gagcagctgt ggaaaaggct tctacaacag gcagggcacc tgtagcgctt 2220
gtgaccaatc ctgtgacagt tgtggcccca gtagccccag gtgtcttacc tgtactgaga 2280
agacagtgtc gcatgatggg aaatgcatgt ctgaatgccc tggcgggtac tatgctgatg 2340
ccactggcag gtgcaaagtt tgtcataact catgtgccag ctgctctggg cccacaccct 2400
ctcactgtac agcctgcagc cccccaagg ctctgcgtca aggccactgt ctgccccgct 2460
gtggagaggg tttctactct gaccacggag tctgcaaagc ctgtcactcc tcctgcctgg 2520
cttgatggg tcccgacccc tctcactgta ctgggtgtaa gaagccagag gaaggactgc 2580
aagtggagca gctgtctggc gtgggcatcc cctctggcga gtgtctagcc cagtgtagag 2640
ccatttttta cttggagagc actggcctat gtgaagggca aaatctggac ttctgtcaga 2700
athtagaaat gattttctgt gtttgccttg gcatatcatc tacagagaat tgatgacatc 2760
ctgaataaat aatttgactc aatagccagg ccatctatga gtggttgagg agatgaaagg 2820
gaagtattat agtttccttt ctgttccac aagtagcctt gctgttgggt gaatagtgtt 2880
actctaaagc tacgtgaaaa aaaaatcatt agtttgtatt ttctattgta aacatatgtt 2940
cattaaaaaa attttataat acaccacta cct 2973

```

<210> 42

<211> 2126

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 6885756CB1

<400> 42

```

gaggagtaat ttgattcagg tgttctagaa gtcattgatgt gggctgtgtc tgttgaattc 60
ccagcgatgc aaggggacac accctgtgac tcattcctta attgagtgtc gatatttgat 120
tggtttatcg cacacctgat ggggtgggtgg ggtgttcgag gttggagggg gtgagttata 180
taagggtga tgcgccaga gagctggtca tttgaagact ctctcggaag agatagcgtc 240
ttgtgcaac ctgcggtccc agcagaaaaa ccttgtgac cttgttgcgg gcgacatgga 300
agacgactca ctctatttgg gaggtgactg gcagttcaat cacttttcaa aactcacatc 360
ttctcggtca gatgcagctt ttgctgaaat ccagcggact tctctctctg aaaagtcacc 420
actctcatct gagaccgtt tcgacctctg tgatgatttg gctcctgtgg caagacagct 480
tgctccagg gagaagcttc ctctgagtag caggagacct gctgcggtgg gggctgggct 540
ccagaagata ggaaatacct tctatgtgaa cgtttcctcg cagtgcctga catacact 600
gccgctttcc aactacatgc tgtcccgga ggactctcaa acgtgtcatc ttcacaagt 660
ctgcatgttc tgtactatgc aagctcacat cacatgggccc ctctaccgtc ctggccatgt 720
catccagccc tcacaggtat tggctgctgg ctcccataga ggtgagcagg aggatgccc 780
tgaatttctc atgtttactg tggatgccat gaaaaaggca tgccttccc ggcacaagca 840
gctagatcat cactccaagg acaccacct catccaccaa atatttgag cgtattggag 900
atctcaaatc aagtatctcc actgccacgg catttcagac acctttgacc cttacctgga 960
catcgccctg gatatccagg cagctcagag tgtcaagcaa gctttggaac agttgggtgaa 1020
gcccagaaga ctcaatggag agaatgccta tcattgtggt ctttgtctcc agaaggcgcc 1080
tgctccaag acgttaactt taccacttc tgccaaggct ctcatcttg tattgaagag 1140
attctccgat gtcacaggca acaaacttgc caagaatgtg caatataccta agtgccgtga 1200
catgcagcca tacatgtctc agcagaacac aggacctctt gtctatgtcc tctatgctgt 1260
gctggtccac gctgggtgga gttgtcacia cggacattac ttctcttatg tcaaagctca 1320
agaaggccag tgggtataaaa tggatgatgc cgaggctact gcctctggca tcacctctgt 1380
cctgagtcaa caggcctatg tcctctttta catccagaag agtgaatggg aaagacacag 1440
tgagagtgtg tcaagaggca gggaaccaag agcccttgggt gctgaagaca cagacaggcc 1500
agcaacgcaa ggagagctca agagagacca cccttgcttc caggtagccg agttggacga 1560
gcacttggtg gaaagagcca ctcaggaaag caccttagac cactggaaat tccccaaaa 1620

```

```

gcaaaacaaa acgaagcctg agttcaacgt cagaaaagtt gaaggtaccc tgcctcccaa 1680
cgtacttgtg attcatcaat caaaatacaa gtgtggtatg aaaaaccatc atcctgaaca 1740
gcaaagctcc ctgctaaacc tctcttogac gaaaccgaca gatcaggagt ccatgaacac 1800
tggcacactc gcttctctgc aaggaggcac caggagatcc aaagggaata acaaacacag 1860
caagagatct ctgcttgtgt gccagtgtac acagtgggaag taccgaccca cactgagggg 1920
tgcacacaca cacacacaca caaacacaaa tacaccacaca agcgcgcacg gaaacacaca 1980
cacaccacaca caaacacgaa caccgtcaat cctacataaa gtaatgagga gccccagttt 2040
ctgtctctac aacagggaca attggatagt gatggctgcg tctcaggatg agcccacaca 2100
tgggaaacat caagttgggg gttcag 2126

```

<210> 43  
 <211> 1973  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7500748CB1

```

<400> 43
ggaagtagcc gcaggcatgg cggcggctat gccgcttgct ctgctcgtcc tgttgcctcc 60
ggggcccgcc ggctggtgcc ttgcagaacc cccacgcgac agcctgcggg aggaacttgt 120
catcaccccg ctgccttcog gggacgtagc cgccacatcc cagttccgca cgcgctggga 180
ttcggagctt cagcgggaag gagtgtccca ttacaggctc tttcccaaag ccctggggca 240
gctgatctcc aagtattctc tacgggagct gcacctgtca ttcacacaag gcttttggag 300
gacccgatac tggggggccac ccttcctgca ggcccatca ggtgcagagc tgtgggtctg 360
gttccaagac actgtcactg atgtggataa atcttgggaag gagctcagta atgtcctctc 420
agggatcttc tgcgcctctc tcaacttcat cgactccacc aacacagtca ctcccactgc 480
ctccttcaaa cccctggggtc tggccaatga cactgaccac tactttctgc gctatgctgt 540
gctgcccgcc gaggtgggtct gcaccgaaaa cctcaccccc tggaagaagc tcttgccctg 600
tagttccaag gcaggcctct ctgtgctgct gaaggcagat cgcttggtcc acaccagcta 660
ccactcccag gcagtgcata tccgcoctgt ttgcagaaat gcacgctgta ctagcatctc 720
ctgggagctg aggcagaccc tgtcagttgt atttgatgcc ttcatcacgg ggcagggaaa 780
gaaagactgg tccctcttcc ggatgttctc ccgaaccctc acggagccct gccccctggc 840
ttcagagacc cgagtctatg tggacatcac cacctacaac caggacaacg agacattaga 900
ggtgcaccca ccccgcacca ctacatatca ggacgtcatc ctaggcactc ggaagacctc 960
tgccatctat gacttgcttg acaccgccat gatcaacaac tctcgaaaac tcaacatcca 1020
gctcaagtgg aagagacccc cagagaatgg ttacatccac taccagcctg cccaggaccg 1080
gctgcaaccc cacctcctgg agatgctgat tcagctgcog gccaaactcag tcaccaaggt 1140
ttccatccag tttgagcggg cgctgctgaa gtggaccgag tacacaccag atcctaacca 1200
tggcttctat gtcagcccat ctgtcctcag cgcccttggt cccagcatgg tagcagccaa 1260
gccagtggac tgggaagaga gtccctctt caacagcctg ttcccagctc ctgatggctc 1320
taactacttt gtgcggctct acacggagcc gctgctgggt aacctgccga caccggactt 1380
cagcatgccc tacaacgtga tctgcctcac gtgcactgtg gtggcctgtg gctacggctc 1440
cttctacaat ctctcaccoc gaaccttcca catcgaggag ccccgcacag gtggcctggc 1500
caagcggctg gccaacctta tccggcgcgc ccgaggtgtc ccccaactct gattcttggc 1560
ctttccagca gctgcagctg ccgtttctct ctggggaggg gagcccaagg gctgtttctg 1620
ccacttgctc tcctcagagt tggcttttga accaaagtgc cctggaccag gtcagggcct 1680
acagctgtgt tgtccagtac aggagccacg agccaaatgt ggcatttgaa tttgaattaa 1740
cttagaaatt catttctca cctgtagtgg ccacctctat attgaggtgc tcaataagca 1800
aaagtggctg gtggctgctg tattggacag cacagaaaaa gatttccatc accacagaaa 1860
ggtcggctgg cagcactggc caaggtgatg ggggtgtgcta cacagtgtat gtcactgtgt 1920
agtggatgga gtttactgtt tgtggaataa aaacggctgt ttccgtggaa aaa 1973

```

<210> 44  
 <211> 1884

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500749CB1

&lt;400&gt; 44

```
ggcatggcgg cggctatgcc gcttgctctg ctcgctcctgt tgctcctggg gccgggaggc 60
tggtgccttg cagaaccccc acgcgacagc ctgcgaggagg aacttgatcat ccccccgctg 120
ccttcggggg acgtagccgc cacattccag ttccgcacgc gctgggattc ggagcttcag 180
cggaaggag acactgacca ctactttctg cgctatgctg tgctgccgcg ggaggtggtc 240
tgcaccgaaa acctcacccc ctggaagaag ctcttgccct gtagttccaa ggcaggcctc 300
tctgtgctgc tgaaggcaga tcgcttgctc cacaccagct accactccca ggcagtgcac 360
atccgccctg tttgcagaaa tgcacgctgt actagcatct cctgggagct gaggcagacc 420
ctgtcagttg tatttgatgc cttcatcacg gggcagggaa agaaagactg gtccctcttc 480
cggatgttct cccgaacctt caccggagccc tgccccctgg cttcagagag ccgagtctat 540
gtggacatca ccacctacaa ccaggacaac gagacattag aggtgcaccc acccccgacc 600
actacatatc aggacgtcat cctaggcact cggaagacct atgccatcta tgacttgctt 660
gacaccgcca tgatcaacaa ctctcgaaac ctcaacatcc agctcaagtg gaagagacc 720
ccagagaatg agggcccccc agtgcccttc ctgcatgccc agcggtagct gagggtgctat 780
gggctgcaga agggggagct gagcacactg ctgtacaaca cccaccata ccgggccttc 840
ccggtgctgc tgctggacac cgtaccctgg tatctgcggc tgtatgtgca caccctcacc 900
atcacctcca agggcaagga gaacaaacca agttacatcc actaccagcc tgcccaggac 960
cggctgcaac cccacctcct ggagatgctg attcagctgc cggccaactc agtcaccaag 1020
gtttccatcc agtttgagcg ggcgctgctg aagtggaccg agtacacacc agatcctaac 1080
catggcttct atgtcagccc atctgtcctc agcgcccttg tgcccagcat ggtagcagcc 1140
aagccagtgg actgggaaga gagtccctc ttcaacagcc tgttccagct ctctgatggc 1200
tctaactact ttgtgaggct ctacacggag ccgctgctgg tgaacctgcc gacaccggac 1260
ttcagcatgc cctacaacgt gatctgcctc acgtgcactg tggtagcctg gtgctacggc 1320
tccttctaca atctcctcac ccgaaccttc cacatcgagg agccccgcac aggtggcctg 1380
gccaagcggc tggccaacct tatccggcgc gcccgagggtg tccccccact ctgattcttg 1440
ccctttccag cagctgcagc tgccgtttct ctctggggag gggagcccaa gggctgtttc 1500
tgccacttgc tctcctcaga gttggctttt gaaccaaaagt gccctggacc aggtcagggc 1560
ctacagctgt gttgtccagt acaggagcca cgagccaaat gtggcatttg aatttgaatt 1620
aacttagaaa ttcatttcct cacctgtagt ggccacctct atattgaggt gctcaataag 1680
caaaagtggc cgggtggctgc tgtattggac agcacagaaa aagatttcca tcaccacaga 1740
aaggctggct ggcagcactg gccaagggtga tggggtgtgc tacacagtgt atgtcactgt 1800
gtagtggtat gagtttactg tttgtggaat aaaaacggct gtttccgtgg aaaaaaaaaa 1860
aaaaaaaaaa aaaaaaaaaa aaag 1884
```

&lt;210&gt; 45

&lt;211&gt; 1581

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503401CB1

&lt;400&gt; 45

```
cgccaccacc tcagctgcgg accgaggcga gatggcggcc accgaggggg tcggggaggc 60
tgcgcaaggg ggcgagcccg ggcagccggc gcaacccccg cccagccgc acccaccgcc 120
gccccagcag cagcacaagg aagagatggc ggccgaggct ggggaagccg tggcgtcccc 180
catggacgac gggtttgtga gcctggactc gccctcctat gtctgtaca gggacagagc 240
agaatgggct gatatagatc cgggtgccga gaatgatggc cccaatcccc tgggtccagat 300
```

```

catttatagt gacaaattta gagatgttta tgattacttc cgagctgtcc tgcagcgtga 360
tgaaagaagt gaacgagctt ttaagctaac ccgggatgct attgagttaa atgcagccaa 420
ttatacagtg tggcatcata ggcgagtatt agtggaaatgg ctaagagatc catctcagga 480
gcttgaattht attgctgata ttcttaatac ggatgcaaag aattatcatg cctggcagca 540
tcgacaatgg gttattcagg aatttaaaact ttgggataat gagctgcagt atgtggacca 600
acttctgaaa gaggatgtga gaaataaact tgtctggaac caaagatact tcgttatttc 660
taacaccact ggctacaatg atcgtgctgt attggagaga gaagtccaat acactctgga 720
aatgattaaa ctagtaccac ataataaag tgcatggaac tatttgaaag ggattttgca 780
ggatcggtgt ctttccaaat atcctaactc gttaaatcaa ttacttgatt tacaaccaag 840
tcatagttcc cctacctaata ttgcctttct tgtggatata tatgaagaca tgctagaaaa 900
tcagtgtgac aataaggaag acattcttaa taaagcatta gagttagtg aaatcctagc 960
taaagaaaag gacactataa gaaaggaata ttggagatac attggaagat cccttcaaag 1020
caaacacagc acagaaaatg actcaccaac aaatgtacag caataacacc atccagaaga 1080
acttgatgga atgctttttat tttttattaa gggaccctgc aggagtttca cacgagagtg 1140
gtccttccct ttgcctgtgg tgtaaaagtg catcacacag gtattgcttt ttaacaagaa 1200
ctgatgctcc ttgggtgctg ctgctactca gactagctct aagtaatgtg attcttctaa 1260
agcaaagtca ttgatggga ggaggaagaa aaagtcccat aaaggactt ttgtagtctt 1320
atcaacatat aatctaatac cttagcatca gctcctccct cagtggtaga tgcgtcaaga 1380
tttgtagcag taataactgc aggtcacttg tatgtaatgg atgtgaggta gccgaagttt 1440
ggttcagtaa gcagggaata cagtcgttcc atcagagctg gtctgcacac tcacattatc 1500
ttgctatcac tgtaaccaac taatgccaaa agaacggttt tgtaataaaa ttatagctgt 1560
atctaaaaaa aaaaaaaaaa g 1581

```

&lt;210&gt; 46

&lt;211&gt; 1996

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503485CB1

&lt;400&gt; 46

```

gttcagcccc cgtctacact ggggtgggtgc ttagccggcg ccagaccgac cctcgacttc 60
ggagaggcag cgcgggttcc ctgggtgctt ccgcctcccc ttctcctgct tctccagcct 120
cttcggcctc ctgcgccgcc gcgggaaccc gagaccccag tgtatgcccc acccctgacc 180
ccgctcgcga catgtccacc ccggtcgggc ggcgctcat gcgggacttc aagagggttg 240
aggaggatcc tccagccgga gtcagcgggg ctccgtccga gaacaacata atggtgtgga 300
acgcgggtcat tttcgggcct gaagggaccc cgtttgagga tgtctatgca gatggttagta 360
tatgtctgga catacttcag aaccgttgga gtccaacct tgaatgtgtc tccattctaa 420
catccatata gtctctgttg gatgaaccca atcccaatag tccagcaaac agccaggctg 480
ctcagctgta ccaggagaac aaacgggaat atgaaaagcg tgtttctgca atagtagaac 540
aaagctggcg tgattgttga cccgggttac agtttaaaga agctggccat aagaaaaata 600
tatattgatg tgtttgtcac ctccctactc ctgtcattac atttacttta ttaaaagcaa 660
aataactgtt gtgctgtttc catcttccct gccaaagttt cctaccctct ctaccctctc 720
cttaaacatc agaaaacacc ctctatgaaa tcaaatgtac tgtacctggg ttacttgcaa 780
aaattactaa tgcttcagtt tttctgttgt atttcatttc cagttttcag gcagttattt 840
ttattattgt actttaagct ttaaatgata attgttatat aagagggtgt tatgcttagc 900
ttgatgacca ggatgttatt ttaacaaaa tgattgctga agtgtttcat cctggctggt 960
ccttcacttg tgttgattt agaagtgaat gtgtttggaa tatggcctac agagaataga 1020
aacaatacca tgtaaacatc ttgaaggag gcatgggagc taaaaatcct gtgatactaa 1080
gatctcagtc atatgaatta caacgtagta tttactggca agaaggagaa agttgaagga 1140
ctcagctaaa ggagtacagc aattgtagta actgacacat cctctctttg caagctgctg 1200
actgggcaca ctcatgccaa gtttcagaat tattggtctt ctgggttttt gcttttttaa 1260
agagggtgtg gagcagagga atggaaacaa tctgtagttt ttgagctagg gaaagttgga 1320
gctcctttta tcttttttaa ggatcagtg ctcctaaagt gaataaaact aattgtccat 1380

```

```

ctttatttta gagttttaat gaattcaagg aaggaggcat agcatatctg tggcaaaacta 1440
ttttccactc aaatcctgag ttattgctgc atgctttaat ttcttccctt tcagcatctg 1500
agaaccttaa agccaatgtc tgcgatcttt ttttggatat ttatactttt agatatatag 1560
tacctttaag tagcagtatg ggacaaggct tgtaaagtgt ttgtctaagt ttctattgtc 1620
accttttatg cttttatcac ttccaaatct aactttgcac aagtaacca tgtaaaaaaa 1680
aaatgtacat ttttcaaaag ttgtaaataa aaataacctt aaaatttcaa aaaaaaaaaa 1740
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1800
aaaaaaaaatt ttttcttgtg ggacaaaaaa aaaaaagcgg cgggtgtgtgt gtgttttaat 1860
agccaaacaa aatttttttt cgggtatattt tgggggggtgg ccccccttt tttagcaaaa 1920
tgagaaaaac tgtacacgat gtataaacgg cggaggagga ataaaaatat ttttatcaaa 1980
aagaaaaaaa ttatca                                     1996

```

&lt;210&gt; 47

&lt;211&gt; 1232

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7504076CB1

&lt;400&gt; 47

```

ggcaagcgca ggtcaggcgg tgcagctggg cggccagcgg atcgtgccgc ggcgcccgag 60
cgcagctaca ggagggtgtc cagaagccac aagccatggc tgtggggaac atcaacgagc 120
tgcccgagaa catcctgtct gagctgttca cgcacgtgcc cgcccgccag ctgctgctga 180
actgccgcct ggtctgcagc ctctggcggg acctcatcga cctcgtgacc ctctggaaac 240
gcaagtgcct gcgagagggc ttcatcactg aggactggga ccagcccgtg gccgactgga 300
agatcttcta cttcttacgg agcctgcaca ggaacctcct gcacaacccg tgcgctgaag 360
aggggttcga gttctggagc ctggatgtga atggaggcga tgagtggag gtggaggatc 420
tctctcgaga ccagaggaag gaattcccca atgaccaggt caagaaatac ttcgttactt 480
catattacac ctgcctcaag tcccagggtg tggacctcaa ggccgaaggg tattgggagg 540
agctgatgga taccacacgg ccggacatcg aggtcaagga ctggttcgca gccaggccag 600
attgcggttc caagtaccag ctgtgcgttc agctcctgtc gtccgcgcac ggcctctg 660
ggaccttcca gccagaccgg gcgaccatcc agcagaagag cgatgccaa tggaggaggg 720
tctccacac attctccaac taccgccccg gcgtccgcta catctggttt cagcacggcg 780
gcttgacac tcattactgg gccggctggt acggccccg ggtcaccaac agcagcatca 840
ccatcggggc cccgctgccc tgacaccccc tgagccccca tctgctgaac cctgactgct 900
ttacggacat tggatgaagc cgaagcattt agaatgggtc ctggcacaca gttggtgcgt 960
gatatgggta agcttttgtt ccccaccac atctcatctt gaatgtgac gtttcccccg 1020
ctccctcctg ccgccatgtg aagaaggtcg ttgcttcccc ttcacctcc accaccatga 1080
tttagagatg gagtttcacc atgattggac acagggtggt ctcaatctcc tgaacctcgt 1140
gatccacca cctcgacctc ccatagtgtc gagattaaca tggcgtggcc accgcgctct 1200
accgcttgtg tcttgacggc tcccagcct ca                                     1232

```

&lt;210&gt; 48

&lt;211&gt; 810

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500926CB1

&lt;400&gt; 48

```

ggggtccgga actgcttgtt ccggcagtg aagagacggc ccggcggttg ccgctgctgc 60
tagcagcttg aaccccgagg tcgggaccga tgtcggcttg ggctgctgcc agcctaagca 120

```

```

gggccgctgc cccgatgcttg ctggcacgag gccccggggt cagggcggt cctccgcgcg 180
acccccggcc ctcccacccc gagccccggg gctgcggtgc cgctccgggc aggacgctgc 240
actttaccgc ggctgtcccc gccgggcaca acaagtgggc caaagtcagg cacatcaagg 300
gtccgaagga cgtcgaaagg agtcgcatct tctccaaact ctgtttgaac atccgcctgg 360
cagtgaagga aggaggcccc aaccctgagc acaacagcaa cctggccaat atcttagagg 420
tgtgtcgcag caaacatatg cccaagtcaa cgattgagac agcactgaaa atggagaaat 480
ccaaggacac ttatttgctg tatgagggtc gaggccctgg tggctcttct ctgctcatcg 540
aggcattatc taacagtagc cacaagtgcc aagcagactt gcgacctga agccaaagga 600
atctcacttg tggggcctcc ttgtcagctc tgctgctgctc tcagagccat ctggatgagt 660
gtcccagcac cctctcggat gcagggcagg accaccagc tggtcagact ctgatgttgg 720
gtagctggcc tctgtgggga ttgtaagtgc cctgaggcgc tctgtactag aaactgctct 780
taatagtaac ggtgattatt ggttgctgca 810

```

&lt;210&gt; 49

&lt;211&gt; 2625

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503216CB1

&lt;400&gt; 49

```

gcggcggcgg cagcagcagc agcagcatta gcagcagcag cttctctcaa tgctgagtag 60
tatatactca agataaatga aaacacatgt ccacacaaaa gcttgctcat gaattgtcat 120
agaaacattc ttcacaatat ccaaaagtgg aagcaacca aatgtccatc aactgatgaa 180
tggaataaaca aagtaatatt ctgtacgatg gaattttaat tggcataattt ggtaataaaa 240
aggaatcaag tactgataca tcctacaaca tggatacatc ttgaaaatgc tatgctaaat 300
gaaagagggt gaatcaccag gattttcaac tgagaaattt aagaataatt gaacctaacg 360
aggtgacaca ctacaggagc acaggtgtgg aaacagacgg cagaatgcct ccaaagggtga 420
cttcagagct gcttcggcag ctgagacaag ccatgaggaa ctctgagtat gtgaccgaac 480
cgatccaggc ctacatcatc ccacgggag atgtctcatc gagtgagtat attgctccat 540
gtgactgtcg gcgggctttt gtctctggat tcgatggctc tgcgggcaca gccatcatca 600
cagaagagca tgcagccatg tggactgacg ggcgctactt tctccaggct gccaaagcaa 660
tggaagcaaa ctggacactt atgaagatgg gtctgaagga cacaccaact caggaagact 720
ggctggtgag tgtgtctcct gaaggatcca gggttggtgt ggacccttg atcattccta 780
cagattattg gaagaaaatg gccaaagtgc tgagaagtgc cggccatcac ctattcctg 840
tcaaggagaa cctcgttgac aaaatctgga cagaccgtcc tgagcgccct tgcaagcctc 900
tcctcacact gggcctggat tacacagggc tattttaatct ccgaggatca gatgtggagc 960
acaatccagt atttttctcc tacgcaatca taggactaga gacgatcatg ctcttcattg 1020
atggtgaccg catagacgcc ccagtggtga aggagcacct gcttcttgac ttgggtctgg 1080
aagccgaata caggatccag gtgcatccct acaagtccat cctgagcgag ctcaaggccc 1140
tgtgtgctga cctctcccca agggagaagg tgtgggtcag tgacaaggcc agctatgctg 1200
tgagcgagac catccccaag gaccaccgct gctgtatgcc ttacaccccc atctgcatcg 1260
ccaaagctgt gaagaattca gctgagtcag aaggcatgag gcgggctcac attaaagatg 1320
ctgttgctct ctgtgaactc tttaactggc tggagaaaga ggttcccaaa ggtggtgtga 1380
cagagatctc agctgtgac aaagctgagg agtttcgcag gcaacaggca gactttgtgg 1440
acctgagctt occaacaatt tcagtacgg gaccacacgg cgccatcatt cactacgcgc 1500
cagtcctga gacgaatagg accttgctcc tggatgaggt gtacettatt gactcgggtg 1560
ctacatacaa ggatggcacc acagatgtga cgcggacaat gcattttggg acccctacag 1620
cctacagagaa ggaatgcttc acatatgtcc tcaagggcca catagctgtg agtgacggc 1680
ttttcccgac tggaaacaaa ggtcaccttc ttgactcctt tgcccgttca gctttatggg 1740
attcaggcct agattacttg cacgggactg gacatggtgt tgggtctttt ttgaatgtcc 1800
atgagggtcc ttgcggcatc agttacaaaa cattctctga tgagcccttg gaggcaggca 1860
tgattgtcac tgatgagccc ggggtactatg aagatggggc ttttggaatt cgcattgaga 1920
atgttgcctc tgtggttcct gtgaagacca agtataattt taataaccgg ggaagcctga 1980

```

```

cctttgaacc tctaacattg gttccaattc agacccaaat gatagatgtg gattctctta 2040
cagacaaaga gtgcgactgg ctcaacaatt accacctgac ctgcagggat gtgattggga 2100
aggaattgca gaaacagggc cgccaggaag ctctcgagtg gctcatcaga gagacgcaac 2160
ccatctccaa acagcattaa taaatacctc cccggttttg tttttgtaa atgctctgga 2220
ggaaggaaga aacgtggcag atccctgaca tctttccctt ttcctttcct tcttccctac 2280
ctcccccttt tacttttagac ttttaagaaga acagaaaatc ttcttatcct ctttgatatt 2340
ttattgcaaa cactcagtct tttatgattt ttttaattgtt gagaacaagc caagaataaa 2400
attgctgcac cagaaggagg gtccctccaa agttgaacac ttggtgaaag gaagatgccc 2460
cgacttcttt ggccagtgat ggggaatcag tgagtgtctc atgatgggtc tggtccagggt 2520
gctagtacat cattcatgat caccttaatg ctcatgagac tatatttatg atcagtgaat 2580
aaaaatgtca gaactgtgaa aaaaaaaaaa aaaaaaaaaa aaaag 2625

```

&lt;210&gt; 50

&lt;211&gt; 2432

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503233CB1

&lt;400&gt; 50

```

gcgctcttcc tgggtggggc ctgcccctgag ctgccaccgg gaagccagcc tcaggggactg 60
cagcgacccc caaacacccc tccccagga tgctcgaggat gatcatcacg ccggtgtact 120
gcaactgggt gtcagcccaa gtgcagaagc agcgggcccag ggagctgggc ctgggcccgc 180
atgagaatgc catcaagtac ctggggccagg attatgagca gctgcgggtg cgatgcctgc 240
agagtgggac cctcttccgt gatgaggcct tccccccggt accccagagc ctggggttaca 300
aggacctggg tcccaattcc tccaagacct atggctatgc cggcatcttc catttccagc 360
tgtggcaatt tggggagtgg gtggacgtgg tctgggatga cctgctgccc atcaaggacg 420
ggaagctagt gtctgtgcac tctgccgaag gcaacgagtt ctggagcgcc ctgcttgaga 480
aggcctatgc caaggtaa atggcagctac aggcctgtc agggggcagc acctcagagg 540
gctttgagga cttcacaggc ggggttacgg agtggtacga gttgcgcaag gctcccagtg 600
acctctacca gatcatctc aaggcgctgg agcggggctc cctgctgggc tgctccatag 660
acatctccag cgttctagac atggaggcca tcactttcaa gaagtgggtg aaggggccatg 720
cctactctgt gaccggggcc aagcaggtga actaccgagg ccagggtggg agcctgatcc 780
ggatgcggaa cccctggggc gaggtggagt ggacgggagc ctggagcgac agctcctcag 840
agtggaaaca cgtggaccca tatgaacggg accagctccg ggtcaagatg gaggacgggg 900
agttctggat gtcattccga gacttcatgc gggagttcac cgcctggag atctgcaacc 960
tcacacccga cgcctcaag agccggacca tccgcaaatg gaacaccaca ctctacgaag 1020
gcacctggcg gcggggggagc acccgggggg gctgccgaaa ctaccagcc accttctggg 1080
tgaacctca gttcaagatc cggctggatg agacggatga cccggacgac tacgggggacc 1140
gcgagtcagg ctgcagcttc gtgctcggcc ttatgcagaa gcaccgtcgc cgcgagcgcc 1200
gcttcggccg cgacatggag actattggct tcgcggtcta cgaggctcct ccggagctgg 1260
tgggccagcc ggccgtacac ttgaagcgtg acttcttcc tggccaatgcg tctcggggcg 1320
gctcagagca gttcatcaac ctgcgagagg tcagcaccgg cttccgcctg ccaccggggg 1380
agtatgtggt ggtgccctcc accttcgagc ccaacaagga gggcgacttc gtgctgcgt 1440
tcttctcaga gaagagtgt gggactgtgg agctggatga ccagatccag gccaatctcc 1500
ccgatgagca agtgctctca gaagaggaga ttgacgagaa cttcaaggcc ctcttcaggc 1560
agctggcagg ggaggacatg gagatcagcg tgaaggagtt gcggacaatc ctcaatagga 1620
tcacagcaa acacaaagac ctgcggacca agggcttcag cctagagtcg tgccgcagca 1680
tggtgaacct catggatcgt gatggcaatg ggaagctggg cctgggtggg ttcaacatcc 1740
tgtggaaccg catccggaat tacctgtcca tcttccggaa gtttgacctg gacaagtcgg 1800
gcagcatgag tgcctacgag atgcggatgg ccattgagtc ggcaggcttc aagctcaaca 1860
agaagctgta cgagctcatc atcaccgct actcggagcc cgacctggcg gtcgactttg 1920
acaatttcgt ttgctgcctg gtgcggctag agaccatgtt ccgatttttc aaaactctgg 1980
acacagatct ggatggagtt gtgacctttg acttgtttaa gtggttgcag ctgacctatg 2040

```

```

ttgcatgagg cagggactcg gtcccccttg cegtgtctcc ctccctcctc gtctgccaaag 2100
cctcgctccc taccacacca caccaggcca cccagctgc aagtgccttc cttggagcag 2160
agaggcagcc tcgtcctcct gtcccccttc ctcccagcca ccatcgttca tctgctccgg 2220
gcagaactgt gtggccccctg cctgtgccag ccatgggctc gggatggact ccctgggccc 2280
cacccattgc caagccagga aggcagcttt cgcttggttc tgctcggga cagccccggg 2340
tttccccagc atcctgatgt gtcccccttc cccacttcag aggccacca ctcagacca 2400
acgggcttgg ccttgcttgc agactataaa ct 2432

```

&lt;210&gt; 51

&lt;211&gt; 3969

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7726576CB1

&lt;400&gt; 51

```

ccgggcaggg tcgcctctag gtgctcacct ccgccacttc gccatggcgg gtccctggccc 60
ggggcgcggtg ctggagtgccc cccggcagct gctgggcccgc gtgcgcttct tggcagaggc 120
agcgcgaggc ctccgcgcgc ggccggcgcct gccagcagcg ctggcttttc tgccgcgaga 180
gggtgctctac aagctttaca aggaccagc gggaccgtcg cgcgtgcttc tgccgggtgtg 240
ggaggcagag ggccctggggc tgctgtgtgg cgccgcaggg ccagcccccg gtaccggctc 300
cggggccctc cgccgcgcgc gcgacagcat tgagctccgg cgccggcgct gcgtgcgcac 360
cacgggcgag gagctgtgca atggccacgg gctctgggtg aagctgacaa aggagcagct 420
ggcagagcac ctggggcgact gcgggctgca ggaaggctgg ctgctgggtg gccgcccggc 480
ggaggggcgga gcccgcttg taccatcga cactcccaac cactccagc ggcagcagca 540
gctctttggc gtggattatc ggccgggtgct cagggtggaa cagggtggtg acctgacata 600
ctcacatcgc ctgggatcga gacctcagcc ggcagaggca tacgcagaag ctgtacaaag 660
gctactctat gtacccccga catggacctc cgagtgcgac gaggacctga tccacttctt 720
gtatgaccac ctgggcaagg aggatgagaa cctgggtagc gtgaagcagt atgtggagag 780
catagacgtt tcctcctaca cggaggagtt caacgtgtcc tgctgacag acagcaatgc 840
cgatacctac tgggagagcg atgggtccca gtgccaacac tgggtacggc ttactatgaa 900
gaagggcacc attgtcaaga agctgtact cacagtggat accacagatg acaactttat 960
gccaaagcgg gtggtggtct atgggggtga aggggacaac ctgaagaagc tgagtgcagt 1020
gagcattgac gagacctca tcggggatgt ctgtgtcctg gaggacatga ccgtccacct 1080
cccgatcatc gagatccgca tcgtggagtg ccgagatgat gggattgatg ttctgtctccg 1140
aggggtcaag atcaagtcac ctagacagcg ggaactaggg ttgaatgcag acctgttcca 1200
gccaaactagt ctggtgcat atccacgcct agaaggcacc gacctgaag tactgtaccg 1260
cagagctgtc ctctgcaga gactcatcaa gatcctcgat agtgtcctgc accacctggt 1320
acctgcctgg gaccacacac tgggcacctt cagtgcagat aagcaagtga agcagttcct 1380
actgctgtcc cgccagcggc caggcctggt ggctcagtc ctgcgtgact ctgagagcag 1440
caagcccagc ttcattgccac gcctatacat caaccgccgt cttgccatgg aacaccgtgc 1500
ctgcccctct cgagacctg cctgcaagaa tgcagtcttc acccaggtat atgaaggcct 1560
caagccctct gacaaatatg aaaagccct ggactacagg tggcccatgc gctatgacca 1620
gtggtgggag tgtaaattta ttgcagaagg catcattgac caagggggtg gtttccggga 1680
cagcctggca gatatgtcag aagagctgtg ccctagctca gcggataccc ccgtgccctt 1740
gcccttcttt gtacgcacag ccaaccaggg caatggcact ggtgaggctc gggacatgta 1800
tgtacccaac ccctcctgcc gagactttgc caagtatgaa tggatcggac agctgatggg 1860
ggctgccctt cggggtgaagg agttcctggt cctggccctg cctggttttg tgtggaagca 1920
gctttctggt gagggagtga gctggagcaa ggacttccca gctgtggact ctgtgctggt 1980
gaagctcctg gaagtgtgg aaggaatgga caaggagacg tttgagttca agtttgggaa 2040
ggaactaaca ttcaccactg tactgagtga ccaacagggt gtggagctga tccctggggg 2100
tgcaggcatc gtctgtggat atggggaccg ttctcgtttc atccaactgg tccagaaggc 2160
acggctagag gagagcaagg agcagggtggc agctatgcag gcaggctctc tgaagggtgt 2220
accacaggct gtgctggact tgctgacctg gcaagagttg gagaagaaag tgtgtgggga 2280

```



```

tccagaggtc actgtggatg ctctgcgcaa gctcaccogg tttgaggact tcgagccatc 2340
tgactcgcgg gtgcagtatt tctgggaggg actgaacaac ttcaccaacg aggaccggag 2400
ccgcgtcctg cgctttgtca cgggccgcag tcgcctgcca gcacggatct acatctaccc 2460
agacaagctg ggctacgaga ccacagacgc gctgcccag tcttcacttt gctccagcac 2520
cctcttcctg ccacactatg ccagtgcgaa ggtatgagag gagaagctcc gctatgcggc 2580
ctacaactgc gtggccatcg acactgacat gagcccttgg gaggagttag gcgtgccggc 2640
ggctgtggga ccagcaagac tgcacgtgtc cctcttgccc ttgcccaggg cgaagacacc 2700
ttccctgccc tgggttggtg gacgtgctca gcaaaaaccc atgtgccctg ctccctgtgtg 2760
cagttggggg aggggcagct ggcatgggtc ggtaacacta gtggcccagc ccgcgagacc 2820
cacaagccct acccgtgctg gggcttgctt cccgaggtat ttcacctctt aagagggaat 2880
cttccacaag cccagcaca gctgccaggc ctgagctact tgaagggggc catctaggtc 2940
cccaacccat ggactttgcc tccattttca gctccgcctt ttttctctta ttttctctct 3000
ggctttcttc agccatgact cacaactaaa aacataaaac actggagggtt agtggaggcc 3060
cctcccaag caggagacct gggatgggca gggagtata gccaaactcc ttggtcacct 3120
gctccaagaa ggaagcagta gctgagcacc tgccctcaca tactgctctt tccctctctc 3180
cctccacacc agagatgtgg tgagctctgt tcttctacca acccagtctc aacacacaaa 3240
gtgccaccac cttccctgac tcagaaccca catccactca atgtgaactc tactaccacg 3300
acctcccat attcctcact tctccatcac ctccagcctg actccctgtc tgccctttca 3360
cccccaagat tttgcacagg ttaaggccag ttatggcctt tttgaaatct gtaatagctc 3420
ccctttcccc aactctaaag cctagacctt aaacctgttc ctagaactct ggccccacc 3480
attcctcagt gccacctttc tgctgtgaa aggccacagt gatgcccccc agtgtgaggc 3540
gggaggtgtg cctcttccc cagccaagcc tttttacca ctcccagggt ggcagctatg 3600
caggcaggtc tgctgaagggt ggtaccacag gctgtgctgg acttgctgac ctggcaagag 3660
ttggagaaga aagtgtgtgg ggatccagag gtcactgtgg atgctctgag caagctcacc 3720
cgggttgagg acttcgagcc atctgactcg cgggtgcagt atttctggga ggcactgaac 3780
aacttcacca acgaggaccg gagccgcttc ctgcgctttg tcacgggccc cagtgcctg 3840
ccagcacgga tctacatcta cccagacaag ctgggctacg agaccacaga cgcgctgccc 3900
gagtcttcca cttgctccag caccctctct tgccagcaca ctgcgccgta taagttagcg 3960
agctcgtec 3969

```

&lt;210&gt; 52

&lt;211&gt; 2537

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503507CB1

&lt;400&gt; 52

```

gggttcgggg gcgcccgcgt gtgaggccgg ggcctagagc cagccgcggc cgcgcaggag 60
gggcccaggg cccgcgctcg cccgcgtccc cgccttcctc cgcgctcag cccgcctcg 120
gctcgctgcc cttggctctc gtgcgcatgg cctccgtcgc ccaggagagc gcgggctcgc 180
agcgcgggct accgcgcgct cagggggcgc tgccggggct gctactgtc tgctgtggc 240
tgccaagcgg ccgtgcggcc ttgccgcccg cggcgccgct gtccgaactg cagcgcagc 300
tgtcgggctg ggagcagctg ctggaggagt tccgcggca actgcagcag gagcggcctc 360
aggaggagct ggagctggag ctgcgcgcgg cgggcggccc ccaggaggac tgccggggcc 420
ggggcagcgg cggctacagc gcaatgcctg acgccatcat ccgcaccaag gactccctgg 480
cggcctgtgc cagcttcctg cgggcgcggc cggcgtgctg gggctggcgg caatgcgtgg 540
cgccctgctg ctccgagccg cgtgctccg tggcgtggt ggagctgcc cggcgccccg 600
cgccccggc agcgtgtctc ggctgtctacc tcttcaactg cagggcgcg gcgcgcaacg 660
tctgcaagtt cgcgtgcac agcggctaca gcagctacag cctcagccgc gcgcgggacg 720
gcgcggccct ggccaccgcg cgcgcctcgc cccggcagga aaaggatgcg cctccactta 780
gcaaggctgg gcaggatgtg gttctgcatc tgcccacaga cggggtgggt ctagacggcc 840
gcgagagcac agatgaccac gccatcgtec agtatgagt ggcactgctg cagggggacc 900
cgtcagtgga catgaagggt cctcaatcag gaacctgaa gctgtcccac ctacaggagg 960

```

```

gaacctacac cttccagctg accgtgacgg aactgcccgg gcagagaagc tctgacaacg 1020
tgtcagtgac agtgcttcgc gcagcctact ccacaggagg atgtttgcac acttgctcac 1080
gctaccactt cttctgtgac gatggctgct gcattgacat cacgctcgcc tgcgatggag 1140
tgcagcagtg tctgatggg tctgatgaag acttctgcca gaatctgggc ctggaccgca 1200
agatggtaac ccacacggca gctagtctcg cctgccaag aaccacaggg ccgagtgaag 1260
atgcaggggg tgactccttg gtggaaaagt ctgagaaagc cactgcccc aacaagccac 1320
ctgcattatc aaacacagag aagaggaaaag ttatatattt gagtcaaagg gtgatggagg 1380
aggaggggaa caccagccc cagaaacagg tgcagtgcta cccctggcgc tgggtttggc 1440
tatcactgct ctgctgcttc tcatggttgc atgcccacta cgactggtga aacagaaact 1500
gaaaaaagct cgtcccatca catctgagga atcggactac ctcataaatg ggatgtatct 1560
atagtaatgt aatttcaata ccttggggca gggacatgtt ttgtttataa tttatacatc 1620
tattaagttc tggatattta cagcttcttt tgtttttaat tgggccagaa gattctgcaa 1680
atcccaaatc tttctttatt atttattgta aaaaaagttt ccttagaagt cataaaatat 1740
tttgaaattt agagaggaat tcatgattaa agattcctaa aaatataatt ctgatttatg 1800
taagctgtcc ctgaaaatag aaatgtgtac ttagctgaga gaaaattcag catctcagga 1860
ggtggtatta ggatgactgt gttaacccat taccttttag aagccaactg ttggccctt 1920
accatgctgg actgctatag gccagcttc ccttgttct gtggccctt tcttctct 1980
tgaagctccc agtattcttt ttcttttccc ctctaaacct gtttctgaga gtggatctca 2040
agcaagtcca tgccttcaat cagatgttac ttaggtggg tatacctaaa ttataaacct 2100
tatgtacaag tcagttaagg tttagggaagg tgagtgtggg tcttctctaa tccctctgac 2160
gtcatgtcat ataggtggct gcctccttag actgacctt gggagaaaaa aacccagac 2220
tttgaattag taacagctct aagatggtca tgcagtgaga taggaaatca agatggaagc 2280
agagaatctg gcatgcaaaa aactaacaga aacttagttg aaggcaaaga gagcaaggag 2340
aaagtttaat acttcattac atcaaatcaa cactgctcca tggtagagc acagcaactc 2400
atztatatat atatatatag gctttgttga tgaaaaacaa caattgaaga gaggacgttg 2460
agtggattcc tgggtacagc ttttgtaaaa atgtcaccat ggctttcatc caatggaatg 2520
agtcgatgtt ttttaat 2537

```

&lt;210&gt; 53

&lt;211&gt; 2526

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503506CB1

&lt;400&gt; 53

```

gggttcgggg gcgcgcgct gtgaggccgg ggcctagagc cagccgcggc cgcgcaggag 60
ggggccaggg ccgcgcctcg ccgcgtccc cgccttctc ccgcgctcag cccgcctcg 120
gctcgctgcc cttggctctc gtgcctatgg cctccgtcgc ccaggagagc gcgggctcgc 180
agcgccggct accgcgcgt cagggggcgc tgcgcgggct gctactgctc tgcctgtggc 240
tgccaagcgg ccgtgcggcc ttgcccgcgc cggcgccgct gtccgaactg cagcgcgacg 300
tgtcggggct ggagcagctg ctggaggagt tccgcggca actgcagcag gagcggcctc 360
aggaggagct ggagctggag ctgcgcgcgg gcggcgccc ccaggaggac tgcccgggccc 420
ggggcagcgg cggctacagc gcaatgcctg acgccatcat ccgcaccaag gactccctgg 480
cggcggtgct cagcttccctg cgggcgcggc cggcgtgctg gggctggcgg caatgcgtgg 540
cggcctgctg ctccgagccg cgtgctcgc ttggcgtggt ggagctgccc cggcgccccg 600
cgccccgggc agcgtgctc ggctgctacc tcttcaactg cagggcgcg gcgcgcaacg 660
tctgcaagtt cgcgctgcac agcggctaca gcagctacag cctcagccgc gcgcggagc 720
gcgcgcctt ggccaccgcg cgcgcctcgc cccggcagga aaaggatgct cctccactta 780
gcaaggctgg gcaggatgtg gttctgcatc tgcccacaga cggggtggtt ctgacggcc 840
gcgagagcac agatgaccac gccatcgctc agtatgagtg ggcaactgct cagggggacc 900
cgtcagtgga catgaaggct cctcaatcag gaaccctgaa gctgtcccac ctacaggagg 960
gaacctacac cttccagctg accgtgacgg aactgcccgg gcagagaagc tctgacaacg 1020
tgtcagtgac agtgcttcgc gcagcctact ccacaggagg atgtttgcac acttgctcac 1080

```

```

gctaccactt cttctgtgac gatggctgct gcattgacat cacgctcgcc tgcgatggag 1140
tgcagcagtg tctgatggg tctgatgaag acttctgcca gaatctgggc ctggaccgca 1200
agatggtaac ccacacggca gctagtcctg ccctgccaag aaccacaggg ccgagtgaag 1260
atgcaggggg tgactccttg gtggaaaagt ctcagaaaag cactgcccc aacaagccac 1320
ctgcattatc aaacacagag aagaggaatc attccgcctt ttggggacca gagagtcaaa 1380
tcattcctgt gatgccaggt gcagtgtac ccctggcgct gggtttggct atcactgctc 1440
tgctgcttct catggttgca tgccgactac gactggtgaa acagaaactg aaaaaagctc 1500
gtcccattac atctgaggaa tcggactacc tcataaatgg gatgtatcta tagtaatgta 1560
atttcaatac cttggggcag ggacatgttt tgtttataat ttatacatct attaagttct 1620
ggatattttac agcttctttt gtttttaatt gggccagaag attctgcaaa tcccaaattct 1680
ttctttatta ttattgttaa aaaaagtttc cttagaagtc ataaaatatt ttgaaattta 1740
gagaggaatt catgattaaa gattcctaaa aatataattc tgatttatgt aagctgtccc 1800
tgaaaataga aatgtgtact tagctgagag aaaattcagc atctcaggag gtggtattag 1860
gatgactgtg ttaaccatt accttttaga agccaactgt tggcccctta ccatgctgga 1920
ctgctatagg ccagcttcc ccttggtctg tggccctttt cttcctcctt gaagctccca 1980
gtattccttt tcttttcccc tctaaacctg tttctgagag tggatctcaa gcaagttcat 2040
gccttaacat agatgttact taggggtgggt atacctaaat tataaacctt atgtacaagt 2100
cagtaagcct tagggaaagt gagtgtgggt ccttcctaata cctctgacg tcatgtcata 2160
taggtggctg cctccttaga ctgaccttgg ggagaaaaaa accccagact ttgaattagt 2220
aacagctcta agatgggtcat gcagtgtgat aggaaatcaa gatggaagca gagaatctgg 2280
catgccaaaa actaacagaa acttagttga aggcaaagag agcaaggaga aagtttaata 2340
cttcattaca tcaaatcaac actgctccat ggtgagagca cagcaactca tttatatata 2400
tatatatagg ctttgttgat gaaaaacaac aattgaagag aggacgttga gtggattcct 2460
gggtacagct tttgtaaaaa tgtcaccatg gctttcatcc aatggaatga gtcgatgttt 2520
tttaat 2526

```

&lt;210&gt; 54

&lt;211&gt; 2464

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503509CB1

&lt;400&gt; 54

```

gggttcgggg gcgcgcgcgt gtgaggccgg ggcctagagc cagcccgggc gcgcaggag 60
ggggccaggg ccgcgcgtcg ccgcgcgtccc gccttctc cgcgcgtcag ccccgctcg 120
gtcgcgtgcc cttggctctc gtgcgcattg cctccgtcgc ccaggagagc gcgggctcgc 180
agcgcgcgt accgcgcgt cacggggcgc tgcgcgggct gctactgctc tgcctgtggc 240
tgccaagcgg ccgtgcggcc ttgcgcgccg cgcgcgcgt gtccgaactg cacgcgcagc 300
tgtcgggctg ggagcagctg ctggaggagt tccgcggca actgcagcag gagcggcctc 360
aggaggagct ggagctggag ctgcgcgcgg gcgcgcggcc ccaggaggac tgcccgggcc 420
cgggcagcgg cggctacagc gcaatgcctg acgccatcat ccgcaccaag gactccctgg 480
cggcggtgct cagcttctc cgggcgcggc cggcgtgctg gggctggcgg caatgcgtgg 540
cggcctgctg ctccgagcgg cgctgctcgg tggcgtgggt ggagctgccc cggcgccccg 600
cgccccggc agcctgtctc ggctgtacc tcttcaactg cacggcgccg gcccgcaacg 660
tctgcaagtt cgcgtgac agcggttaca gcagctacag cctcagccgc gcccggaacg 720
gcgcgcctt ggccaccgcg cgcgcctcgc cccggcaggg tgctcaatc aggaacctg 780
aagctgtccc acctacagga gggaacctac acctccagc tgaccgtgac ggacactgcc 840
gggcagagaa gctctgacaa cgtgtcagtg acagtgttc gcgcagccta ctccacagga 900
ggatgtttgc acacttgctc acgctaaccac ttcttctgtg acgatggctg ctgcattgac 960
atcacgctcg cctgcgatgg agtgcagcag tgtcctgatg ggtctgatga agacttctgc 1020
cagaatctgg gcctggaccg caagatggta acccacagg cagctagtcc tgccctgcca 1080
agaaccacag ggccgagtga agatgcaggg ggtgactcct tgggtggaaa gtctcagaaa 1140
gccactgccc caaacaagcc acctgcatta tcaaacacag agaagaggaa tcattccgcc 1200

```

```

ttttggggac cagagagtc aatcattcct gtgatgccag atagtagttc ctcaggggaag 1260
aacagaaaaag aggaaagtta tatatttgag tcaaagggtg atggaggagg aggggaacac 1320
ccagccccag aaacagggtgc agtgetaccc ctggcgctgg gtttggctat cactgctctg 1380
ctgcttctca tgggttgcag ccgactacga ctgggtgaaac agaaactgaa aaaagctcgt 1440
cccattacat ctgaggaatc ggactacctc ataaatggga tgtatctata gtaatgtaat 1500
ttcaatacct tggggcaggg acatgttttg ttataaattt atacatctat taagttctgg 1560
atatttacag cttcttttgg ttttaattgg gccagaagat tctgcaaate ccaaactctt 1620
ctttattatt tattgtaaaa aaagtttcct tagaagtcac aaaatatttt gaaatttaga 1680
gaggaattca tgattaaaga ttctaaaaa tataattctg atttatgtaa gctgtccctg 1740
aaaatagaaa tgtgtactta gctgagagaa aattcagcat ctcaggagggt ggtatttaga 1800
tgactgtgtt aacccattac ctttttagaag ccaactgttg gcccttacc atgctggact 1860
gctataggcc cagcttcccc ttgttctgtg gcccttttct tctccttga agctcccagt 1920
attcttttct ttttccccctc taaacctgtt tctgagagtg gatctcaagc aagttcatgc 1980
cttcaatcag atgttactta ggggtgggtat acctaaatta taaaccttat gtacaagtca 2040
gtaagcctta gggaagggtga gtgtgggtcc ttctaatcc ctctgacgtc atgtcatata 2100
gggtggctgc tccttagact gaccttttggg agaaaaaac cccagacttt gaattagtaa 2160
cagctctaag atgggtcatgc agtgagatag gaaatcaaga tgaagcaga gaatctggca 2220
tgccaaaaac taacagaaac ttagttaga gcaaagagag caaggagaaa gtttaatact 2280
tcattacatc aaatcaacac tgcctcatgg tgagagcaca gcaactcatt tatatatata 2340
tatataggct ttgttgatga aaaacaacaa ttgaagagag gacgttagt ggattcctgg 2400
gtacagcttt tgtaaaaatg tcaccatggc tttcatcaa tggaatgagt cgatgttttt 2460
taat

```

&lt;210&gt; 55

&lt;211&gt; 1452

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7505800CB1

&lt;400&gt; 55

```

ccgaggcgag atggcgggcca ccgaggggggt cggggagggt gcgcaagggg gcgagcccg 60
gcagccggcg caacccccgc ccagccgcga cccaccgccc cccagcagc agcacaagg 120
agagatggcg gccgaggctg gggaagccgt ggcgtcccc atggacgacg ggtttgtgag 180
cctggactcg ccctoctatg tcctgtacag gcatttccgg agagttcttt tgaagtcact 240
tcagaaggat ctacatgagg aaatgaacta catcactgca ataattgagg agcagcccaa 300
aaactatcaa gtttggcatc ataggcgagt attagtggaa tggctaagag atccatctca 360
ggagcttgaa tttattgctg atattcttaa tcaggatgca aagaattatc atgcctggca 420
gcatcgacaa tgggttattc aggaatttaa actttgggat aatgagctgc agtatgtgga 480
ccaacttctg aaagaggatg tgagaaataa ctctgtctgg aaccaaagat acttcgttat 540
ttctaaccac actggctaca atgatcgtgc tgtattggag agagaagtcc aatacactct 600
ggaaatgatt aaactagtac cacataatga aagtgcagtg aactatttga aagggatttt 660
gcaggatcgt ggtctttcca aatatectaa tctgttaaat caattacttg atttacaacc 720
aagtcatagt tccccctacc taattgcctt tottgtggat atctatgaag acatgctaga 780
aaatcagtg gacaataagg aagacattct taataaagca ttagagttat gtgaaatcct 840
agctaaagaa aaggacacta taagaaagga atattggaga tacattggaa gatcccttca 900
aagcaaacac agcacagaaa atgactcacc aacaaatgta cagcaataac accatccaga 960
agaacttgat ggaatgcttt tattttttat taagggacc tgcaggagt tccacagaga 1020
gtggtccttc cctttgcctg tgggtgtaaaa gtgcatcaca caggatttgc tttttaacaa 1080
gaactgatgc tccttgggtg ctgctgctac tcagactagc tctaagtaat gtgattcttc 1140
taaagcaaag tcattggatg ggaggaggaa gaaaaagtcc cataaaggaa cttttgtagt 1200
cttatcaaca tataatctaa tcccttagca tcagctcctc cctcagtggt acatgcgtca 1260
agattttag cagtaataac tgcaggtcac ttgtatgtaa tggatgtgag gtagccgaag 1320
tttggttcag taagcaggga atacagtcgt tccatcagag ctggtctgca cactcacatt 1380

```

atcttgctat cactgtaacc aactaatgcc aaaagaacgg ttttgtaata .aaattatagc 1440  
tgtatctaaa aa 1452

<210> 56  
<211> 1802  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 7503141CB1

<400> 56  
cgccggtgcc gggcgaacat ggccggcggcc accggaccct cgttttggct ggggaatgaa 60  
acctgaagg tgccgctggc gctctttgcc ttgaaccggc agcgccctgtg tgagcggctg 120  
cggaagaacc ctgctgtgca ggccggctcc atcgtggtcc tgcagggcgg ggaggagact 180  
cagcgctact gcaccgacac cggggtcctc ttccgccagg agtccttctt tcaactggcg 240  
ttcgggtgca ctgagccagg ctgctatggt gtcacgatg ttgacactgg gaagtcgacc 300  
ctgtttgtgc ccaggcttcc tgccagccat gccacctgga tgggaaagat ccattccaag 360  
gagcacttca aggagaagta tgccgtggac gacgtccagt acgtagatga gattgccagc 420  
gtcctgacgt cacagaagcc ctctgtcctc ctcaacttgc gtggcgtaa cacggacagc 480  
ggcagtgctc gcagggaggc ctcccttgac ggcatcagca agttcgaagt caacaatacc 540  
attcttcacc cagagatcgt tgagtgcctc ttccagcact actgctactc ccggggcggc 600  
atgcgccaca gctcctacac ctgcatctgc ggcatggtg agaactcagc cgtgctacac 660  
tacggacacg ccggagctcc caacgaccga acgatccaga atggggatat gtgcctgttc 720  
gacatggcgg gtgagtatta ctgcttcgct tccgacatca cctgctcctt tcccgccaac 780  
ggcaagttca ctgcagacca gaaggccgtc tatgaggcag tgctgcggag ctcccgtgcc 840  
gtcatgggtg ccatgaagcc aggtgtctgg tggcctgaca tgcaccgcct ggctgaccgc 900  
atccacctgg aggagctggc ccacatgggc atcctgagcg gcagcgtgga cgccatggtc 960  
caggctcacc tggggggcgt gtttatgcct caggggcttg gccacttctt gggcattgac 1020  
gtgcacgacg tgggaggcta ccagaggggc gtggagcgca tcgacgagcc cggcctgcgg 1080  
agcctgcgca ctgcacggca cctgcagcca ggcatggtgc tcaccgtgga gccgggcata 1140  
tacttcatcg accacctcct ggatgaggcc ctggcggacc cggcccgccg ctccctcctt 1200  
aaccgcgagg tctgcagcg ctttcgcggt tttggcgggg tccgcatcga ggaggacgtc 1260  
gtggtgactg acagcggcat agagctgctg acctgcgtgc cccgactgt ggaagagatt 1320  
gaagcatgca tggcaggctg tgacaaggcc tttaccccct tctctggccc caagtagagc 1380  
cagccagaaa tcccagcgca cctggggggc tggccttgca acctcttttc gtgatgggca 1440  
gcctgctggt cagcactcca gtagcgagag acggcaccga gaatcagatc ccagcttcgg 1500  
catttgatca gaccaaacag tgctgtttcc cggggaggaa acactttttt aattaattac 1560  
ccttttgtag gctcccacct ttaatctggt ttataccttg cttattaaat gagcgactta 1620  
aaatgattga aaataatgct gttcttttagt agcaactaaa atgtgtcttg ctgtcattta 1680  
tattcctttt ccaggaag aagcatttct gatactttct gtcaaaaatc aatatgcaga 1740  
atggcatttg caataaaagg tttcctaaaa tgaaaaaaa aaaaaaaaaa aaaaaaaaaa 1800  
aa 1802

<210> 57  
<211> 1833  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 7500362CB1

<400> 57  
cgggcgaaca tggcggcggc caccggaccc tcgttttggc tggggaatga aacctgaag 60

```

gtgccgctgg cgctctttgc cttgaaccgg cagcgcctgt gtgagcggct gcggaagaac 120
cctgctgtgc aggcgcggctc catcgtgtcc ttctttcact gggcggtcgg tgtcactgag 180
ccaggctgct atgggtgtcat cgatgttgac actgggaagt cgaccctgtt tgtgccagg 240
cttcctgccca gccatgccac ctggatggga aagatccatt ccaaggagca cttcaaggag 300
aagtatgccg tggacgacgt ccagtacgta gatgagattg ccagcgtcct gacgtcacag 360
aagccctctg tcctcctcac tttgctgggc gtcaacacgg acagcggcag tgtctgcagg 420
gaggcctcct ttgacggcat cagcaagttc gaagtcaaca ataccattct tcaccagag 480
atcgttgagt gccagtggtt taagacggat atggagctgg aggttctgct ctataccaat 540
aaaatctcca gcgaggccca ccgtgaggta atgaaggctg taaaagtggg aatgaaagaa 600
tatgagttgg aaagcctctt cgagcactac tgctactccc ggggcggcat gcgccacagc 660
tcctacacct gcatctgcgg cagtggtag aactcagccg tgctacacta cggacacgcc 720
ggagctccca acgaccgaac gatccagaat ggggatatgt gcctgttcga catgggcggg 780
gagtattact gcttcgcttc cgacatcacc tgctcctttc ccgccaacgg caagttcact 840
gcagaccaga aggcggtcta tgaggcagtg ctgaggagct ccggtgccgt catgggtgcc 900
atgaagccag gtgtctgggt gcctgacatg caccgcctgg ctgaccgcat ccacctggag 960
gagctggccc acatgggcat cctgagcggc agcgtggacg ccatgggtcca ggctcacctg 1020
ggggcctgtt ttatgcctca cgggcttggc cacttcctgg gcattgacgt gcacgacgtg 1080
ggaggctacc cagagggcgt ggagcgcac gacgagcccg gcctgcggag cctgcgcact 1140
gcacggcacc tgcagccagg catggtgtct accgtggagc cgggcatcta cttcatcgac 1200
cacctcctgg atgaggccct ggcggaaccg gccgcgcct ccttccttaa ccgcgaggtc 1260
ctgcagcgtt ttgcgggttt tggcgggttc cgcatcgagg aggacgtcgt ggtgactgac 1320
agcggcatag agctgctgac ctgctgtccc cgactgtgg aagagattga agcatgcatg 1380
gcaggctgtg acaaggcctt taccctcttc tctggcccca agtagagcca gccagaaatc 1440
ccagcgcacc tgggggcctg gccttgcaac ctcttttctg gatgggcagc ctgctggtca 1500
gcactccagt agcgagagac ggcaccagca atcagatccc agcttcggca tttgatcaga 1560
ccaaacagtg ctgtttcccg gggaggaaac acttttttaa ttaccctttt gcaggctccc 1620
acctttaatc tgttttatac cttgcttatt aaatgagcga cttaaaatga ttgaaaataa 1680
tgctgttctt tagtagcaac taaaatgtgt cttgctgtca tttatattcc ttttccagg 1740
aaagaagcat ttctgatact ttctgtcaaa aatcaatatg cagaatggca tttgcaataa 1800
aaggtttcct aaaatggtca aaaaaaaaaa aaa 1833

```

&lt;210&gt; 58

&lt;211&gt; 2465

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7503328CB1

&lt;400&gt; 58

```

ggcccttccg cgggtgatca gctggtctgc gctccctga cgtgggctgg ggcacgtcac 60
cgccgaatgg cagcctccag aaagccaccg cgagtaaggg tgaatcacca ggattttcaa 120
ctgagaaatt taagaataat tgaacctaac gaggtgacac actcaggaga cacagggtgtg 180
gaaacagacg gcagaatgcc tccaaagggtg acttcagagc tgcttcggca gctgagacaa 240
gccatgagga actctgagta tgtgaccgaa ccgatccagg cctacatcat cccatcgga 300
gatgctcatc agagttagta tattgctcca tgtgactgtc ggcgggcttt tgtctctgga 360
ttcgtggtct ctgcgggcac agccatcatc acagaagagc atgcagccat gtggactgac 420
gggcgctact ttctccaggc tgccaagcaa atggacagca actggacact tatgaagatg 480
ggtctgaagg acacaccaac tcaggaagac tggctggtga gtgtgcttcc tgaaggatcc 540
agggttggtg tggaccctt gatcattcct acagattatt ggaagaaaat ggccaaagtt 600
ctgagaagtg ccggccatca cctcattcct gtcaaggaga acctcgttga caaaatctgg 660
acagagcgtc ctgagcggcc ttgcaagcct ctctcacac tgggcctgga ttacacaggc 720
atctcctgga aggacaaggt tgcagacttt cggttgaaaa tggctgagag gaacgtcatg 780
tggtttgggt tcaactgcctt ggatgagatt gcgtggctat ttaatctccg aggatcggat 840
gtggagcaca atccagtatt tttctcctac gcaatcatag gactagagac gatcatgctc 900

```

```

ttcattgatg gtgaccgcat agacgcccc agtgtgaagg agcacctgct tcttgacttg 960
ggtctggaag ccgagtacag gatccagggt catccctaca agtccatcct gagcgagctc 1020
aaggccctgt gtgctgacct ctccccaagg gagaagggtgt gggtcagtga caaggccagc 1080
tatgctgtga gcgagacct ccccaaggac caccgctgct gtatgcctta ccccccatc 1140
tgcacgcca aagctgtgaa gaattcagct gagtcagaag gcatgaggcg ggctcacatt 1200
aaagatgctg ttgctctctg tgaactcttt aactggctgg agaaagaggt tcccaaaggt 1260
ggtgtgacag agatctcagc tgctgacaaa gctgaggagt ttcgcaggca acaggcagac 1320
tttgtggacc tgagcttccc aacaatttcc agccagtccc tgagacgaat aggaccttgt 1380
ccctggatga ggtgtacctt attgactcgg gtgctcaata caaggatggc accacagatg 1440
tgacgaggac aatgcatttt gggaccctta cagcctacga gaaggaaatgc ttcacatatg 1500
tcctcaaggg ccacatagct gtgagtgcag ccgttttccc gactggaacc aaaggtcacc 1560
ttcttgactc ctttgcccggt tcagctttat gggattcagg cctagattac ttgcacggga 1620
ctggacatgg tgttgggtct tttttgaatg tccatgaggg tccttgccgc atcagttaca 1680
aaacattctc tgatgagccc ttggaggcag gcatgattgt cactgatgag cccgggtact 1740
atgaagatgg ggcttttggg attcgcattg agaatgttgt ccttgtggtt cctgtgaaga 1800
ccaagtataa ttttaataac cggggaagcc tgaccttga accttaaca ttggttccaa 1860
ttcagaccaa aatgatagat gtggattctc ttacagacaa agagtgcgac tggctcaaca 1920
attaccacct gacctgcagg gatgtgattg ggaaggaatt gcagaaaacag ggccgccagg 1980
aagctctcga gtggctcatc agagagacgc aactcatctc caaacagcat taataaatac 2040
ctccccggtt ttgtttttgt aaaatgctct ggaggaaagga agaaacgtgg cagatccctg 2100
acatctttcc ctttctcttt ccttcttccc cactccctct ttttacttta gactttaaga 2160
agaacagaaa atcttcttat cctctttgat attttattgc aaacactcag tcttttatga 2220
tttttaattt gttgagaaca agccaagaat aaaattgctg caccagaagg aggggtccctc 2280
caaagttgaa cacttggtga aaggaagatg ccccgacttc tttggccagt gatggggaat 2340
cagtgaagtgc tccatgatgg tcatgttcca ggtgctagta catcattcat gatcacctta 2400
atgctcatga gactatattt atgatcagtg aataaaaatg tcagaactgt gaaaaaaaaa 2460
aaaaa 2465

```

<210> 59

<211> 2560

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 7510464CB1

<400> 59

```

ggcccttccg cgggtgatca gctggtctgc gctccctga cgtgggctgg ggcaagtcac 60
cgccgaatgg cagcctccag aaagccaccg cgagtaaggg tgaatcacca ggattttcaa 120
ctgagaaaatt taagaataat tgaacctaac gaggtgacac actcaggaga cacagggtgtg 180
gaaacagacg gcagaatgcc tccaaagggtg acttcagagc tgcttcggca gctgagacaa 240
gcatgagga actctgagta tgtgaccgaa ccgatccagg cctacatcat cccatcggga 300
gatgctcatc agagtgaagta tattgctoca tgtgactgtc ggcgggcttt tgtctctgga 360
ttcgatggct ctgcggggcag agccatcatc acagaagagc atgcagccat gtggactgac 420
ggcgctact ttctccaggc tgccaagcaa atggacagca actggacact tatgaagatg 480
ggtctgaagg acacaccaac tcaggaagac tggctggtga gtgtgcttc tgaaggatcc 540
agggttggtg tggaccctt gatcatctct acagattatt ggaagaaaat ggccaaagtt 600
ctgagaagtg ccggccatca cctcattcct gtcaaggaga acctcgttga caaaatctgg 660
acagaccgtc ctgagcgccc ttgcaagcct ctctcacac tgggcctgga ttacacaggc 720
atctcctgga aggacaaggt tgcagacctt cggttgaaaa tggctgagag gaacgtcatg 780
tggtttggg tcaactgcctt ggatgagatt gcgtggctat ttaatctccg aggatcagat 840
gtggagcaca atccagtatt tttctcctac gcaatcatag gactagagac gatcatgctc 900
ttcattgatg gtgaccgcat agacgcccc agtgtgaagg agcacctgct tcttgacttg 960
ggtctggaag ccgaatacag gatccagggt catccctaca agtccatcct gagcgagctc 1020
aaggccctgt gtgctgacct ctccccaagg gagaagggtgt gggtcagtga caaggccagc 1080

```

```

tatgtctgtga gcgagaccat ccccaaggac caccgctgct gtatgcctta cacccccata 1140
tgcattcgcca aagctgtgaa gaattcagct gagtcagaag gcatgaggcg ggctcacatt 1200
aaagatgctg ttgtctctctg tgaactcttt aactggctgg agaaagaggt tcccaaagg 1260
gggtgtgacag agatctcagc tgctgacaaa gctgaggagt ttgcaggca acaggcagac 1320
tttgtggacc tgagcttccc aacaatttcc agtacgggac ccaacggcgc catcattcac 1380
tacgcgccag tccttgagac gaataggacc ttgtccctgg atgagggtgta ccttattgac 1440
tcgggtgtct aatacaagga tggcaccaca gatgtgaogc ggacaatgca ttttgggacc 1500
cctacagcct acgagaagga atgcttcaca tatgtcctca agggccacat agctgtgagt 1560
gcagccgttt tcccgactgg aaccaaagg  cacttcttg actcctttgc ccgttcagct 1620
ttatgggatt caggcctaga ttacttgac gggactggac atgggtgttg gtcttttttg 1680
aatgtccatg agggctcctg cggcatcagt taaaaacat tctctgatga gcccttggag 1740
gcaggcatga ttgtcactga tgagccggg tactatgaag atggggcttt tggaaatcgc 1800
attgagaatg ttgtccttgt ggttcctgtg aagaccaagt ataatttta taaccgggga 1860
agcctgacct ttgaacctct aacattgggt ccaattcaga ccaaaatgat agatgtggat 1920
tctcttacag acaaaagga gctgtggaat gggattctcc cagctagaag cctcttctgc 1980
ctgttccagt tcacagtgcg actggtcaa caattaccac ctgacctgca gggatgtgat 2040
tgggaaggaa ttgcagaaac agggccgcca ggaagctctc gactggctca tcagagagac 2100
gcaacccatc tccaaacagc attaataaat acctccccg ttttgttttt gtaaaatgct 2160
ctggaggaag gaagaaacgt ggcagatccc tgacatcttt cccctttcct ttcttcttc 2220
cctacctccc ctttttactt tagactttta gaagaacaga aaatcttctt atcctctttg 2280
atattttatt gcaaacactc agtcttttat gattttttta ttgttgagaa caagccaaga 2340
ataaaattgc tgcaccagaa ggagggtccc tccaaagtgt aacacttggg gaaaggaaga 2400
tgccccgact tctttggcca gtgatgggga atcagtgagt gctccatgat ggtcatgttc 2460
caggtgctag tacatcattc atgatcacct taatgtcat gagactatat ttatgatcag 2520
tgaataaaaa tgtcagaact gtgaaaaaaa aaaaaaaaaa 2560

```

&lt;210&gt; 60

&lt;211&gt; 2254

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7510394CB1

&lt;400&gt; 60

```

gcaggcatgg cggcggtctat gccgcttgct ctgctcgctc tgttgctcct ggggcccggc 60
ggctgggtgcc ttgcagaacc cccacgcgac agcctgcggg aggaacttgt catcacccc 120
ctgccttccg gggacgtagc cgccacattc cagttccgca cgcgctggga ttcgagctt 180
cagcgggaag gactgtccca ttacaggctc ttcccaaag ccctggggca gctgatctcc 240
aagtattctc tacgggagct gcacctgtca ttacacaag gcttttggag gaacccgatac 300
tggggggccac ctttcttgca ggccccatca ggtgcagagc tgtgggtctg gttccaagac 360
actgtcactg agtttagcag ccagctgtgg actttgaaag agggagcaga ggtagcccca 420
ggacagttag tggatttgtg tctctatcca gtgtggataa atcttggaag gagctcagta 480
atgtcctctc agggatcttc tgccctctc tcaacttcat cgactccacc aacacagtca 540
ctcccactgc ctcttcaaa cccctgggtc tggccaatga cactgaccac tactttctgc 600
gctatgctgt gctgccgagg gaggtgggtc gcaccgaaaa cctcaccccc tggaagaagc 660
tcttgccctg tagttccaag gcaggcctct ctgtgctgct gaaggcagat cgcttgttcc 720
acaccagcta ccaactccag gcagtgcata tccgccctgt ttgcagaaat gcacgctgta 780
ctagcatctc ctgggagctg aggcagaccc tgtcagttgt atttgatgcc ttcatcacgg 840
ggcagggaaa gaaagactgg tccctcttcc ggatgttctc cogaaccctc acggagccct 900
gccccctggc ttcagagagc cgagtctatg tggacatcac cactacaac caggacaacg 960
agacattaga ggtgcaccca cccccgacca ctacatatca ggacgtcatc ctaggcactc 1020
ggaagacctc tgccatctat gacttgcttg acaccgccat gatcaacaac tctcgaaacc 1080
tcaacatcca gctcaagtgg aagagacccc cagagaatga ggcccccca gtgcccttcc 1140
tgcatgcccc gcggtacgtg agtggctatg ggctgcagaa gggggagctg agcacactgc 1200

```



```

tgtacaacac ccaccatac cgggccttcc cgggtgctgct gctggacacc gtaccctggt 1260
atctgcggct gtatgtgcac accctcacca tcacctcaa gggcaaggag aacaaaccaa 1320
gttacatcca ctaccagcct gccaggacc ggctgcaacc ccacctcctg gagatgctga 1380
ttcagctgcc ggccaactca gtcaccaagg tttccatcca gtttgagcgg gcgctgctga 1440
agtggaccga gtacacgcca gatcctaacc atggcttcta tgtcagccca tctgtcctca 1500
gcgcccctgt gccagcatg gtagcagcca agccagtga ctgggaagag agtcccctct 1560
tcaacagcct gttcccagtc tctgatggct ctaactactt tgtgcggctc tacacggagc 1620
cgctgctggt gaacctgccg acaccggact tcagcatgcc ctacaacgtg atctgcctca 1680
cgtgcactgt ggtggccgtg tgctacggct ccttctacaa tctcctcacc cgaaccttcc 1740
acatcgagga gccccgcaca ggtggcctgg ccaagcggct ggccaacctt atccggcgcg 1800
cccaggtgt cccccactc tgattcttgc cctttccagc agctgcagct gccgtttctc 1860
tctggggagg ggagcccaag ggctgtttct gccacttgct ctctcagag ttggcttttg 1920
aaccaaagt ccctggacca ggtcagggcc tacagctgtg ttgtccagta caggagccac 1980
gagccaaatg tggcatttga atttgaatta acttagaaat tcatttcctc acctgtagt 2040
gccacctcta tattgaggtg ctcaataagc aaaagtggtc ggtggctgct gtattggaca 2100
gcacagaaaa agatttccat caccacagaa aggtcggctg gcagcactgg ccaaggtgat 2160
ggggtgtgct acacagtgtg tgctactgtg tagtggatgg agtttactgt ttgtggaata 2220
aaaacggctg tttccgtgga aaaaaaaaaa aaaa 2254

```

&lt;210&gt; 61

&lt;211&gt; 2139

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7500745CB1

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 2126

&lt;223&gt; a, t, c, g, or other

&lt;400&gt; 61

```

gccgcaggca tggcggcggc tatgccgctt gctctgctcg tctgtttgct cctggggccc 60
ggcggctggt gccttgacaga acccccacgc gacagcctgc gggaggaact tgtcatcacc 120
ccgctgcctt ccggggagct agccgccaca ttccagttcc gcacgcgctg ggattcggag 180
cttcagcggg aaggagtgtc ccattacagg ctctttccca aagccctggg gcagctgac 240
tccaagtatt ctctacggga gctgcacctg tcattcacac aaggcttttg gaggaccga 300
tactgggggc cacccttctc gcaggcccca tcagtgtgga taaatcttgg aaggagctca 360
gtaatgtcct ctcagggatc ttctgcgctt ctctcaactt catcgactcc accaacacag 420
tactccacc tgccctcttc aaacccttgg gtctggccaa tgacactgac cactacttct 480
tgcgctatgc tgtgctgccg cgggaggtgg tctgcaccga aaacctcacc ccctggaaga 540
agctcttgcc ctgtagttcc aaggcaggcc tctctgtgct gctgaaggca gatcgcttgt 600
tccacaccag ctaccactcc caggcagtgc atatccgccc tgtttgcaga aatgcacgct 660
gtactagcat ctctggggag ctgaggcaga cctgtcagt tgtatttgat gccttcatca 720
cggggcaggg aaagaaagac tggctcctct tccggatgtt ctcccgaacc ctcacggagc 780
cctgccccct ggcttcagag agccgagtct atgtggacat caccacctac aaccaggaca 840
acgagacatt agaggtgcac ccaccctga ccactacata tcaggacgtc atcctaggca 900
ctcgaagac ctatgccatc tatgacttgc ttgacaccgc catgatcaac aactctcgaa 960
acctcaacat ccagctcaag tggaagagac ccccagagaa tgaggccccc ccagtgcctt 1020
tctgcatgc ccagcggtag gtgagtggct atgggctgca gaagggggag ctgagcacac 1080
tgctgtacaa caccaccca taccgggctt tcccgggtgt gctgctggac accgtaccct 1140
ggtatctgcy gctgtatgtg cacacctca ccatcacctc caagggcaag gagaacaaac 1200
caagttacat ccactaccag cctgccagg accggctgca accccacctc ctggagatgc 1260
tgattcagct gccggccaac tcagtcacca aggtttccat ccagtttgag cgggcgctgc 1320

```

```

tgaagtggac cgagtacacg ccagatccta accatggcct ctatgtcagc ccatctgtcc 1380
tcagcgccct tgtgccagc atggtagcag ccaagccagt ggactgggaa gagagtcccc 1440
tcttcaacag cctgttccca gtctctgatg gctctaacta ctttgtgcgg ctctacacgg 1500
agccgctgct ggtgaacctg ccgacaccgg acttcagcat gccctacaac gtgatctgcc 1560
tcacgtgcac tgtggtggcc gtgtgctacg gctccttcta caatctctc acccgaacct 1620
tccacatcga ggagccccgc acaggtggcc tggccaagcg gctggccaac cttatccggc 1680
gcgcccagg tgtcccccca ctctgattct tgccctttcc agcagctgca gctgccgttt 1740
ctctctgggg aggggagccc aagggtgtt tctgccactt gctctctca gaggttggctt 1800
ttgaacccaa gtgccctgga ccaggtcagg gcctacagct gtgttgcca gtacaggagc 1860
cacgagccaa atgtggcatt tgaatttgaa ttaacttaga aattcatttc ctcacctgta 1920
gtggccacct ctatattgag gtgctcaata agcaaaagtg gtcggtggct gctgtattgg 1980
acagcacaga aaaagatttc catcaccaca gaaaggtcgg ctggcagcac tggccaaggt 2040
gatggggtgt gctacacagt gtatgtcact gtgtagtggg tggagtttac tgtttggga 2100
ataaaaacgg ctgtttccgt gaaaanaaaa aaaaaaagg 2139

```

<210> 62

<211> 648

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 7500929CB1

<400> 62

```

gtggaagaga cgcgccggcg ttggccgctg ctgctagcag cttgaacccc agggtcggga 60
ccgatgtcgg cttgggctgc tgccagccta agcagggccg ctgcccgatg cttgctggca 120
cgaggccccg gggtcagggc ggctcctccg cgcgaccccc ggccctccca ccccgagccc 180
cggggctgcg gtgccgctcc gggcaggacg ctgcacttta ccgcggtgt ccccgccggg 240
cacaacaagt ggtccaaagt caggcacatc aagggtccga aggacgtcga aaggagtgcg 300
atcttctcca aactctgttt gaacatccgc ctggcagtga aagccaggag gccaaggac 360
aggacttgcg accttgaagc caaaggaatc tcacttggtg ggctccttg tcagctctgc 420
tgctgtctca gagccatctg gatgagtgtc ccgacaccct ctgggatgca gggcaggacc 480
accagctgg tcagactctg atgttgggta gctggcctct gtggggattg taagtgccct 540
gaggcgctct gtactagaaa ctgctcttaa taataacggt gattattggg tgetgcaaaa 600
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa 648

```